

Direct Explorations of Exoplanets with the Subaru Telescope and Beyond

Motohide Tamura¹, Lyu Abe¹
and Subaru HiCIAO/SPICA coronagraph/JTPF teams

¹Exoplanet Project Office, National Astronomical Observatory,
Osawa 2-21-1, Mitaka, Tokyo 181-8588, Japan
email: hide@subaru.naoj.org

Abstract. We present a Japanese “roadmap” on direct extrasolar planet studies spanning from the current ground-based telescope to future IR/Opt space-based telescopes. Several searches for young planets and disks have been conducted with the Subaru 8.2m telescope with adaptive optics (AO) infrared coronagraph, CIAO. The instrument will be soon upgraded to a new AO and a coronagraph with simultaneous spectral and polarimetric differential imaging modes (HiCIAO), which will significantly improve the contrast performance and hence the capability of young planet detection. A sensitive unbiased survey for extrasolar zodiacal emission around nearby stars will be conducted with the ASTRO-F space mission (0.7m telescope, $\sim 2\text{--}200\ \mu\text{m}$) to be launched around the beginning of 2006. A successor space mission, SPICA (3.5m, $5\text{--}200\ \mu\text{m}$), is also planned; its high sensitivity will enable the detection and characterization of outer-most planets around nearby stars, if any. For the studies of extrasolar terrestrial planets, a high contrast space telescope (HCST; 3.5m, $\sim 0.3\text{--}2\ \mu\text{m}$). We are also seeking for collaborations with or are considering to join to foreign missions. We describe an outline, status, and role of each project on the extrasolar planet studies.

Keywords. instrumentation: high angular resolution, planetary systems: formation, planetary systems: protoplanetary disks, stars: low-mass, brown dwarfs.

1. A roadmap

Figure 1 is a roadmap showing our approach to the explorations of exoplanets in Japan.

Our ground-based direct explorations are concentrated on the fully-operated and successful Subaru 8.2m telescope at Mauna Kea, Hawaii. Making use of the good observing site and the excellent image quality, the infrared coronagraph CIAO (Coronagraphic Imager with Adaptive Optics) has been used for various kinds of surveys, which is described in detail in the next section. Its description and scientific results are introduced in §2. Several indirect efforts are also sought on the Subaru telescope: an international collaboration on the N2K survey on the Doppler searches has been most productive (Sato *et al.* 2005). HiCIAO (Next generation High Contrast Imager with Adaptive Optics for the Subaru telescope) is the Subaru next generation coronagraph which is used with the new AO and its near-future upgrade to an extreme-AO system. This will be used for the more extensive surveys of young exoplanets, as described in §3. Regarding space-based explorations, the MIR and FIR space telescope, *ASTRO-F*, is developed by JAXA/ISAS and will be launched soon (Murakami 2000). This telescope has a similar size to the *Spitzer* telescope, but is optimized to all-sky survey observations. Although its low spatial resolution is not suitable for direct explorations, it will conduct an unbiased survey for faint disk emission. This enables us to make census of exo-zodi and disk evolution with a large sample of stars. As a next project in the OPT/IR community in Japan after Subaru/ASTRO-F, SPICA is currently heavily discussed, a cooled large single mirror




Figure 1. A roadmap of exoplanet explorations in Japan.

space-telescope. Because of its high sensitivity and acceptable resolution, the mission will serve for direct exoplanet explorations. Its coronagraph and exoplanet science are discussed in §4. A space-telescope dedicated for direct explorations of Earth-like planets is also under discussion in the OPT/IR community in Japan, as described in §5.

2. Subaru CIAO – First dedicated infrared coronagraph on 8–10 m class telescopes

Observationally, the direct imaging of both exoplanets and protoplanetary disks is extremely difficult: it requires a high sensitivity, a high resolution, and a high dynamic range or contrast at the same time. Such high sensitivity and high resolution are now simultaneously realized with the 8–10 meter class telescopes with adaptive optics (AO). The AO corrects for atmospheric perturbations in real-time, by the means of wavefront sensors and deformable mirrors. In the current Subaru AO system, 36-elements sensors and 36-elements mirrors are used. With the Subaru adaptive optics and infrared camera we regularly achieve a resolution of 0.07–0.10 arcsec and a sensitivity of 22 mag (5σ in 1 hour) around 1.5–2 μm (H and K bands). However, a high contrast, that is, a capability to observe faint objects near bright objects, is another thing.

CIAO is the first among those equipped with a full cold coronagraph mode (with various cold occulting masks and Lyot stop optics) on the world 8-m class telescopes. Compared to previous coronagraphs, it has unique features of near-infrared operation and small occulting masks (down to 0.2 arcsec in diameter). See Tamura *et al.* (1998, 2000) for the instrument details.

CIAO has a comparable or higher contrast than other coronagraphs on the 8-m class telescopes (Table 1). See Nakajima *et al.* (2005) for a detailed performance discussion of CIAO.

Table 1. Contrast comparison (from Nakajima *et al.* 2005)

Telescope/Instrument	Contrast	Separation
Subaru/CIAO	$\Delta K=13$	$r>2.5''$
Subaru/CIAO	$\Delta K=11$	$r=1.2''$
VLT/AO+Mask	$\Delta K=13$	$r>3.0''$
VLT/AO+Mask	$\Delta K=10$	$r=1.0''$
VLT/AO+SDI	$\Delta H=9.5$	$r=0.5''$

2.1. CIAO disk searches

CIAO has been successful to *directly* reveal the morphological diversity of protoplanetary disks, such as rings, spirals, and even others, with a high spatial resolution of $\sim 0.1''$. As shown in an inset of Figure 1, besides a well known ring-like disk around GG Tau, a clear spiral disk with several arms has been revealed around AB Aur, and a new-type of disk morphology, banana-split disk, has been recently discovered around HD 142527 (Fukagawa *et al.* 2004 and 2005). In addition, first $2\text{-}\mu\text{m}$ imaging polarimetry of β Pic has demonstrated that the disk is composed of several planetesimal belts with a gap at $r \sim 100$ AU (Tamura *et al.* 2005).

2.2. CIAO companion searches

Companion searches are greatly merited if the targets are younger; the younger companion is brighter and the contrast between the primary and the companion becomes lower (e.g., Burrows *et al.* 2001). This is one of the reasons why we explore YSO companions in nearby star forming regions.

DH Tau is a classical T Tauri star with an age of 0.1-4 Myr and a mass of 0.25-0.5 solar mass. CIAO coronagraphic images have revealed a companion of 15 mag at H at $2.3''$ (330 AU) from the central star. This star is originally selected because it has a relatively large infrared polarization (Tamura & Sato 1989), suggestive of the circumstellar structures. However, interestingly, no significant circumstellar disks or envelopes are detected except for the companion.

The companion shows the same proper motion of the Taurus cloud members, distinct from the background stars. Its near-infrared spectra show deep absorption band due to water and other metal features, all indicating low effective temperatures, but not as low as those of planetary mass objects ($T_{eff} \sim 2700$ K). With recent theoretical models, the companion is concluded as a companion young brown dwarf of about $40 M_{JUP}$ (Itoh *et al.* 2005). This source is more or less similar to the recently detected companion around GQ Lup and other handful companions around YSOs (Neuhauser *et al.* 2005; Liu 2005 in this proceedings).

Similar surveys are also nearby M stars using CIAO (Nakajima *et al.* 2005).

3. HiCIAO – the Subaru next high contrast instrument

HiCIAO is a new high-contrast instrument for the Subaru telescope, which is currently developed by NAOJ and University of Hawaii. NAOJ PI is M. Tamura and UH PI is K. Hodapp. HiCIAO will be used in conjunction with the new adaptive optics system (188 actuators and/or its laser guide star - AO188/LGSAO188) at the Infrared Nasmyth platform. It is designed as a flexible camera comprising several modules that can be configured into different modes of operation. The main modules are the AO module with its future extreme AO capability, the warm coronagraph module, the high contrast optics module, and the cold infrared camera module. HiCIAO will be the first instrument on

the 8-m class telescopes which can combine coronagraphic techniques with simultaneous polarization and spectral differential imaging modes which minimizes the common path errors. The basic concept of such differential imaging is to split up the image into two or more images, and then use either different planes of polarization or different spectral filter bandpasses to produce a signal that distinguishes faint objects near a bright central object from scattered halo or residual speckles. This enables us to achieve a contrast improvement of at least 10 times better than before. The instrument specification is summarized in Table 2.

Table 2. HiCIAO overall specifications.

Platform	Subaru Nasmyth focus (IR)
Combined adaptive optics	188-elements curvature-sensing AO
Wavelength coverage	<i>zJHKs</i> (optimized at H)
Detector	2048 × 2048 HgCdTe (Rockwell HAWAII-2RG)
Detector controller	ASIC (Rockwell “Sidecar”)
Pixel scale	0.01"/pixel (proper PSF sampling at J-band)
Spatial resolution	0.03" (<i>J</i>), 0.04" (<i>H</i>), 0.055" (<i>Ks</i>) FWHM
Strehl ratio	0.6 (<i>J</i>), 0.7 (<i>H</i>), 0.8 (<i>Ks</i>) w/ 0.5" optical seeing
Sensitivity	<i>H</i> > 23 mag (5 σ , 1 hr)
Contrast goal	10 ⁴ at 0.1", 10 ⁶ at 1" (w/ SDI mode + coronagraph)
DI mode	Direct imaging 20" × 20" FOV, general purpose imaging High throughput, w/wo coronagraph
PDI mode	Polarimetric (o-ray and e-ray) dual imaging (Wollaston prism) 20" × 10" FOV, polarized target imaging
SDI mode	Spectral simultaneous differential imaging (Double Wollaston prism) 6" × 6" FOV, unpolarized target imaging
Coronagraph*	
Occulting masks	3 sizes with real-time remote choice
Lyot stops	Stop with spider/center-hole blocks & rotation compensation

* Starting with the classical Lyot coronagraph. More sophisticated coronagraph will be implemented later.

The main sciences with Subaru/HiCIAO are summarized as follows.

- For directly detecting young (*age* ~ 100 Myr) extrasolar planets and brown dwarfs, HiCIAO can be used to conduct a survey of ~120 nearby relatively young dwarfs. About 60 stars younger than 700 Myr with spectral types between A and K are accessible from Subaru. These 60 stars and the 350 Myr M dwarf sample (~60 stars) will serve as the prime targets for the brown dwarf/planet. The inner radius of search will be reduced from 25 AU to 10 AU for a target at 10 pc by the transition from CIAO to HiCIAO.

- For directly detecting very young (*age* ~ 1–10 Myr) extrasolar planets and brown dwarfs, HiCIAO can be used to conduct a survey of ~200 T Tauri stars in nearest star forming regions accessible from the northern hemisphere and to detect young Jupiter-mass companions down to an inner radius of 30 AU. With the availability of the laser guide star, the target can be extended to the optically faint sources including Class I YSOs.

- HiCIAO will be efficiently used to detect or set a limit on scattered light from disks around the young stars in nearby star forming regions. The spatial resolution is 0.03" (80 AU at the distance of the Taurus dark cloud) in the *J*-band, which is better than the

HST resolution. The large sample of disk data will be used to trace the morphological evolution and diversity of YSO disks at the age of age <1 Myr to 10 Myr.

- HiCIAO will be extremely useful to detect or set a limit on scattered light from the known and new, large samples of Vega-like disks from the *Spitzer* and the *ASTRO-F* missions, which requires both high sensitivity and high contrast. The achievable spatial resolution is 0.6 AU at the distance of β Pic. The polarization differential imaging is crucial to study the small-scale and inner disk structures and dust grain properties. A synergy observation with the *ASTRO-F* will be considered.

- The targets for TPF are well pre-surveyed with HiCIAO for characterizing their circumstellar matter and detecting relatively “low”-contrast objects including background stars and galaxies.

An extensive survey dedicated to these sciences on the Subaru telescope will be a significant contribution to astronomy before the high contrast space telescope era. The application to other targets might include not only other galactic objects such as late-type stars but also extragalactic objects such as AGNs and quasars, especially with the laser guide stars, where the AO application has been relatively limited. A high throughput direct imaging will be useful for such objects.

4. SPICA coronagraph

The SPICA mission plans to launch a cooled (4.5K) large (3.5m) single-mirror telescope into a halo orbit around the Sun-Earth L2 orbit around the beginning of 2010s (Nakagawa 2000). Its unprecedented sensitivity at MIR (and FIR) and its simple telescope pupil make SPICA to be one of the best platforms to implement coronagraph instrument for exoplanet studies. The preliminary specification of the SPICA coronagraph is summarized in Table 3.

Table 3. SPICA coronagraph preliminary specification.

Wavelength coverage	5–27 μm
Spectral resolution	<200
Inner working distance	$3\lambda/D$
Contrast goal	10^6
Detector	1Kx1K Si:AS IBC
Adaptive Optics	Cold Tip-Tilt mirror (to correct for the jitters) Cold Deformable Mirror (to correct for the mirror wavefront errors)

SPICA will target direct observations of self-luminous planets at $r >$ a few to ~ 20 AU of nearby (<10 pc) stars. The detectable planets depend on their mass, age, and separation. If we assume the inner working distance of $3\lambda/D$, then at $\lambda=5$ μm we can detect 1 Gyr, 2 M_{JUP} planets at $r > 9$ AU around 10 pc stars (~ 30 G-M type stars). At $\lambda=20$ μm we can detect 5 Gyr, 2 M_{JUP} planets at $r > 18$ AU around 10 pc stars (~ 150 G-M type stars). Very young planets and sub-brown dwarfs in nearby star forming regions and cold brown dwarfs are also good targets.

5. JTPF

JTPF (Japanese Terrestrial Planet Finder) is a future space mission whose main science driver is the extrasolar terrestrial planet studies. The project is officially approved both by NAOJ and ISAS. In NAOJ, its activity is regarded as one of the main themes of the Phase A project which has started in April 2005. In JAXA/ISAS, the JTPF Working Group

has been approved by the ISAS Science Steering Committee in 2002. Discussion has been made among both astronomy and planetary communities in Japan, and regarded as one of the important programs for optical and infrared astronomy. It could be either an independent mission or a joining collaboration to TPF/Darwin missions.

At present, an optical high contrast space telescope (HCST) which realizes clean and stable images is under main discussions for the JTPF architecture among JTPF WG (see Figure 1). JTPF/HCST will make full use of the SPICA bus system with an optimization to shorter wavelengths. The current plan of HCST is shown in Table 4.

Table 4. JTPF/HCST preliminary specification.

Telescope	3.5m off-axis single aperture
Wavelength coverage	Optical and near-IR w/ limited UV capability
Orbit	Sun-Earth L2
Instruments	Opt-NIR coronagraph, Wide-field NIR/Opt instruments

Various JTPF-related activities in Japan include studies of:

- combinations of nulling interferometer and modified pupil (Nishikawa *et al.* 2005),
- a chromatic nulling of on-axial light to detect off-axial light by means of 3D common-path interferometer (Tavrov *et al.* 2002), and
- polarization differential objective spectroscopy with nulling coronagraph (Murakami *et al.* 2005).

References

- Burrows, A., Hubbard, W. B., Lunine, J. I., & Liebert, J. 2001, *Rev. of Modern Physics* 73, 719
- Fukagawa, M., Hayashi, M., Tamura, M. *et al.* 2004, *ApJ* 605, L53
- Fukagawa, M., Tamura, M., Hayashi, M. *et al.* 2005, *ApJ* submitted
- Itoh, Y., Hayashi, M., Tamura, M. *et al.* 2005, *ApJ* 620, 984
- Murakami, H. 2000, *ISAS Report SP* 14, 267
- Murakami, N., Baba, N., Tate, Y. *et al.* 2005, *PASP* submitted
- Nakagawa, T. 2000, *ISAS Report SP* 14, 189
- Nakajima, T., Morino, J., Tsuji, T. *et al.* 2005, *Astronomical Notes* in press
- Neuhauser, R., Guenther, E. W., Wuchterl, G., Mugrauer, M., Bedalov, A., Hauschildt, P. H. 2005, *A&A* 435, L13
- Nishikawa, J., Kotani, T., Murakami, N., Baba, N., Itoh, Y., & Tamura, M. 2005, *A&A* 435, 379
- Sato, B. *et al.* 2005, *ApJ* in press
- Tamura, M., Suto, H., Takami, H. *et al.* 1998, *SPIE* 3354, 845
- Tamura, M., Suto, H., Itoh, Y. *et al.* 2000, *SPIE* 4008, 1153
- Tamura, M., Fukagawa, M., Kimura, H. *et al.* 2005, *ApJ* submitted
- Tamura, M. & Sato, S. 1989, *AJ* 98, 1368
- Tavrov, A. *et al.* 2002, *Opt. Lett.* 27, 2070