

SMOOTHNESS OF SOLUTIONS OF PARABOLIC EQUATIONS IN REGIONS WITH EDGES

A. AZZAM AND E. KREYSZIG*

§1. Introduction

We consider the mixed initial-boundary value problem for the parabolic equation

$$(1.1) \quad Lu = \sum_{i,j=1}^2 a_{ij}(x, t)u_{x_i x_j} + \sum_{j=1}^2 a_j(x, t)u_{x_j} + b(x, t)u - u_t = f(x, t)$$

in a region $\Omega \times (0, T]$, where $x = (x_1, x_2)$ and $\Omega \subset \mathbb{R}^2$ is a simply-connected bounded domain having corners.

Our main objective will be the study of smoothness properties of solutions of that problem. Early investigations of that type concern *elliptic* equations in domains with a *smooth* boundary, starting with the Dirichlet problem for the Laplace and Poisson equations and proceeding to general second order elliptic equations as well as general boundary conditions; see S. Agmon, A. Douglis and L. Nirenberg [1]. A more recent survey of the elliptic case and further references are given by D. Gilbarg and N. S. Trudinger [6].

Similar work on *parabolic* equations appeared later; we mention in particular investigations by A. Friedman [5] on the first boundary value problem, by Z. Itô [8] and L. I. Kamynin and V. N. Maslennikova [9] on the second boundary value problem and by N. V. Zitarasû [15] on general boundary value problems. Further references are given in [12].

The case of a *nonsmooth boundary* was treated by E. A. Volkov [14], V. A. Kondrat'ev [10] and others whose work is discussed or mentioned in [7]; all these papers concern *elliptic* equations, whereas we shall deal with *parabolic* equations.

We want to mention that those problems in regions with edges and

Received December 21, 1979.

* Work supported by the N.S.E.R.C. of Canada under grant no. A9097.

corners are also of practical importance in applications, for instance in heat flow (cf. [4]), mechanics of continua and numerical analysis (difference methods, subtraction of singularities, acceleration of convergence; cf. [11], which treats elliptic equations but involves ideas which are also relevant in connection with parabolic equations).

It is known that in the case of a *smooth boundary* $\partial\Omega$ of Ω , the smoothness of solutions increases with that of the coefficients of (1.1) and the boundary data. Indeed, if $\partial\Omega$ is of class $C^{2+\alpha}$, $0 < \alpha < 1$, the coefficients of (1.1) are of class $C^\alpha(\bar{G})$, $G = \Omega \times (0, T]$, and u is a solution of (1.1) in G satisfying

$$\begin{aligned} u(x, 0) &= 0 && \text{on } \bar{\Omega} \\ \beta(x, t)u + \eta(x, t)u_n &= 0 && \text{on } \partial\Omega \times (0, T], \end{aligned}$$

where u_n is the outer normal derivative and $\beta \in C^{2+\alpha}(\partial\Omega \times (0, T])$, $\eta \in C^{1+\alpha}(\partial\Omega \times (0, T])$, then $u \in C^{2+\alpha}(\bar{G})$.

In this paper we show that the increase of the smoothness of solutions with that of the coefficients of (1.1) and the boundary data is no longer true if $\partial\Omega$ has corners, and obtain a smoothness theorem for this case.

§2. Main result

Let $\Omega \subset \mathbb{R}^2$ be a simply connected bounded domain with boundary $\partial\Omega$. For simplicity and without loss of generality we assume that $\partial\Omega$ has a single corner at the origin with interior angle $\gamma > 0$. We assume that the two arcs Γ_1 and Γ_2 forming the corner have the representation

$$\begin{aligned} \Gamma_1: & x_1 = g_2(x_2) \\ \Gamma_2: & x_2 = g_1(x_1) \end{aligned}$$

with g_1 and g_2 of class $C^{2+\alpha}$ and $g_1(0) = 0$, $g_2(0) = 0$, $g_1'(0) = 0$, and $g_2'(0) = \cot \gamma$.

We consider the problem [cf. (1.1)]

$$(2.1) \quad Lu = \sum_{i,j=1}^2 a_{ij}(x, t)u_{x_i x_j} + \sum_{j=1}^2 a_j(x, t)u_{x_j} + b(x, t)u - u_t = f(x, t) \quad \text{in } G,$$

$$(2.2a) \quad u(x, 0) = 0 \quad \text{on } \bar{\Omega},$$

$$(2.2b) \quad \beta(x, t)u + \eta(x, t)u_n = 0 \quad \text{on } S = (\partial\Omega \setminus \{0\}) \times (0, T],$$

where $\beta \in C^{2+\alpha}(S)$, $\eta \in C^{1+\alpha}(S)$, and

$$\beta = 1, \eta = 0 \text{ on } \Gamma_1, \quad \beta = 0, \eta = 1 \text{ on } \Gamma_2 .$$

If $a_{ij}, a_j, b, f \in C^\alpha(\bar{G})$, then for any bounded solution of (2.1)–(2.2) we have

$$(2.3) \quad u \in C^{2+\alpha}(G_1) \cap C^0(\bar{G})$$

where $G_1 = \Omega_1 \times (0, T]$ and Ω_1 is any compact subregion of $\bar{\Omega}$ having positive distance from 0.

To investigate the smoothness of solutions near the edge, we take any fixed $t_0 \in [0, T]$ and transform the equation

$$(2.4) \quad \sum_{i,j=1}^2 a_{ij}(0, t_0) u_{x_i x_j} = 0$$

to canonical form. Then γ at $(0, t_0)$ is transformed to

$$\omega(t_0) = \arctan \frac{[a_{11}(0, t_0)a_{22}(0, t_0) - a_{12}^2(0, t_0)]^{1/2}}{a_{22}(0, t_0) \cot \gamma - a_{12}(0, t_0)} .$$

Clearly, the value $\omega(t_0)$ does not depend on the particular choice of that transformation.

We now state our main result. From (2.3) it follows that it suffices to consider the smoothness of solutions near the edge.

THEOREM 1. *Let u be a bounded solution of (2.1), (2.2) in $\mathcal{G}_c = \Omega_c^* \times (0, T]$, where*

$$\Omega_c^* = \{x | x \in \Omega, 0 < |x| \leq c\} .$$

Assume $a_{ij}, a_j, b, f \in C^\alpha(\bar{\mathcal{G}}_c)$ in (2.1), where $0 < \alpha < 1$, and, furthermore, $\omega(t) < \frac{1}{2}\pi$ for every $t \in [0, T]$. Then u , as a function of x , satisfies

$$(2.5) \quad u \in C^\nu(\bar{\mathcal{G}}_c) ,$$

for some $\nu \in (1, 2]$.

The proof of this theorem will result from Theorem 2 (below) and will be given at the end.

§3. The case of a cylindrical sector

Let $t_0 \in [0, T]$ be fixed and define

$$G_s = \Omega_s \times I_s ,$$

where

$$\Omega_\sigma = \{(r, \theta) | 0 < r \leq \sigma, \frac{1}{2}\pi - \omega < \theta < \frac{1}{2}\pi\}$$

with $\omega = \omega(t_0) < \frac{1}{2}\pi$ and r, θ given by $x_1 = r \cos \theta, x_2 = r \sin \theta$, and

$$I_\sigma = \{t | t \in (0, T], |t - t_0| \leq \sigma\}.$$

Furthermore, let

$$\Gamma_1 = \{(r, \theta) | r \leq \sigma, \theta = \frac{1}{2}\pi - \omega\},$$

$$\Gamma_2 = \{(r, \theta) | r \leq \sigma, \theta = \frac{1}{2}\pi\}.$$

As the mixed initial-boundary conditions we take

$$(3.1a) \quad u(x, 0) = 0 \quad \text{on } \Omega_\sigma,$$

$$(3.1b) \quad u|_{\Gamma_1} = u_n|_{\Gamma_2} = 0, \quad 0 < t \leq T.$$

Then we have the following result.

THEOREM 2. *Let u be a bounded solution of the problem (2.1), (3.1) in G_σ with $\omega < \frac{1}{2}\pi$. Suppose that $a_i, a_j, b, f \in C^\alpha(\bar{G}_\sigma), 0 < \alpha < 1$, and $a_{ij}(0, t_0) = \delta_{ij}$. Then u , as a function of x , satisfies*

$$(3.2) \quad u \in C^\nu(\bar{G}_\sigma),$$

where $\nu < \sigma, \nu = \min(2, \pi/2\omega - \varepsilon)$ and $\varepsilon > 0$ is arbitrarily small.

We shall obtain a proof of this theorem by first proving two lemmas in the next two sections.

§4. Estimation of solutions

LEMMA 1. *Under the assumptions of Theorem 2 there exists a positive $a < \sigma$ such that in \bar{G}_a (cf. Sec. 3)*

$$(4.1) \quad |u(x, t)| \leq Mr^\nu,$$

where $\nu = \min(2, \pi/2\omega - \varepsilon)$ and $\varepsilon > 0$ is arbitrarily small.

Proof. In G_σ we consider the function

$$v(x) = -Mr^\nu \cos \lambda(\frac{1}{2}\pi - \theta),$$

$$\nu = \min\left(2, \frac{\pi}{2\omega} - \varepsilon\right) < \frac{\pi - 2\delta}{2\omega} = \lambda, \quad 0 < \delta < \frac{1}{2}\pi.$$

We shall prove that $v(x)$ serves as a barrier for the solution u of (2.1), (3.1). We have

$$Lv = M[(\lambda^2 - \nu^2) \cos \lambda(\frac{1}{2}\pi - \theta) - h(x, t)]r^{\nu-2} + Mh_1(x, t)r^{\nu-1} + Mh_2(x, t)r^\nu,$$

where $h(x, t)$ is continuous, $h(0, t_0) = 0$, and h_1 and h_2 are bounded. Thus for every $\epsilon_1 > 0$ we can find an $a < \sigma$ such that in G_a ,

$$Lv \geq M[(\lambda^2 - \nu^2) \cos \lambda(\frac{1}{2}\pi - \theta) - \epsilon_1]r^{\nu-2}.$$

Since for $\frac{1}{2}\pi - \omega \leq \theta \leq \frac{1}{2}\pi$ we have

$$\sin \delta \leq \cos \lambda(\frac{1}{2}\pi - \theta) \leq 1,$$

in G_a we thus obtain

$$Lv \geq M[(\lambda^2 - \nu^2) \sin \delta - \epsilon_1]r^{\nu-2}$$

with arbitrarily small $\epsilon_1 > 0$ and sufficiently small $a < \sigma$. We now take $\epsilon_1 < (\lambda^2 - \nu^2) \sin \delta$ and use the fact that $f(x, t)$ is bounded in G_a , say, $|f(x, t)| \leq K$. Consequently, by taking M sufficiently large (if $\nu = 2$) and a sufficiently small (if $\nu < 2$), we obtain in G_a

$$Lv \geq K \geq f(x, t).$$

We now want to apply in that region the maximum principle to $w = u - v$. We have $Lw \leq 0$ in G_a . Furthermore, $w_n = 0$ on $\Gamma_2 \times I_a$ and $w \geq 0$ on $\Gamma_1 \times I_a$. Since u is bounded in G_a , by taking M sufficiently large we can make $w \geq 0$ on

$$\{(r, \theta) | r = a, \frac{1}{2}\pi - \omega < \theta < \frac{1}{2}\pi\} \times I_a.$$

Hence, by the maximum principle (cf. [13]) we have $w \geq 0$ in \bar{G}_a with sufficiently small a ; that is,

$$u \geq -Mr^\nu \cos \lambda(\frac{1}{2}\pi - \theta) \geq -Mr^\nu.$$

The other part of (4.1) can be obtained similarly. This proves Lemma 1.

§5. Proof of Theorem 2

To prove Theorem 2, we need another lemma, as follows.

LEMMA 2. *Let u be a bounded solution of (2.1), (2.2) in G_a . Suppose that $a_{i,j}, a_j, b, f \in C^\alpha(\bar{G}_a)$. Then if for some $\mu, 0 < \mu \leq 2$, and $a < \sigma$ we have*

$$|u(x, t)| \leq Mr^\mu \quad \text{in } \bar{G}_a,$$

it follows that

$$(5.1) \quad \left| \frac{\partial^k u(x, t)}{\partial x_1^{k_1} \partial x_2^{k-k_1}} \right| \leq M_k r^{\mu-k} \quad \text{in } \bar{G}_a$$

where $k_1 = 0, 1$ or $2, k_1 \leq k, k = 1, 2$.

Proof. It suffices to indicate the basic idea since similar proofs were used in [2] and [3]. In Ω_a we define

$$D_s = \{(r, \theta) | 2^{-s-2}a \leq r \leq 2^{-s-1}a, \frac{1}{2}\pi - \omega < \theta < \frac{1}{2}\pi\},$$

$$D'_s = D_{s-1} \cup D_s \cup D_{s+1}, \quad s = 0, 1, \dots,$$

and set

$$R_s = D_s \times I_a \quad R'_s = D'_s \times I_a.$$

The transformation

$$x_i = 2^{-s}y_i, \quad i = 1, 2,$$

transforms (2.1), (2.2) into

$$\sum \tilde{a}_{ij} \tilde{u}_{y_i y_j} + 2^{-s} \sum \tilde{a}_j \tilde{u}_{y_j} + 2^{-2s} \tilde{b} \tilde{u} - 2^{-2s} \tilde{u}_t = 2^{-2s} \tilde{f}, \quad \tilde{\beta} \tilde{u} + \tilde{\eta} \tilde{u}_n = 0.$$

Furthermore, it maps R_s onto R_0 and R'_s onto R'_0 . In R_0 and R'_0 we apply Schauder estimates for the solution of the transformed problem to obtain

$$\|\tilde{u}\|_{2+\alpha}^{R_0} \leq C[\|\tilde{u}\|_{R'_0} + 2^{-2s} \|\tilde{f}\|_{R'_0}].$$

This yields in R_s

$$\left| \frac{\partial^k u(x, t)}{\partial x_1^{k_1} \partial x_2^{k-k_1}} \right| \leq c_0 2^{-(\mu-k)s},$$

which entails the assertion of Lemma 2.

From this lemma (with $\mu = \nu$) we shall now obtain Theorem 2.

Proof of Theorem 2. It is sufficient to show that

$$(5.2) \quad \frac{|u_x(P) - u_x(Q)|}{d(P, Q)^{\nu-1}} \leq H$$

for any two points $P: (r_1, \theta_1, t_1)$ and $Q: (r_2, \theta_2, t_2)$ in \bar{G}_a . Here

$$d(P, Q)^2 = (x_1 - x_2)^2 + (y_1 - y_2)^2 + |t_1 - t_2|,$$

as usual. Without restriction, let $0 \leq r_2 \leq r_1$. If $r_2 \leq r_1/2$ or $|t_1 - t_2|^{1/2} \geq r_1/2$, then $d(P, Q) \geq r_1/2$ and (5.2) can be obtained using the bound (5.1) of u_x . Now let $r_2 > r_1/2$ and consider the region

$$S_P = \{(x, t) | (x, t) \in G_a, \frac{1}{2}r_1 \leq r \leq r_1, |t - t_1| \leq \frac{1}{4}r_1^2\}.$$

The transformation

$$x_i = 2r_1 z_i / a, \quad i = 1, 2,$$

maps S_P onto

$$G'_P = \{(z, t) | \frac{1}{4}a \leq \rho \leq \frac{1}{2}a, |t - t_1| \leq \frac{1}{4}r_1^2\},$$

where $\rho^2 = z_1^2 + z_2^2$. It transforms equation (2.1) to

$$\sum a_{ij}^* u_{z_i z_j}^* + \frac{2r_1}{a} \sum a_j^* u_j^* + \left(\frac{2r_1}{a}\right)^2 b^* u^* - \left(\frac{2r_1}{a}\right)^2 u_t^* = \left(\frac{2r_1}{a}\right)^2 f^*$$

In G'_P and

$$G''_P = \{(z, t) | \frac{1}{8}a \leq \rho \leq a, |t - t_1| \leq \frac{1}{4}r_1^2\}$$

we apply a Schauder estimate to get

$$\|u^*\|_{2+\alpha}^{G'_P} \leq C \left[\|u^*\|_{\alpha}^{G''_P} + \left(\frac{2r_1}{a}\right)^2 \|f^*\|_{\alpha}^{G''_P} \right].$$

As in the proof of Lemma 2 we obtain

$$(5.3) \quad \|u^*\|_{2+\alpha}^{G'_P} \leq M_2 r_1^\nu.$$

Now

$$H_{\nu-1}^{G_P}(u_x) = \left(\frac{2r_1}{a}\right)^\nu H_{\nu-1}^{G'_P}(u_x^*),$$

where $H_{\nu-1}^{G'_P}(u_x^*)$ is the Hölder coefficient of the Hölder condition for u_x in G'_P (with exponent $\nu - 1$). From (5.3) we thus obtain

$$H_{\nu-1}^{G_P}(u_x) \leq M_3.$$

This proves Theorem 2.

§ 6. Proof of Theorem 1. Concluding remarks

Consider any point $(0, t_0)$, $t_0 \in [0, T]$. We straighten the boundary around $(0, t_0)$ by the transformation

$$(6.1) \quad y_1 = x_1 - g_2(x_2), \quad y_2 = x_2 - g_1(x_1).$$

Equation (2.1) is then transformed to a parabolic equation with principal part

$$\sum_{i,j=1}^2 A_{ij}(y, t)U_{y_i y_j} .$$

Clearly, (6.1) is of class $C^{2+\alpha}$ and the value of the Jacobian at $(0, t_0)$ is 1. By another linear transformation we cast

$$\sum_{i,j=1}^2 A_{ij}(0, t_0)U_{y_i y_j} = 0$$

to canonical form. These two transformations map Ω_c^* in Theorem 1 onto a sector Ω_σ of angle $\omega(t_0)$. In $G_\sigma = \Omega_\sigma \times (0, T]$ with a suitable $\sigma > 0$ the transformed functions satisfy all the conditions of Theorem 2. Hence the conclusion of Theorem 2 applies to the transformed solution. Since the composite of those transformations is of class $C^{2+\alpha}$ and locally injective, Theorem 1 follows.

Remark 1. In [3] we were concerned with the initial Dirichlet problem for a parabolic equation with coefficients a_{ij} independent of t . Our present results now permit us to extend our considerations in [3] for the case $n = 2$ as follows.

THEOREM 3. *Let u be a bounded solution of the initial-Dirichlet problem*

$$\begin{aligned} Lu &= f && \text{in } G_c \\ u(x, 0) &= 0 && \text{on } \bar{\Omega} \\ u &= 0 && \text{on } S \end{aligned}$$

with L , Ω and S as in Sec. 2. Suppose that a_{ij} , a_j , b , $f \in C^\alpha(\bar{G}_c)$ and $\omega(t) < \pi$ for every $t \in [0, T]$. Then

$$(6.2) \quad u \in C^\nu(\bar{G}_c)$$

for some $\nu \in (1, 2]$.

To prove this, find a bound for u , using the idea of the proof of the above Lemma 1 and then continue as in [3].

Remark 2. If u and the coefficients of the equation in Theorem 2 are extended by symmetry across $\theta = \frac{1}{2}\pi$, then in $\bar{G}_c \setminus J$ with $\bar{G}_c = \bar{\Omega}_c \times I_c$ and

$$\begin{aligned} \bar{\Omega}_c &= \{(r, \theta) | 0 < r \leq c, \frac{1}{2}\pi - \omega < \theta < \frac{1}{2}\pi + \omega\} , \\ J &= \{(\theta, t) | \theta = \frac{1}{2}\pi, t \in I_c\} \end{aligned}$$

the extended function satisfies an initial *Dirichlet* problem (instead of a *mixed* problem), and we may still apply the maximum principle for gener-

alized solutions to obtain a bound for the solution. However, in Lemma 2 it is required that we have a bound for the solution *everywhere*. Hence the approach just mentioned would not be of help in the present case.

Remark 3. Combining Theorems 1 and 3 we obtain

THEOREM 4. *Let $\Omega \subset \mathbb{R}^2$ be a bounded domain whose boundary $\partial\Omega$ is a simple polygon, with sides $\Gamma_1 \dots, \Gamma_p$ of class $C^{2+\alpha}$ and vertices $P_j = \Gamma_j \cap \Gamma_{j+1}$, $j = 1, \dots, p$, ($\Gamma_{p+1} = \Gamma_1$) with angles γ_j such that for the corresponding $\omega_j = \omega(P_j, t)$ we have*

$$\begin{aligned} \omega_j &= \frac{1}{2}\xi_j(t)(\kappa_j + \kappa_{j+1}), & 0 < \xi_j < \pi, \\ \kappa_j &= 0 \text{ or } 1, & \kappa_j + \kappa_{j+1} \neq 0, \quad j = 1, \dots, p \quad (\kappa_{p+1} = \kappa_1). \end{aligned}$$

Let u be a bounded solution of the problem

$$\begin{aligned} Lu &= f \text{ in } \Omega \times I, \quad I = (0, T], \\ u(x, 0) &= 0 \quad \text{in } \bar{\Omega}, \\ \kappa_j u + (1 - \kappa_j)u_n &= 0 \quad \text{on } \Gamma_j \times I, \end{aligned}$$

with L as in (2.1) and $a_{ij}, a_j, b, f \in C^\alpha(\bar{\Omega} \times \bar{I})$, $0 < \alpha < 1$. Then

$$(6.3) \quad u \in C^{2+\alpha}(\bar{\Omega}_1 \times \bar{I}),$$

where $\bar{\Omega}_1$ is any compact subregion of $\bar{\Omega}$ with positive distance from the corners, and

$$(6.4) \quad \begin{aligned} u &\in C^{\nu_j}([\bar{\Omega} \cap N_j] \times \bar{I}) \\ \nu_j &= \min_{t \in I} (2, \pi/\xi_j(t) - \epsilon), \end{aligned}$$

where $\epsilon > 0$ is arbitrarily small and N_j is a sufficiently small closed disk centered at P_j .

The proof is obvious; indeed, for $\kappa_j + \kappa_{j+1} = 1$ (mixed data) it follows from Theorem 1 and for $\kappa_j + \kappa_{j+1} = 2$ (Dirichlet data) from Theorem 3.

Acknowledgement. We want to thank the Reviewer for valuable comments which entailed a substantial improvement of this paper.

REFERENCES

[1] S. Agmon, A. Douglis and L. Nirenberg, Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions, I., *Comm. Pure Appl. Math.*, **12** (1959), 623–727.

- [2] A. Azzam, On Dirichlet's problem for elliptic equations in sectionally smooth n -dimensional domains, *SIAM J. Math. Anal.*, **11** (1980), 248–253.
- [3] A. Azzam and E. Kreyszig, On parabolic equations in n space variables and their solutions in regions with edges, in press.
- [4] H. S. Carslaw and J. C. Jaeger, *Conduction of heat in solids*, Clarendon Press, Oxford (1959).
- [5] A. Friedman, Boundary estimates for second order parabolic equations and their applications, *J. Math. Mech.*, **7** (1958), 771–791.
- [6] D. Gilbarg and N. S. Trudinger, *Elliptic partial differential equations of second order*, Springer, Berlin etc., 1977.
- [7] P. Grisvard, Behavior of the solutions of an elliptic boundary value problem in a polygonal or polyhedral domain; in: *Numerical solutions of partial differential equations III* (ed. B. Hubbard), Academic Press, New York (1976).
- [8] Z. Itô, A boundary value problem of partial differential equations of parabolic type, *Duke Math. J.*, **24** (1957), 299–312.
- [9] L. I. Kamynin and V. N. Maslennikova, Boundary estimates for the solution of the third boundary-value problem for a parabolic equation, *Dokl. Akad. Nauk SSSR*, **153** (1963), 526–529.
- [10] V. A. Kondrat'ev, Singularities of a solution of Dirichlet's problem for a second order elliptic equation in the neighborhood of an edge, *Diff. Uravnen.*, **13** (1977), 2026–2033.
- [11] P. Laasonen, On the degree of convergence of discrete approximations for the solutions of the Dirichlet problem, *Ann. Acad. Fenn. Ser. A. I. No. 246* (1957), 19 pp.
- [12] O. A. Ladyženskaja, V. A. Solonnikov and N. Ural'ceva, *Linear and quasi-linear equations of parabolic type*, American Mathematical Society, Providence, R. I. 1968.
- [13] M. H. Protter and H. F. Weinberger, *Maximum principles in differential equations*, Prentice-Hall, Englewood Cliffs, N. J. (1967).
- [14] E. A. Volkov, Differentiability properties of solutions of boundary value problems for the Laplace equation on a polygon, *Proc. Steklov Inst. Math.*, **77** (1965), 127–159.
- [15] N. V. Zitarăşu, Schauder estimates and solvability of general boundary problems for general parabolic systems with discontinuous coefficients, *Soviet Math. Dokl.*, **7** (1966), 952–956.

*Department of Mathematics
University of Windsor
Windsor, Ontario, Canada*