

Relationship of socio-economic factors and parental eating habits with children's food intake in a population-based study in a metropolitan area of Brazil

Gabriela dos Santos Barroso¹, Rosely Sichieri² and Rosana Salles-Costa^{3,*}

¹Post-Graduate Program in Nutrition, Nutrition Institute Josué de Castro, Federal University of Rio de Janeiro, Rio de Janeiro/RJ and Antonio Pedro Hospital – Federal Fluminense University, Niterói/RJ, Brazil; ²Department of Epidemiology, Institute of Social Medicine, Universidade do Estado do Rio de Janeiro, Rio de Janeiro/RJ, Brazil; ³Department of Public Nutrition, Nutrition Institute Josué de Castro, Federal University of Rio de Janeiro, Av. Carlos Chagas Filho 373, Edifício do Centro de Ciências da Saúde, Bloco J, 2° andar, Cidade Universitária, Rio de Janeiro/RJ, CEP 21941-902, Brazil

Submitted 1 September 2011: Final revision received 5 September 2012: Accepted 11 September 2012: First published online 16 October 2012

Abstract

Objective: To evaluate the association of sociodemographic factors and parental food consumption with children's food intake.

Design: A cross-sectional survey.

Setting: A population-based study with a representative sample in a metropolitan region of Rio de Janeiro, Brazil. Parents' socio-economic variables, age and education level and children's age were obtained by face-to-face interviews. The parental food intake was assessed using an FFQ and the children's food intake was assessed using two 24 h recalls.

Subjects: Children (*n* 366) aged 6–30 months and their parents.

Results: The hierarchical regression analysis indicated that parents' age was positively associated with the intake of vegetables among children ($\beta = 0.73$, 95% CI 0.11, 1.34), while parents' educational level was positively associated with the intake of fats ($\beta = 3.52$, 95% CI 0.04, 7.01) and negatively associated with the intake of beans ($\beta = -13.98$, 95% CI -27.94 , -0.03). The age of the children was positively associated with the intakes of meats and eggs ($\beta = 2.88$, 95% CI 1.55, 4.22), sugars ($\beta = 5.08$, 95% CI 1.85, 8.30) and coffee ($\beta = 1.77$, 95% CI 0.71, 2.84), and negatively associated with the intake of vegetables ($\beta = -2.12$, 95% CI -3.20 , -1.05). The influence of parental food intake was observed for the food groups of breads, cereals and tubers ($\beta = 0.06$, 95% CI 0.003, 0.12), beans ($\beta = 0.11$, 95% CI -0.003 , 0.22) and fruits ($\beta = 0.10$, 95% CI 0.03, 0.16). Unfavourable socio-economic variables were associated with intakes of breads, cereals and tubers, vegetables, fruits, meats, sugars and coffee by children.

Conclusions: Parental food intake is associated with children's intake of cereals, beans and fruits independent of socio-economic status.

Keywords
Parental dietary intake
Children diet
Socio-economic factors
Food insecurity

The quality of health and nutrition in a population reflects its food intake. This is especially true for children, whose growth and development depend on nutrition as a fundamental condition⁽¹⁾. Inadequate feeding in the first 2 years of life, especially in disadvantaged populations, is closely associated with increased morbidity and may lead to problems such as Fe deficiency, malnutrition or obesity, and other nutritional disorders⁽²⁾.

Dietary intake can be influenced by cultural, social, demographic, economic and political factors in a society, and food habits and attitudes become consequences of these characteristics⁽³⁾. Therefore, studies have analysed the effect of social inequality on infant feeding in

terms of family characteristics such as low income⁽⁴⁾, family food insecurity^(5–8) and low levels of parental education^(9–11).

Family eating habits also influence children's food intake, as indicated in studies conducted primarily in developed countries^(12–14). However, the children's age when parental influence on children's food intake is most important has not been well established^(15,16). In addition, most of the previous studies have been conducted in middle-class families with almost no data on low-income families. Thus, the present cross-sectional study was conducted to identify the associations of socio-economic and demographic variables and parents' food intake with

*Corresponding author: Email rosana@nutricao.ufrj.br

the food intake of children under 3 years old in a metropolitan area of Brazil with a high percentage of families living in poverty.

Methods

Study population and procedures

The data included in the present analysis were obtained from a population-based cross-sectional study using a representative sample of households in the district of Campos Elíseos, in the city of Duque de Caxias, in Rio de Janeiro metropolitan area, Brazil. This place was in seventh position in 2003 among the most extreme areas of poverty in the State of Rio de Janeiro⁽¹⁷⁾. Details of the sample design have been previously described by Salles-Costa *et al.*⁽¹⁸⁾. The sample analysed comprised 1085 households, and 37% of these households (n 402) included children aged from 6 to 30 months. Three hundred and eighty-three households were able to provide complete information on the food intake of both children and their parents. Seventeen households were excluded because the adult interviewed was not the primary caregiver. One child and one adult from each household were randomly selected. Ultimately, 366 households with children and their parents (284 mothers and eighty-two fathers) were evaluated. Data collection was conducted from May to December 2005. The interview team comprised fifteen local residents who had graduated from high school and five nutritionists (whose role was to evaluate the information on children's food intake). All interviewers were previously trained in the dietary measurement and socio-economic and demographic questionnaire. The interviewers were trained by nutritionists with expertise in population surveys for the administration of the questionnaire (demographic information and food intake). All adult participants provided signed informed consent and the research was approved by the Ethics Committee of the State University of Rio de Janeiro.

Socio-economic and demographic characteristics

A questionnaire was administered to the adult responsible for the family (e.g. child's father or mother). The variables included in the present study were as follows: (i) monthly per capita family income (total family income divided by the number of residents who depend on this income); (ii) household food insecurity estimated using the Brazilian Food Insecurity Scale (Escala Brasileira de Segurança Alimentar, EBIA), which was adapted and validated for Brazil by Pérez-Escamilla *et al.*⁽¹⁹⁾; (iii) number of residents in the household; (iv) parental age in years; (v) parental education level; and (vi) child's age in months.

Food intake of children and parents

Children's food intake was estimated using the average of two 24 h recalls administered on non-consecutive days by

trained nutritionists. Portion sizes of the reported foods were estimated using household measures. Energy and food intakes were estimated using the software program NutWin (2005; Department of Information Science applied to Health, Federal University of São Paulo, Brazil), which is based on the US Department of Agriculture database. For items that were not included in NutWin software food composition table, the nutritional composition was obtained from the Brazilian food composition table⁽²⁰⁾ (5% of foods analysed). Standard recipes and serving sizes were used to estimate the nutritional composition of preparations that were not included in the software database.

Parents' food intake was estimated using a semi-quantitative FFQ validated for Rio de Janeiro's adult population⁽²¹⁾. The FFQ was administered by trained interviewers. The FFQ consisted of eighty-two items with pre-set usual portions and eight options for the frequency of consumption (ranging from 3 times/d to never/almost never). Optical reading of the FFQ was performed to produce data about the portion sizes and frequencies of intake for each food. The values were converted to g/d.

In order to analyse the consumption data, the foods reported in the 24 h recalls and the FFQ were converted into grams and then into eight food groups based on the recommendations of the Ministry of Health of Brazil⁽²²⁾ for children: (i) breads, cereals and tubers (rice, flour, bread, potatoes, yam, polenta, pasta, salt biscuits, sweet biscuits without filling, children's flour); (ii) beans; (iii) greens and vegetables; (iv) fruits; (v) milk and dairy products; (vi) meat (beef, chicken, fish and sausage) and eggs; (vii) sugars (chocolate powder, ice cream, candy, cookies, jam, jelly, chocolate bars, pies and puddings, sweets, processed juices and soft drinks); and (viii) oils and fats (oil, margarine, fried foods and pizza). Coffee consumption was evaluated separately due to the high consumption of this product reported in previous studies⁽⁷⁾. The food groups provided mutually exclusive estimates of intake.

Data analysis

For the statistical analysis, mean values and their standard errors or proportion distributions and their 95% confidence intervals were estimated for each category of variables in order to characterize the study population. The intra-class correlation coefficient between the first and second 24 h recalls for total energy and each of the food groups was calculated to estimate the reliability of dietary assessment.

A hierarchical model was used to evaluate how socio-demographic variables and parental food intake were associated with children's food intake. This analysis was achieved using linear regression with three explanatory levels: (i) parents' sociodemographic characteristics (age and educational level) and children's age; (ii) parents' food intake; and (iii) socio-economic variables (monthly per capita family income, household food insecurity and

number of residents in the household; Table 1). In the first stage, for each of the nine food groups, linear regression models were used to test each variable, employing a significance level of less than 20% ($P < 0.20$) to include the variable in the model. Multivariate analysis was then performed at each hierarchical level with the inclusion of all significant variables from the bivariate analysis. First, analysis included the sociodemographic characteristics of the parents (age and educational level) and the age of the children (first level). The variables of this level that retained significance ($P < 0.05$) were maintained in the model. Parents' food intakes were added to this model (second level) and a further analysis was processed; variables with statistical significance previously stipulated ($P < 0.05$) were kept in the model. Then, the same procedures described to include variables of levels 1 and 2 were used to add socio-economic variables (third level) and a new analysis was processed.

The variables that remained associated in each level constituted the final hierarchical model. All analyses were adjusted for the children's energy intake as a continuous variable in each hierarchical level.

The database was developed and recorded in duplicate by previously trained staff, the analytical procedures were performed using the statistical software package Stata 11.0.

Results

The characteristics of the study population are presented in Table 2. The mean monthly per capita family income was \$US 93.90, and most households reported some food insecurity (71.3%). The average composition of the households was five members. The parents were, on average, 34 years old, and most had less than 8 years of education.

Table 1 Theoretical model of hierarchical analysis and variables considered in the associations with food intake in children; Duque de Caxias, Rio de Janeiro, Brazil, 2005

Step	Study variables
Step 1: Sociodemographic characteristics	Parental education (years) Parental age (years) Age of children (months)
Step 2: Parental food intake	Food groups (g): (i) breads, cereals and tubers (rice, flour, bread, potatoes, yam, polenta, pasta, salt biscuits, sweet biscuits without filling, children's flour); (ii) beans; (iii) greens and vegetables; (iv) fruits; (v) milk and dairy products; (vi) meat (beef, chicken, fish and sausage) and eggs; (vii) sugars (chocolate powder, ice cream, candy, cookies, jam, jelly, chocolate bars, pies and puddings, sweets, processed juices and soft drinks); (viii) oils and fats (oil, margarine, fried foods and pizza); and (ix) coffee
Step 3: Socio-economic variables	Monthly per capita family income (\$US) Household food insecurity‡ Number of residents in household

†Adjusted for children's total energy intake.

‡Food insecurity of the household estimated with the Brazilian Food Insecurity Scale⁽¹⁹⁾.

Table 2 Sample household sociodemographic characteristics; Duque de Caxias, Rio de Janeiro, Brazil, 2005

	Mean or %	SE or 95% CI
Socio-economic variables (<i>n</i> 366)†		
Monthly per capita family income (\$US)‡	93.90	132.30
Number of residents in household	4.9	1.7
Proportion of food insecurity (%)§		
Food security	28.7	24.0, 33.3
Mild food insecurity	35.8	30.8, 40.7
Moderate food insecurity	20.5	16.3, 24.6
Severe food insecurity	15.0	11.3, 18.7
Parental variables (<i>n</i> 357)†		
Age (years)	33.8	11.1
Parental education (%)		
<8 years	58.1	52.9, 62.3
≥8 years	41.9	36.7, 47.0
Children's variables (<i>n</i> 366)†		
Age (months)	19.9	6.5
Total energy intake (kJ/d)	4259	1791
Total energy intake (kcal/d)	1018	428

†The values differ due to losses.

‡\$US 1.00 = 1.61 Brazilian reais (25 August 2011).

§Food insecurity of the household estimated with the Brazilian Food Insecurity Scale⁽¹⁹⁾.

The reliability of the two 24h recalls was measured by the intra-class correlation coefficient and indicated greater values for energy compared with the food groups, but all measures were statistically significant ($P < 0.05$; energy = 0.68, cereals = 0.15, beans = 0.21, vegetables = 0.40, fruits = 0.46, milk and dairy products = 0.82, meats and eggs = 0.34, sugars = 0.12, fats = 0.90 and coffee = 0.47).

The results of the associations between the children's intake and the study variables after the hierarchical-level adjustments are displayed in Table 3. The age of the parents was positively associated with vegetable intake among the children. A greater level of education was directly associated with the intake of fat and negatively associated with the intake of beans. An increase in age of the children was positively associated with the intakes of meats and eggs, sugars and coffee, and negatively associated with vegetable intake. Regarding the influence of parental food intake on children's food intake, the results revealed positive associations with the intakes of breads, cereals and tubers, beans and fruits, independently of the other variables. The increase in monthly per capita family income was positively associated with fruit intake and inversely associated with the intakes of breads, cereals and tubers, and coffee. Furthermore, the increase of food insecurity in the household was inversely associated with the intakes of vegetables, meats and eggs, and sugars. The number of residents in the household demonstrated a positive association with coffee intake.

Discussion

In the present study, both sociodemographic level and parental food intake were independently associated with the food intake of children under 3 years of age. Older parents included a greater number of healthy items in their children's diet, even in a population with a high prevalence of food insecurity. In contrast to other studies^(10,11,23), a positive association between parental educational level and healthy foods in the children's diet was not observed. The data indicated an increase in fat intake and a reduced intake of beans with increasing income. In the population with a high level of food insecurity, the educational level was a marker for the inclusion of processed foods in the diet, which tends to have greater fat concentrations. Budget surveys in Brazil have revealed a trend towards a reduction in unprocessed food items, such as beans, with increasing family income^(24,25).

The association observed between child age and food intake is likely due to the introduction of complementary foods and a greater participation of children in family meals. However, the results revealed some negative aspects of infant feeding. The first one refers to the lower consumption of vegetables with advancing age, a result corroborated by other studies⁽²⁶⁻²⁸⁾, which may be related

Table 3 Final hierarchical regression model of the determinants of children's food intake for each food group; Duque de Caxias, Rio de Janeiro, Brazil, 2005

	Food group (g/d)																			
	Breads, cereals and tubers		Beans		Vegetables		Fruits		Milk and dairy products		Meats and eggs		Sugars		Fats		Coffee			
	β	95% CI	β	95% CI	β	95% CI	β	95% CI	β	95% CI	β	95% CI	β	95% CI	β	95% CI	β	95% CI		
Step 1: Sociodemographic characteristics																				
Parental age (years)	-	-	-	-	0.73*	0.11, 1.34	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Parental education (years)	-	-	-13.98*	-27.94, -0.03	-	-	-2.12**	-3.20, -1.05	-	-	-	-	-	-	3.52*	0.04, 7.01	-	-	-	-
Age of children (months)	-	-	0.11*	-0.003, 0.22	-	-	-	-	0.10*	0.03, 0.16	-	-	-	-	-	-	-	-	1.77**	0.71, 2.84
Step 2: Parental food intake	0.06*	0.003, 0.12	-	-	-	-	-	-	-	-	-	-	5.08*	1.85, 8.30	-	-	-	-	-	-
Step 3: Socio-economic factors	-0.09*	-0.18, -0.001	-	-	-	-	-	-	0.23*	0.09, 0.38	-	-	-	-	-	-	-	-	-	-
Monthly per capita family income (\$US)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-0.05*	-0.10, -0.001
Food insecurity (ordinal variable)†	-	-	-	-	-1.76*	-3.37, -0.14	-	-	-	-	-3.11*	-5.08, -1.13	-5.82*	-10.60, -1.03	-	-	-	-	-	-
Number of residents in household	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	7.87**	3.97, 11.78
R ² (explained variance)	0.34		0.04		0.07		0.16				0.28		0.22		0.06				0.09	

* $P < 0.05$, ** $P < 0.001$.
 †Adjusted for children's total energy intake.
 ‡Food insecurity of the household estimated with the Brazilian Food Insecurity Scale⁽¹⁹⁾.

to the replacement of the soup offered when the child is younger by the food consumed by adults. Diets with low amounts of vegetables are likely deficient in essential nutrients such as vitamins and minerals, which may increase the risk of disease⁽²⁹⁾. Another issue is the increase in consumption of sugars found in studies conducted in Brazil^(30,31) and elsewhere^(27,32), and the influence of the advertising market, globalization and the fast pace of life in large cities, which may be some of the primary factors responsible for such increase⁽³³⁾.

The positive association between parents' dietary intake and greater consumption of fruits by the children is corroborated by Hart *et al.*⁽¹⁶⁾ and Cooke *et al.*⁽⁹⁾. In the present study, the dietary intake of parents was also associated with children's consumption of breads, cereals and tubers and beans, likely because these are traditional foods for this population⁽²⁴⁾. Moreover, the combination of rice and beans is more common among low-income families⁽²⁴⁾, where the financial cost and energy density are important factors in food selection.

In the present study, all variables related to socio-economic context were included in the model due to the great sample homogeneity in relation to family income, which makes the use of other indices important. As observed in the analysis, the employed variables present different associations in relation to children's food intake, which reflects different meanings in the families' socio-economic level.

The high prevalence of household food insecurity may have influenced the low intakes of vegetables, fruits and meats, as observed in other studies^(7,26,34). In fact, among the socio-economic factors, income is a delimiter in food choices towards the availability of resources that allow access to food. According to Panigassi *et al.*⁽⁵⁾, in situations of food insecurity, the individual responsible for the household food demand optimizes his/her financial resources by buying basic and inexpensive food, which may be a possible explanation for the results. Considering this, the high cost of meats, fruits and vegetables for families with food insecurity, as well as the population studied in the city of Rio de Janeiro, may lead consumers to purchase foods that can meet the family's essential satiety and energy needs, such as rice, beans, potatoes, pasta and flour, as described by Lignani *et al.*⁽³⁵⁾.

Family composition was evaluated because crowding in the home is another variable linked to food insecurity⁽³⁶⁾. Considering the difficulty of access to food, an assumption is that in a smaller family, individuals are able to obtain a more varied diet; however, our results demonstrated an association only for a greater consumption of coffee as the number of household members increased.

Limitations of the present study should be noted, including the cross-sectional design, which precludes the observation of a cause-and-effect relationship. Another limitation is the use of different means to assess parents' and children's food intake, although assessment of the

food groups included in the FFQ for the parents minimized this limitation.

Another limitation in the study was the use of the 24 h recall to evaluate children's food intake. The estimated consumption of energy and nutrients in infancy is particularly challenging due to the greater number of errors in measuring the diet, which compromises the accuracy of the methods for assessing food intake in this age group⁽³⁷⁾. However, the 24 h recall is the main method used to evaluate children's food intake⁽³⁸⁾. According to Salles-Costa *et al.*⁽³⁹⁾, the use of two 24 h recalls among children is appropriate for evaluating the intakes of energy, carbohydrates, protein, lipids and other micronutrients (Ca, Fe, vitamin C). In Salles-Costa *et al.*'s study, which used the same sample as the present study, the authors observed that the CV ratios (within- to between-person variation) for most nutrients were <1 among the younger (6 to 17 months) and older (18 to 30 months) children, considering the average of two days of 24 h recalls. Considering this result, the average of two 24 h recalls may be a good measure for stable long-term diet.

Conclusions

The present results indicate that for the promotion of healthy eating habits in childhood, programmes should consider all aspects related to children's food intake. The improvement of socio-economic indicators is essential for ensuring greater access to food, but the importance of the food choices of parents and their influence on infant feeding cannot be forgotten. Therefore, strategies should be adopted for the nutritional guidance offered to parents, especially during the introduction of complementary foods in children's diet.

Acknowledgements

Sources of funding: This study was supported by the National Research Council (CNPq; Grant CT-Agronegócio MCT/CNPq/MDS-2003) and the Brazilian National Cancer Institute – Ministry of Health. *Conflicts of interest:* No conflicts of interest are declared. *Authors' contributions:* G.d.S.B. participated in data collection, the manuscript concept, statistical analysis, and writing and revising the manuscript. R.S. was responsible for support, conception and coordination on the study design, and writing and revising the manuscript. R.S.-C. participated in the concept and design of the study, the coordination and supervision of the data collection, and writing and revising the manuscript.

References

1. US Department of Agriculture & US Department of Health and Human Services (2010) *Dietary Guidelines for Americans*. Washington, DC: US Government Printing Office; available at <http://www.dietaryguidelines.gov>

2. American Dietetic Association (2008) Position of the American Dietetic Association: nutrition guidance for healthy children ages 2 to 11 years. *J Am Diet Assoc* **108**, 1038–1047.
3. Osório MM, Ribeiro MA, Costa EC *et al.* (2009) Disponibilidade familiar de alimentos na Zona da Mata e Semi-Árido do Nordeste do Brasil. *Rev Nutr* **22**, 319–329.
4. Patrick H & Nicklas TA (2005) A review of family and social determinants of children's eating patterns and diet quality. *J Am Coll Nutr* **24**, 83–92.
5. Panigassi G, Segall-Corrêa AM, Marin-León L *et al.* (2008) Insegurança alimentar intrafamiliar e perfil de consumo de alimentos. *Rev Nutr* **21**, Suppl., S135–S144.
6. Rosas LG, Harley K, Fernald LCH *et al.* (2009) Dietary associations of household food insecurity among children of Mexican descent: results of a binational study. *J Am Diet Assoc* **109**, 2001–2009.
7. Antunes MML, Sichieri R & Salles-Costa R (2010) Consumo alimentar de crianças menores de três anos residentes em área de alta prevalência de insegurança alimentar domiciliar. *Cad Saude Publica* **26**, 1642–1650.
8. Santos JV, Gigante DP & Domingues MR (2010) Prevalência de insegurança alimentar em Pelotas, Rio Grande do Sul, Brasil, e estado nutricional de indivíduos que vivem nessa condição. *Cad Saude Publica* **26**, 41–49.
9. Cooke LJ, Wardle J, Gibson EL *et al.* (2004) Demographic, familial and trait predictors of fruit and vegetable consumption by pre-school children. *Public Health Nutr* **7**, 295–302.
10. Hendricks K, Briefel R, Novak T *et al.* (2006) Maternal and child characteristics associated with infant and toddler feeding practices. *J Am Diet Assoc* **106**, 1 Suppl. 1, S135–S148.
11. Sausenthaler S, Kompauer I, Mielck A *et al.* (2007) Impact of parental education and income inequality on children's food intake. *Public Health Nutr* **10**, 24–33.
12. Vereecken CA, Keukekier E & Maes L (2004) Influence of mother's education level on food parenting practices and food habits of young children. *Appetite* **43**, 93–103.
13. Robinson S, Marriott L, Poole J *et al.* (2007) Dietary patterns in infancy: the importance of maternal and family influences on feeding practice. *Br J Nutr* **98**, 1029–1037.
14. Fisk CM, Crozier SR, Inskip HM *et al.* (2011) Influences on the quality of young children's diets: the importance of maternal food choices. *Br J Nutr* **105**, 287–296.
15. Pappas MA, Hurlley KM, Quigg AM *et al.* (2009) Low-income, African American adolescent mothers and their toddlers exhibit similar dietary variety patterns. *J Nutr Educ Behav* **41**, 87–94.
16. Hart CN, Raynor HA, Jelalian E *et al.* (2010) The association of maternal food intake and infants' and toddlers' food intake. *Child Care Health Dev* **36**, 396–403.
17. Instituto Brasileiro de Geografia e Estatística (2000) *Censo Demográfico*. Rio de Janeiro: IBGE.
18. Salles-Costa R, Pereira RA, Vasconcellos MTL *et al.* (2008) Associação entre fatores socioeconômicos e insegurança alimentar: estudo de base populacional na Região Metropolitana do Rio de Janeiro, Brasil. *Rev Nutr* **21**, 99–109.
19. Pérez-Escamilla R, Segall-Corrêa AM, Kurdian Maranhã L *et al.* (2004) An adapted version of the US Department of Agriculture Food Insecurity module is a valid tool for assessing household food insecurity in Campinas, Brazil. *J Nutr* **134**, 1923–1928.
20. Núcleo de Estudos e Pesquisas em Alimentação (2004) *Tabela de Composição de Alimentos – TACO*. Campinas: Flamboyant.
21. Sichieri R & Everhart J (1998) Validity of a Brazilian food frequency questionnaire against dietary recalls and estimated energy intake. *Nutr Res* **18**, 1649–1659.
22. Ministério da Saúde (2010) *Dez passos para uma alimentação saudável: guia alimentar para crianças menores de dois anos*. Brasília: Ministério da Saúde.
23. Moreira P, Santos S, Padrão P *et al.* (2010) Food patterns according to sociodemographics, physical activity, sleeping and obesity in Portuguese children. *Int J Environ Res Public Health* **7**, 1121–1138.
24. Instituto Brasileiro de Geografia e Estatística (2004) *Pesquisa de Orçamentos Familiares – POF 2002/2003: Análise da disponibilidade domiciliar de alimentos e do estado nutricional no Brasil*. Rio de Janeiro: IBGE.
25. Instituto Brasileiro de Geografia e Estatística (2010) *Pesquisa de Orçamentos Familiares – POF 2008/2009: Avaliação nutricional da disponibilidade domiciliar de alimentos no Brasil*. Rio de Janeiro: IBGE.
26. Lorson BA, Melgar-Quinonez H & Taylor C (2009) Correlates of fruit and vegetable intakes in US children. *J Am Diet Assoc* **109**, 474–478.
27. Fox MK, Pac S, Devaney B *et al.* (2004) Feeding infants and toddlers study: what foods are infants and toddlers eating? *J Am Diet Assoc* **104**, 1 Suppl. 1, S22–S30.
28. Siega-Riz AM, Deming DM, Reidy KC *et al.* (2010) Food consumption patterns of infants and toddlers: where are we now? *J Am Diet Assoc* **110**, 12 Suppl., S38–S51.
29. Perry CL, Bishop DB, Taylor GL *et al.* (2004) A randomized school trial of environmental strategies to encourage fruit and vegetable consumption among children. *Health Educ Behav* **31**, 65–76.
30. Corrêa EM, Corso ACT, Moreira EAM *et al.* (2009) Alimentação complementar e características maternas de crianças menores de dois anos de idade em Florianópolis (SC). *Rev Paul Pediatr* **27**, 258–264.
31. Caetano MC, Ortiz TTO, Silva SGL *et al.* (2010) Alimentação complementar: práticas inadequadas em lactentes. *J Pediatr* **86**, 196–201.
32. Webb KL, Lahti-Koski M, Rutishauser I *et al.* (2006) Consumption of 'extra' foods (energy-dense, nutrient-poor) among children aged 16–24 months from western Sydney, Australia. *Public Health Nutr* **9**, 1035–1044.
33. Toloni MHA, Longo-Silva G, Goulart RMM *et al.* (2011) Introdução de alimentos industrializados e de alimentos de uso tradicional na dieta de crianças de creches públicas no Município de São Paulo. *Rev Nutr* **24**, 61–70.
34. Dave JM, Evans AE, Saunders RP *et al.* (2009) Associations among food insecurity, acculturation, demographic factors, and fruit and vegetable intake at home in Hispanic children. *J Am Diet Assoc* **109**, 697–701.
35. Lignani JB, Sichieri R, Burlandy L *et al.* (2011) Changes in food consumption among the Programa Bolsa Família participant families in Brazil. *Public Health Nutr* **14**, 785–792.
36. Instituto Brasileiro de Geografia e Estatística (2010) *Pesquisa Nacional por Amostra de Domicílios – Segurança Alimentar 2004/2009*. Rio de Janeiro: IBGE.
37. Livingstone MBE, Robson PJ & Wallace JM (2004) Issues in dietary intake assessment of children and adolescents. *Br J Nutr* **92**, 213–222.
38. Guinn CH, Baxter SD, Hardin JW *et al.* (2008) Intrusions in children's dietary recalls: the roles of BMI, sex, race, interview protocol, and social desirability. *Obesity (Silver Spring)* **16**, 2169–2174.
39. Salles-Costa R, Barroso GS, Mello MA *et al.* (2010) Sources of variation in energy and nutrient intakes among children from six to thirty months old in a population-based study. *Cad Saude Publica* **26**, 1175–1186.