

## Main Article

Prof E Okur takes responsibility for the integrity of the content of the paper

**Cite this article:** Sivrice ME, Yasan H, Okur N, Okur E, Buyukcelik B, Kumbul YC. Prevalence of chronic maxillary atelectasis: a radiological study. *J Laryngol Otol* 2022;**136**:1240–1244. <https://doi.org/10.1017/S0022215122000056>

Accepted: 6 January 2022  
First published online: 10 January 2022


### Key words:

Chronic Maxillary Atelectasis; Enophthalmos; Hypoglobus; Maxillary Sinus; Silent Sinus Syndrome

### Author for correspondence:

Prof Erdoğan Okur, Department of Oto-Rhino-Laryngology and Head and Neck Surgery, Faculty of Medicine, Süleyman Demirel University, Isparta, Turkey  
E-mail: [erdoganokur@gmail.com](mailto:erdoganokur@gmail.com)  
Fax: +90 2462 112 830

# Prevalence of chronic maxillary atelectasis: a radiological study

M E Sivrice<sup>1</sup> , H Yasan<sup>1</sup>, N Okur<sup>2</sup>, E Okur<sup>1</sup>, B Buyukcelik<sup>1</sup> and Y C Kumbul<sup>1</sup>

Departments of <sup>1</sup>Oto-Rhino-Laryngology and Head and Neck Surgery, and <sup>2</sup>Radiology, Faculty of Medicine, Süleyman Demirel University, Isparta, Turkey

## Abstract

**Objectives.** Chronic maxillary atelectasis is an infrequent entity and data on its prevalence are lacking. This study investigated the prevalence of chronic maxillary atelectasis and aimed to determine the bilaterality of this entity.

**Methods.** The data for 5835 patients who underwent paranasal sinus computed tomography from 2016 to 2020 were retrospectively analysed.

**Results.** Fifty-four patients were diagnosed with chronic maxillary atelectasis; its prevalence was 0.92 per cent. The mean age of these 54 patients was  $42.98 \pm 18.89$  years (range, 18–85 years); 17 of the patients were female and 37 were male. Chronic maxillary atelectasis was unilateral in 42 patients and bilateral in 12 patients (22.2 per cent). Eight patients were found to have enophthalmos with apparent facial asymmetry.

**Conclusion.** The prevalence of bilateral chronic maxillary atelectasis may be higher than previously reported and bilaterality may increase as the number of diagnosed cases increases. A unified classification is also proposed, which describes the silent sinus syndrome as chronic maxillary atelectasis IIIS.

## Introduction

Structurally, the paranasal sinuses can be normal, small (hypoplastic) or large (megasinus or pneumocele), but their size may decrease or increase over time. An example of this increase is pneumosinus dilatans, and an example of its decrease is chronic maxillary atelectasis.<sup>1</sup> Chronic maxillary atelectasis is an unusual and underdiagnosed entity of permanent and increasing reduction in maxillary sinus dimensions that results in an antral wall collapse. Chronic maxillary atelectasis was first reported in 1964 by Montgomery, but the term ‘chronic maxillary atelectasis’ was first described in 1997 by Kass *et al.*<sup>2,3</sup> Kass *et al.* also classified chronic maxillary atelectasis into three progressive grades in accordance with the level of sinus wall collapse.<sup>4</sup>

Previously, other terms have been used for this entity in the literature, such as imploding antrum, silent sinus syndrome and acquired maxillary sinus hypoplasia. In 1994, Soparkar *et al.* used the term ‘silent sinus syndrome’ to describe their cases with spontaneous enophthalmos and unilateral collapse of the maxillary sinus in the absence of sino-nasal symptoms.<sup>5</sup> Although silent sinus syndrome and chronic maxillary atelectasis are still mentioned separately in studies, de Dorlodot *et al.* showed that the only distinction between grade III chronic maxillary atelectasis and silent sinus syndrome is the presence or absence of sinusitis symptoms. Diagnostic criteria for chronic maxillary atelectasis and silent sinus syndrome are described in Table 1.<sup>1</sup> In addition, maxillary sinus hypoplasia is congenital and is included in the differential diagnosis of chronic maxillary atelectasis, so it should be distinguished from maxillary atelectasis.<sup>6</sup>

Chronic maxillary atelectasis has generally been described as a rare and unilateral entity; bilateral chronic maxillary atelectasis cases, as reported in the literature, are extremely uncommon.<sup>4</sup> We believe that there are a lack of data regarding chronic maxillary atelectasis prevalence in the literature. This radiological study investigated the prevalence of chronic maxillary atelectasis encountered in our tertiary hospital. In addition, we aimed to determine whether the bilaterality of this entity is as rare as previously mentioned.

## Materials and methods

This research was conducted in our university hospital between January 2016 and January 2020. The ethics committee of the university approved the research protocol (date: 26 April 2021; approval number: 187). This research was conducted in accordance with the international ethical standards of the Helsinki Declaration.

## Patient selection

Our study included 5835 patients who underwent paranasal sinus computed tomography (CT) for the examination of rhinological or orbital complaints.

**Table 1.** Staging of chronic maxillary atelectasis<sup>1</sup>

Clinical or radiological feature	CMA grade I	CMA grade II	CMA grade III	Silent sinus syndrome CMA IIIS*
Enophthalmos	Absent	Absent	Present	Present
Maxillary sinus volume reduction	Radiological membranous deformity	Radiological bony deformity	Radiological volume reduction with clinical repercussions	Radiological volume reduction with clinical repercussions
Sinusitis or facial pain symptoms	Absent or present	Absent or present	Absent or present	Absent

\*Our unified classification. CMA = chronic maxillary atelectasis

Paranasal CT examinations were performed on a 128-slice CT scanner (Somatom Definition; Siemens Healthcare, Forchheim, Germany), with approximately 120–180 images per CT study. All paranasal sinus scans covered from the roof of the frontal sinuses to the maxillary alveolus just below the hard palate, on the axial plane. The imaging parameters were: tube voltage = 80 kVp, tube current = 120 mA, slice thickness and reconstruction interval = 1 mm, pitch factor = 1, matrix size = 512 × 512, field of view = 14 cm × 17 cm, window width = 2000 HU, and window level = 400 HU. Reconstructed images of coronal and sagittal planes were generated using the separate workstation system.

One radiologist and one ENT specialist evaluated all of the paranasal sinus CT scans. Cases that met the criteria for radiological chronic maxillary atelectasis, such as inward bowing of at least one of the sinus walls and an opacification of the sinus antrum, were included in the study group. Cases with previous facial or orbital trauma (including sinus surgery), congenital facial deformity (such as maxillary hypoplasia) and other causes of enophthalmos were excluded. Data were collected from patients' charts.

### Statistical analyses


Data were transferred to SPSS version 23 software (IBM, Chicago, Illinois, USA) and statistically analysed. Before proceeding to the statistical analysis, controls were made to ensure the absence of data entry errors and confirm that the parameters were within the expected ranges. Regarding descriptive statistics, mean and standard deviation values were calculated for continuous variables, and prevalence (*n*) and percentage values were determined for categorical variables. Relationships between categorical variables were examined by chi-square test analysis. A *p*-value of less than 0.05 was accepted as significant in all analyses.

## Results


### Demographic features and clinical characteristics

The CT scans of 5835 patients were evaluated; 54 of them were diagnosed with chronic maxillary atelectasis. The mean age of these 54 patients was 42.98 ± 18.89 years (range, 18–85 years); 17 patients were female and 37 patients were male. Chronic maxillary atelectasis was unilateral in 42 patients (28 right, 14 left) and bilateral in 12 patients. Eight patients were found to have enophthalmos with apparent facial asymmetry. Two patients had diplopia.

General symptoms of sinusitis were present in 21 patients and absent in 33 patients. Eleven patients had grade I (Figure 1), 35 patients had grade II (Figure 2) and 8 patients had grade III (Figure 3) chronic maxillary atelectasis. Seven



**Fig. 1.** Coronal computed tomography scan of the paranasal sinus demonstrating lateralisation of the right uncinate processes (grade I chronic maxillary atelectasis).




**Fig. 2.** Coronal computed tomography scan of the paranasal sinus demonstrating inward bowing of the sinus walls and opacification of the sinus (left) (grade II chronic maxillary atelectasis).

patients met the silent sinus syndrome criteria; one patient with chronic maxillary atelectasis grade III was symptomatic and therefore not considered to have silent sinus syndrome. Table 2 reports the clinical features.

### Radiological features

Maxillary sinus opacification was complete in 28 patients and partial in 26 patients. Downward displacement of the orbital floor was observed in 15 patients. Other signs of sinus diseases were observed in 19 patients. Table 3 reports the radiological features.



**Fig. 3.** Coronal computed tomography scan of the paranasal sinus demonstrating downward displacement of the orbital floor and right maxillary sinus opacity (right) (grade III chronic maxillary atelectasis).

### Radiographic comparison of unilateral and bilateral cases

Fourteen ipsilateral nasal septal deviations were observed in unilateral cases and two in bilateral cases. There was no statistically significant difference between unilateral and bilateral cases for ipsilateral nasal septal deviation ( $p = 0.322$ ).

Eleven ipsilateral middle turbinate lateralizations were observed in unilateral cases and one in bilateral cases. There was no statistically significant difference between unilateral and bilateral cases for ipsilateral middle turbinate lateralization ( $p = 0.261$ ).

In addition, there was no statistically significant difference between unilateral and bilateral cases in terms of middle turbinate variations (concha bullosa, paradoxical middle turbinate) ( $p = 0.706$ ). Table 4 reports the radiographic comparison of unilateral and bilateral cases.

### Discussion

Chronic maxillary atelectasis is known as a rare and underdiagnosed disease. The detection of chronic maxillary atelectasis in 54 of 5835 CT scans performed in our tertiary care hospital supports the fact that this disease is very rare. Chronic maxillary atelectasis has often been described as a unilateral condition.<sup>7</sup> Although the majority of chronic maxillary atelectasis cases are unilateral, there are also bilateral chronic maxillary atelectasis cases reported in the literature. Kaas *et al.* reported 2 bilateral chronic maxillary atelectasis cases in their series of 22 patients.<sup>3</sup> In 2019, Ho *et al.* published a case series on bilateral chronic maxillary atelectasis with three patients, and conducted a literature review of bilateral chronic maxillary atelectasis cases in their research.<sup>4</sup> Our study comprised 12 bilateral cases; this corresponds to 22.2 per cent of all our cases. We think that the number of bilateral cases may be higher than previously expected, and the rate of bilaterality may increase as the number of diagnosed cases increases.

As previously shown in the literature, chronic maxillary atelectasis and silent sinus syndrome are two terms for the same clinical entity; hence, we think that these two terms should be combined to avoid diagnostic confusion.<sup>1,8</sup> Therefore, silent sinus syndrome is not discussed separately from chronic maxillary atelectasis in our article. We propose

**Table 2.** Clinical features

Parameter	Cases (n (%))
Side affected	
– Right	28 (51.9)
– Left	14 (25.9)
Altered facial appearance?	
– Bilateral	12 (22.2)
– No	46 (85.2)
– Yes	8 (14.8)
Enophthalmos?	
– No	46 (85.2)
– Yes	8 (14.8)
Diplopia?	
– No	52 (96.3)
– Yes	2 (3.7)
General sinusitis symptoms?	
– No	33 (61.1)
– Yes	21 (38.9)
Rhinorrhoea?	
– No	37 (68.5)
– Yes	17 (31.5)
Nasal congestion?	
– No	40 (74.1)
– Yes	14 (25.9)
Post-nasal drip?	
– No	36 (66.7)
– Yes	18 (33.3)
Facial pain or pressure?	
– No	41 (75.9)
– Yes	13 (24.1)
History of sinus disease?	
– No	44 (81.5)
– Yes	10 (18.5)
CMA	
– Grade 1	11 (20.4)
– Grade 2	35 (64.8)
– Grade 3	8 (14.8)
Silent sinus syndrome?	
– No	47 (87.0)
– Yes	7 (13.0)

CMA = chronic maxillary atelectasis

a unified classification describing silent sinus syndrome as chronic maxillary atelectasis IIIS.

There are two main theories for the development of chronic maxillary atelectasis: obstruction of outflow and mechanical theory.<sup>9</sup> Chronic hypoventilation of the maxillary sinus can have mechanical causes such as anatomical variation (septal deviation, concha bullosa of the middle turbinate, paradoxical middle turbinate), or inflammatory causes such as chronic rhinosinusitis.<sup>10</sup> It has been postulated that a deficiency in the posterior attachment of the uncinat process to the inferior

**Table 3.** Radiological features

Parameter	Cases (n (%))
Maxillary opacification	
– Incomplete	26 (48.1)
– Complete	28 (51.9)
Maxillary sinus contraction?	
– No	13 (24.1)
– Yes	41 (75.9)
Downward displacement of orbital floor?	
– No	39 (72.2)
– Yes	15 (27.8)
Other sinus diseases?	
– No	35 (64.8)
– Yes	19 (35.2)
Bone density loss in maxillary wall roof?	
– No	37 (68.5)
– Yes	17 (31.5)
Bone density loss in anterior wall?	
– No	46 (85.2)
– Yes	8 (14.8)
Bone density loss in medial wall?	
– No	33 (61.1)
– Yes	21 (38.9)
Bone density loss in posterolateral wall?	
– No	39 (72.2)
– Yes	15 (27.8)

turbinate, and a lateralised or hypermobile medial infundibular wall, could cause chronic hypoventilation of the sinus cavity via a valve effect.<sup>11,12</sup> Nevertheless, these anatomical variations explain only a few cases of chronic maxillary atelectasis. In our study, ipsilateral nasal septal deviations were

**Table 4.** Radiographic comparison of unilateral and bilateral cases

Parameter	Unilateral cases (n (%))	Bilateral cases (n (%))	P-value
Septal deviation?			
– No	24 (57.1)	10 (83.3)	0.322
– Ipsilateral	14 (33.3)	2 (16.7)	
– Contralateral	4 (9.5)	0 (0.0)	
Ipsilateral middle turbinate lateralisation?			
– No	31 (73.8)	11 (91.7)	0.261
– Yes	11 (26.2)	1 (8.3)	
Middle turbinate variation?			
– No	25 (59.5)	6 (50.0)	0.706
– Concha bullosa	15 (35.7)	6 (50.0)	
– Paradoxical middle turbinate	2 (4.8)	0 (0.0)	

observed in 33.3 per cent of the unilateral cases and 16.7 per cent of bilateral cases. In addition, ipsilateral middle turbinate lateralisation and middle turbinate variations were not observed in the majority of the patients. Other signs of sinus diseases were also absent in most cases. Further research on the aetiology of this entity is needed.

Regardless of its aetiology, blockage to mucus drainage causes chronic hypoventilation of the cavity. This results from the resorption of the mucosal gas and the development of negative pressure within the sinus cavity. Over time, these conditions lead to demineralisation and remodelling of the bone, with subsequent thinning and inward bowing of the maxillary walls.<sup>13–15</sup> This negative pressure theory has been confirmed in animal studies.<sup>16</sup> In addition, Kass *et al.* measured negative pressures during the sinus surgery of chronic maxillary atelectasis patients.<sup>15</sup>

The number of patients with chronic sinusitis and ostial obstruction is higher than the number of patients with chronic maxillary atelectasis. Therefore, it is clear that further explanations are needed. The mechanical theory hypothesises that masticatory muscle contraction generates negative pressure in the infratemporal and pterygopalatine fossa, and the transmitted negative pressure can cause antral wall collapse.<sup>17</sup>

Currently, the accepted method for managing chronic maxillary atelectasis is uncinectomy and middle meatal antrostomy with an endoscopic approach. The primary argument for the treatment is the need for and optimal timing of orbital floor reconstruction.<sup>18</sup> Some researchers prefer a one-stage surgery with endoscopic maxillary antrostomy and orbital floor reconstruction.<sup>11,19</sup> However, most surgeons agree that the one-stage approach with orbital floor reconstruction is not needed in the majority of cases.<sup>20,21</sup> Most patients achieve spontaneous recovery of the enophthalmos and hypoglobus only by re-ventilation of the maxillary sinus.<sup>22</sup> In order to avoid unnecessary surgical interventions, surgeons frequently recommend a six-month delay between the two steps. We accept and use this two-step surgical approach in our clinic.

The main weaknesses of the present study include its retrospective design and the lack of endoscopic records for the patients. Another limitation is that CT was performed on patients who presented with sinus complaints; therefore, the prevalence might only be applied to that group, rather than a healthy population. To the best of our knowledge, this is the first radiological study of the prevalence of chronic maxillary atelectasis. In addition, our study reports on the largest numbers of unilateral and bilateral chronic maxillary atelectasis cases so far.

- Chronic maxillary atelectasis is an infrequent entity and data on its prevalence are lacking
- This study analysed 5835 patients with paranasal sinus computed tomography data to determine its prevalence
- The prevalence of chronic maxillary atelectasis in our study was 0.92 per cent; 22.2 per cent of cases were bilateral
- The prevalence of bilateral chronic maxillary atelectasis may be higher than previously reported, and bilaterality may increase with the number of diagnoses
- Chronic maxillary atelectasis and silent sinus syndrome describe the same clinical entity; these two terms should be combined to avoid diagnostic confusion
- A unified classification system is proposed, describing silent sinus syndrome as chronic maxillary atelectasis IIIS

To date, chronic maxillary atelectasis has been rarely reported in the literature and data on its prevalence are lacking. The prevalence of bilateral chronic maxillary atelectasis may be higher than previously reported, and the rate of

bilaterality may increase as the number of diagnosed cases increases. Chronic maxillary atelectasis and silent sinus syndrome are two terms for the same clinical entity, and we therefore think that these two terms can be combined to avoid diagnostic confusion. We propose a unified classification describing silent sinus syndrome as chronic maxillary atelectasis IIIS. Further studies are needed to evaluate the aetiology, pathogenesis and prevalence of this entity.

**Competing interests.** None declared

## References

- de Dorlodot C, Collet S, Rombaux P, Horoi M, Hassid S, Eloy P. Chronic maxillary atelectasis and silent sinus syndrome: two faces of the same clinical entity. *Eur Arch Otorhinolaryngol* 2017;**274**:3367–73
- Montgomery W. Mucocele of the maxillary sinus causing enophthalmos. *Eye Ear Nose Throat Mon* 1964;**43**:41–4
- Kass ES, Salman S, Weber AL, Rubin PAD, Montgomery WW. Chronic maxillary atelectasis. *Ann Otol Rhinol Laryngol* 1997;**106**:109–16
- Ho JPK, Wong E, Gunaratne DA, Singh N. Chronic maxillary atelectasis (including silent sinus syndrome) can present bilaterally. *J Laryngol Otol* 2019;**133**:251–5
- Soparkar CNS, Patrinely JR, Cuaycong MJ, Dailey RA, Kersten RC, Rubin PA *et al.* The silent sinus syndrome: a cause of spontaneous enophthalmos. *Ophthalmology* 1994;**101**:772–8
- Gunaratne DA, Hasan Z, Floros P, Singh N. Bilateral stage I chronic maxillary atelectasis: a case report. *Int J Surg Case Rep* 2016;**26**:53–6
- Ho JPK, Wong E, Singh NP. Bilateral chronic maxillary atelectasis with a unilateral accessory ostium. *Eur J Rhinol Allergy* 2020;**3**:23–5
- Brandt MG, Wright ED. The silent sinus syndrome is a form of chronic maxillary atelectasis: a systematic review of all reported cases. *Am J Rhinol* 2008;**22**:68–73
- Cobb AR, Murthy R, Cousin GC, El-Rasheed A, Toma A, Uddin J *et al.* Silent sinus syndrome. *Br J Oral Maxillofac Surg* 2012;**50**:81–5
- Blackwell KE, Goldberg RA, Calcaterra TC. Atelectasis of the maxillary sinus with enophthalmos and midface depression. *Ann Otol Rhinol Laryngol* 1993;**102**:429–32
- Rose GE, Lund VJ. Clinical features and treatment of late enophthalmos after orbital decompression: a condition suggesting cause for idiopathic “imploding antrum” (silent sinus) syndrome. *Ophthalmology* 2003;**110**:819–26
- Hobbs CG, Saunders MW, Potts MJ. Spontaneous enophthalmos: silent sinus syndrome. *J Laryngol Otol* 2004;**118**:310–22
- Illner A, Davidson HC, Harnsberger HR, Hoffman J. The silent sinus syndrome: clinical and radiographic findings. *AJR Am J Roentgenol* 2002;**178**:503–6
- Numa WA, Desai U, Gold DR, Heher KL, Annino DJ. Silent sinus syndrome: a case presentation and comprehensive review of all 84 reported cases. *Ann Otol Rhinol Laryngol* 2005;**114**:688–94
- Kass ES, Salman S, Montgomery WW. Manometric study of complete ostial occlusion in chronic maxillary atelectasis. *Laryngoscope* 1996;**106**:1255–8
- Scharf KE, Lawson W, Shapiro JM, Gannon PJ. Pressure measurements in the normal and occluded rabbit maxillary sinus. *Laryngoscope* 1995;**105**:570–4
- Baujart B, Derbez R, Rossarie R, Hardy T, Wagner I, Krastinova D *et al.* Silent sinus syndrome: a mechanical theory. *Orbit* 2006;**25**:145–8
- Cardesin A, Escamilla Y, Romera M, Molina JA. Single surgical step for endoscopic surgery and orbital reconstruction of a silent sinus syndrome. *Acta Otorrinolaringol Esp* 2011;**64**:297–9
- Bossolesi P, Autelitano L, Brusati R, Castelnuovo P. The silent sinus syndrome: diagnosis and surgical treatment. *Rhinology* 2008;**46**:308–16
- Thomas RD, Graham SM, Carter KD, Nerad JA. Management of the orbital floor in silent sinus syndrome. *Am J Rhinol* 2003;**17**:97–100
- Facon F, Eloy P, Brasseur P, Collet S, Bertrand B. The silent sinus syndrome. *Eur Arch Otorhinolaryngol* 2006;**263**:567–71
- Ando A, Cruz AA. Management of enophthalmos and superior sulcus deformity induced by the silent sinus syndrome. *Aesthetic Plast Surg* 2005;**29**:74–7