

***Mycobacterium tuberculosis* exposure of livestock in a German dairy farm: implications for *intra vitam* diagnosis of bovine tuberculosis in an officially tuberculosis-free country**

T. EISENBERG^{1*}, A. NESSELER¹, C. SAUERWALD¹, U. KLING¹, K. RIBE¹,
U. KAIM¹, G. ALTHOFF¹, N. FIEGE¹, K. SCHLEZ¹, H.-P. HAMANN¹,
A. FAWZY^{1,2}, I. MOSER³, R. RIBE⁴, G. KRAFT⁴, M. ZSCHÖCK¹ AND
C. MENGE³

¹Hessian State Laboratory (LHL), Division for Veterinary Medicine, Gießen, Germany

²Department of Medicine and Infectious Diseases, Faculty of Veterinary Medicine, Cairo University, Egypt

³Institute of Molecular Pathogenesis, Friedrich-Loeffler-Institut (Federal Research Institute for Animal Health), Jena, Germany

⁴Veterinary Office, District Vogelsbergkreis, Lauterbach, Germany

Received 20 January 2015; Final revision 2 July 2015; Accepted 7 August 2015;
first published online 7 September 2015

SUMMARY

Germany has been an officially bovine tuberculosis (bTB)-free (OTF) country since 1996. Gradually rising numbers of bTB herd incidents due to *Mycobacterium bovis* and *M. caprae* in North-Western and Southern Germany during the last few years prompted the competent authorities to conduct a nationwide bTB survey in 2013/2014. This led to the detection of a dairy herd in which as many as 55 cattle reacted positively to consecutive *intra vitam* testing. Test-positive animals lacked visible lesions indicative of bTB at necropsy. Extensive mycobacterial culturing as well as molecular testing of samples from 11 tissues for members of the *M. tuberculosis* complex (MTC) yielded negative results throughout. However, caseous lymphadenitis of Ln. mandibularis accessorius was observed during meat inspection of a fattening pig from the same farm at regular slaughter at that time. Respective tissue samples tested MTC positive by polymerase chain reaction, and *M. tuberculosis* T1 family were identified by spoligotyping. Four human reactors within the farmer's family were also found to be immunoreactive. As exposure of livestock to *M. tuberculosis* is not generally considered, its impact may result in regulatory and practical difficulties when using protocols designed to detect classical bTB, particularly in OTF countries.

Key words: Bovine tuberculosis, legislation, monitoring, *Mycobacterium tuberculosis*, pig, skin test.

INTRODUCTION

Bovine tuberculosis (bTB) is one of the most relevant epizootics worldwide and leads to significant losses

in animal production. *Mycobacterium bovis* and *M. caprae* are the primary causative agents in countries of Central and Southern Europe. Both species are zoonotic and 99.9% genetically identical to *M. tuberculosis*. Together with other approved and recently described TB-causing mycobacterial species, i.e. *M. africanum*, *M. microti*, *M. pinnipedii*, *M. orygis*, *M. suricattae*, *M. mungi* and the 'dassie bacillus', they form the *M. tuberculosis* complex (MTC) [1–7]. Bovine *M. microti*

* Author for correspondence: Dr T. Eisenberg, Hessisches Landeslabor, Abteilung Veterinärmedizin, Schubertstr. 60/Haus 13, 35392 Gießen, Germany.
(Email: tobias.eisenberg@lhl.hessen.de)

infections are very rarely seen and appear to be associated with high rodent densities [8]. Occasionally, infections due to *M. tuberculosis* have been observed in cattle [9–12]. The prevalence of bTB caused by *M. tuberculosis* is low, mostly restricted to countries with a high burden of human TB, and normally does not exceed 1% of bTB herd incidents ('breakdowns'; summarized in [13]). The route of *M. tuberculosis* transmission to cattle may involve aerosol inhalation from infected humans, but oral transmission by ingestion of feed or water contaminated with infective human nasal secretions, faeces or urine is also possible [14].

In Germany, bTB prevalence as high as 59% at farm level was recorded for the first half of the last century [15]. National eradication programmes were established on a voluntary basis in 1952 in Western Germany and in 1955 in Eastern Germany and later became mandatory. Programmes were based on regular tuberculin skin testing of all cattle aged >6 weeks and removal of reactor animals. Germany finally met the requirements of Article 2(d) of Directive 64/432/EEC by having 99.9% of cattle herds certified bTB-free for at least 6 consecutive years and was subsequently declared officially bTB-free (OTF), effective from 1 July 1996 (Decision 97/76/EC). Consequently, Germany abandoned the elaborate and costly regular tuberculin testing and bTB surveillance was maintained at the level of official veterinary meat inspection instead.

From 1995 to 2012, the annual number of bTB incidents in cattle herds varied between two and 23. Numbers rose to 46 in 2013 (<http://tsis.fli.bund.de/Reports/Info.aspx>) following an intensified regional surveillance in two counties in Southern Germany bordering Austria. In this region, alpine farming practices were suspected of fostering transmission of *M. caprae* between free-ranging red deer and cattle. Indeed, an apparent local endemic driven by three different molecular types of *M. caprae* was discovered which accounted for 73.9% of German bTB cases in 2013 [16]. Other cases of bTB notified in 2013 affected farms in North-Western Germany with a wide spatial distribution and were exclusively attributed to *M. bovis* [17]. For a deeper insight into the true prevalence of bTB, a nationwide survey was prescribed due to a revision to the German Regulation on bTB issued on 12 July 2013. Between July 2013 and 30 April 2014, approximately 3350 cattle aged >2 years were selected from each federal state for single intradermal comparative cervical tuberculin (SICCT) testing. This nationwide survey revealed a suspicious case of bTB in a dairy farm, which was eventually attributed to

exposure to *M. tuberculosis*. This case, described here, led to significant difficulties in case declaration and implementation of appropriate legal measures and was accompanied by substantial financial losses for the affected farm.

METHODS

In September 2013 the study herd consisted of 234 cattle in total. Besides two fattening pigs for personal consumption, some cats were also kept on the farm and four persons regularly cared for the animals. Fifteen bovines aged >2 years were subjected to an *intra vitam* diagnosis of bTB. SICCT testing was performed with matched equipotent doses of bovine and avian purified protein derivative (PPD), respectively. Increase in skin thickness was measured 72 h after injection, and the reaction was interpreted as positive if clinical signs were observed or there was an increase in the thickness of the skin fold at the injection site together with a positive reaction to bovine PPD being >4 mm greater than the reaction to avian PPD. Inconclusive reactors (1–4 mm thickness increase) were re-tested by interferon-gamma release assay (IGRA; Bovigam[®], Prionics, Germany) within 3 weeks according to the manufacturer's instructions. If the IGRA result was positive, the animal was culled. Animals testing negative by IGRA were re-tested by SICCT and culled if reacting inconclusive. Pathological examinations were performed on the carcasses and organs with special regard to detection of lesions suspicious for bTB. For molecular and histological investigations, tissues of 11 prescribed organs and lymph nodes (lung, intestine, liver, spleen, kidney and respective tributary lymph nodes as well as retropharyngeal lymph nodes) were subjected to polymerase chain reaction (PCR) (duplex real time PCR for IS1081 and a hypothetical helicase (HELI) for the detection of MTC members; [18]) and fixed in neutral buffered formalin, paraffin embedded and routinely stained with haematoxylin and eosin, respectively. Sections showing inflammatory alterations, with granulomas in particular, were Ziehl–Neelsen (ZN)-stained for detection of acid-fast bacilli (AFB). Up to 12 tissue samples per animal were collected from cattle ($n = 55$), pig ($n = 1$) and cat ($n = 1$) between September 2013 and January 2014 and screened for pathological lesions, growth and/or genomic evidence for the presence of MTC members. Caseous lymphadenitis of one Ln. mandibularis accessorius was detected at regular slaughter in the pig. Mycobacterial culture was initiated in duplicate in two different

laboratories using standard procedures as described in the Official Collection of Diagnostic Methods released by the Friedrich-Loeffler-Institut/Federal Research Institute for Animal Health (FLI) based on DIN 222 standard [18] or with minor modifications. Briefly, tissue samples were homogenized and decontaminated and then inoculated onto three solid (slant) culture tubes [Loewenstein–Jensen medium with and without glycerol, Stonebrink medium (Merck, Germany)] and one liquid medium (Kirchner or Middlebrook broth; Merck) and incubated for up to 12 weeks at 37 °C. To inhibit growth of contaminant flora, all media were supplemented with PACT (polymyxin B, amphotericin, carbenicillin, and trimethoprim). Mycobacterial cultures were verified by duplex real-time PCR (*IS1081*, HELI) for MTC members (according to [18]). Molecular differentiation of members of MTC included spoligotyping [19] using a recently developed microarray [20]. Results were recorded automatically according to the nomenclature of international databases (SITVIT; MIRU-VNTRplus; mbovis.org). From cows with evidence of paratuberculosis, faecal samples were screened for the presence of *M. avium* subsp. *paratuberculosis* (MAP) by culture and also directly by real-time PCR (Life Technologies, Germany). Additionally, four milk filters from the milking machine were explicitly tested for MTC members by PCR and culture. Grown colonies of presumptive mycobacteria not belonging to MTC were analysed by 16S rRNA and *rpoB* gene sequence analysis as described previously [21, 22]. Additionally, presumptive mycobacterial isolates were submitted to spectroscopic analysis based on matrix-assisted laser desorption/ionization–time-of-flight mass spectrometry (MALDI–TOF MS; Biotyper, BrukerDaltonics, Germany) using a preparation method for mycobacteria according to the manufacturer’s instructions.

Data on meat inspection and possible condemnations were requested from the abattoir that had slaughtered cattle from the farm under investigation between September 2013 and March 2014.

RESULTS

Herd history, *intra vitam* testing in bovines

During the years 2010–2012, the owner of the herd had bought one bull originating from the same federal state. No other bovines were introduced into the herd. Of 15 cattle selected for testing in the mandated nationwide survey in 2013, two animals showed inconclusive

reactions upon initial SICCT testing. Both cows were re-tested with negative results by IGRA 3 weeks later. The remaining 174 bovines aged >6 weeks were tested by SICCT in mid-October. Of these, five had an inconclusive result and eight a positive result. Of the former, one reacted positively to a subsequent IGRA test and was culled, as were all of the latter. The next herd survey was conducted at end of November 2013, when 46 and 12 cows tested by SICCT were inconclusive and positive, respectively. All SICCT-positive bovines as well as 33 IGRA-positively re-tested cows were necropsied between December 2013 and January 2014. In January, one additional cow was culled after a positive SICCT test. A detailed scheme of SICCT testing reactivity and IGRA results is given in Figure 1. Altogether, 654 SICCT and 60 IGRA tests were performed on this herd producing 577 and 26 negative and 77 and 34 positive results, respectively.

Abnormal findings in bovines and their environment

Necropsies on 55 (23.5%) of 234 animals revealed no evidence of TB or bTB, neither histologically nor microscopically nor by culture or nucleic acid amplification techniques. ZN staining revealed no acid-fast rod-shaped bacteria in tissue sections from cattle. Mycobacterial growth was not observed in any of the tissue samples ($n = 532$) from 55 culled cattle up to 12 weeks of incubation. Altogether, analyses of 452 and 543 tissue samples and nine faecal samples were performed for detection of *IS1081* and HELI as well as for the detection of MAP, with negative results throughout. Meat inspection of 17 bovines sent for regular slaughter up to March 2014 failed to provide any findings indicative of bTB.

AFB were not detected in tissue samples ($n = 9$) from one euthanized cat; however, AFB consistent with fast-growing non-tuberculous mycobacteria could be propagated from milk filter samples ($n = 4$), in which few AFB were detected by microscopy. Partial 16S rRNA and *rpoB* gene sequencing identified all isolates as pure cultures of *M. setense* confirmed by mass spectrometry (data not shown).

Abnormal findings in a fattening pig raised on the farm

After regular slaughtering of one fattening pig, caseous lesions were noted in one Ln. mandibularis accessorius during meat inspection. A high-grade amount of AFB was detected in the affected lymph node tissue by direct ZN staining and histopathology. Mycobacterial culture




Fig. 1. Overview of test results during testing of a dairy herd (animals with negative test results throughout are not included in the Figure) for the presence of bovines having contact with members of the *M. tuberculosis* complex. SICCT, Single intradermal comparative cervical tuberculin test; IGRA, interferon-gamma release assay; PCR, double multiplex real-time PCR targeting *IS1081* and the gene for a hypothetical helicase each in combination with the gene for β -actin as amplification control with tissue obtained during necropsy.

for up to 20 weeks of incubation at 37 °C failed to grow AFB, whereas samples were positive by MTC PCR, yielding positive results with a cycle threshold (C_T) of 33–45. Molecular characterization directly carried out from extracted DNA revealed a *M. tuberculosis* member of the Euro-American SIT (spoligotype international type) T1 family (SITVIT database). The carcass was condemned.

DISCUSSION

None of the tests currently available for the diagnosis of bTB allow a perfectly accurate determination of *M. bovis* infection status of cattle. With 99.5% specificity [23] in TB-free cattle populations, skin tests remain the primary ante-mortem diagnostic tool for bTB. Bovine TB has been eradicated effectively from many countries including Germany following the implementation of programmes of regular tuberculin skin testing and removal of reactors, coupled with slaughterhouse surveillance for undetected infections, and a variety of accompanying measures, e.g. cattle movement restrictions. The deployment of

IGRA as an ancillary *in vitro* test has enhanced the detection of bTB-infected cattle (for a comprehensive overview of different ante-mortem diagnostic techniques for bTB, see de la Rua-Domenech *et al.* [23]).

Using primary skin testing and subsequent IGRA testing of reactor cattle, a case of presumptive exposure of livestock to *M. tuberculosis* in a dairy holding and an infected pig as the probable proximate source of the agent has been uncovered. Although no clinical signs indicative of bTB were detected, exposure of a significant number of cattle was revealed with as many as 55 animals showing delayed type hypersensitivity to bovine PPD. The *M. bovis*-derived PPD test antigens licenced in Germany for both skin and IGRA testing of cattle contain antigens also present in *M. tuberculosis*, which are likely to re-stimulate immune cells sensitized for MTC members other than *M. bovis*. Bovine infections with *M. tuberculosis* have been sporadically reported from European countries previously and, more recently, from developing countries [9, 12, 24–26]. Most infections in cattle seem to be attributed to intensive contact with humans in regions with high incidence of pulmonary TB in

humans [9–12, 27–30], even though the identity of human and bovine strains is only rarely established [13, 31]. Detection of bovine *M. tuberculosis* infections is hampered by the absence or weak expression of clinical symptoms and by SICCT responsiveness occurring only transiently [12, 25, 31]. With the disappearance of the infection source and/or presumptive clearance of the agent, reactivity to tuberculin vanishes quickly [9, 13], thus probably resulting in significant under-reporting. Viable *M. tuberculosis* bacteria could not be isolated in the context of this case. However, sufficient indication exists to conclude that the high number of reactor animals identified upon repeated testing of the herd resulted from exposure to *M. tuberculosis* as detected by PCR in the pig.

The majority of cattle in the herd under investigation were SICCT-negative during initial testing in September 2013. Within the following 2 months, the number of reactors markedly increased. This implies recent introduction and highly efficient spread of the causative agent on the premise or continuous presence of the source [32]. Other authors found the spread of *M. bovis* within cattle herds to be relatively inefficient, but other factors, such as the general health status of infected and/or contact animals or the type of animal husbandry, may facilitate disease transmission [17, 32]. In order to rapidly identify and remove the possible source of infection and considering the high susceptibility of cats towards *M. bovis*, a cat from the farm was also euthanized [33]. None of the cows nor the cat showed typical symptoms of TB and MTC members could neither be detected by PCR nor by bacterial culture, thus strongly arguing against the presence of *M. bovis* in that herd.

The PCR employed in Germany for official diagnosis targets two gene sequences, *IS1081* and *HELI*. Using two target genes instead of a single one can be assumed to enhance sensitivity and specificity of detection, which is of particular relevance for OTF countries. Subsequent species identification and molecular characterization of the pathogen causing the granulomas in the porcine lymph node was conducted by microarray-based spoligotyping. *M. tuberculosis* was identified and the pattern detected was assigned to the T1 family. This lineage has previously been identified in human samples from Germany and many other countries. The microarray possesses a superior sensitivity compared to conventional spoligotyping and was able to generate signal patterns directly from the tissue's DNA extract [20]. In other suspected bTB cases associated with *M. tuberculosis*,

the authors also failed to cultivate the causative organism [9, 11] or only succeeded after initial passage in a guinea pig [9]. Application of highly sensitive molecular typing methods may therefore help resolve more cases of livestock infections involving MTC members other than *M. bovis* and *M. caprae*.

Since the presence of *M. tuberculosis* on the farm became evident, dynamics of the immunoconversion observed in the cattle herd might best be explained by the presence of an unidentified human shedder with 'open' TB on the farm and in proximity to susceptible animals shortly before the nationwide survey started. Quantiferon® (Cellestis Ltd, Germany) testing carried out with the farmer, his family and several contact persons including contract workers and several other contact persons indeed led to the detection of four human reactors within the farmer's family. Although all four tested negative by X-ray examination, the possibility exists that these persons had experienced recent *M. tuberculosis* infection. The exact route of pathogen introduction remains obscure leaving the question unanswered whether the patients were source or target of bovine exposure to *M. tuberculosis*. Pigs are more susceptible to *M. tuberculosis* infection than cattle [33]. Thus, the possibility exists that infection of the fattening pig from a human source resulted in amplification of the agent including induction of histopathological lesions, shedding to the farm environment, as well as exposure and subsequent conversion of bovines and probably (even more likely) humans. The lack of positive culture results despite pathological lesions compatible with TB and positive PCR tests has been previously reported for wild boar [34, 35]. These authors assumed that active TB was no longer present at the time of investigation.

Based on observations during the German bTB eradication campaign in the 1960s, Schliesser expressed the opinion that bovine infections with *M. tuberculosis* were principally restricted to the formation of primary complexes with limited propagation of the agent in the host [36], which would result in absent or minor lesions detected at necropsy [13, 25]. Infections do not develop into open stages, thereby confining the infection to single animals and leading to the assumption that cattle represent dead-end hosts for *M. tuberculosis* [36]. Indeed, numerous reports on asymptomatic as well as clinical cases of bTB due to *M. tuberculosis* exist [9, 12, 13, 24–26, 30, 31] that merely agree in cattle-to-cattle transmission being uncommon. However, more recent cases with clinical and morphological symptoms, indistinguishable from 'classical' bTB, and even detection of

M. tuberculosis in bovine milk have been reported repeatedly from countries with a high TB burden [9, 25–30, 37]. These findings might indicate cattle as a potential source of *M. tuberculosis* transmission to humans. The TB incidence in humans is rising by approximately 1% per year [38] and a continuous spillover to cattle might even cause a better adaptation of *M. tuberculosis* to the bovine host [30]. This situation may create novel threats, especially in the light of multidrug-resistant strains being isolated from cattle [31]. Immunoconversion in cattle to bTB antigens, which cannot be immediately substantiated by direct detection of *M. bovis*, should not be dismissed easily as resulting from exposure to environmental mycobacteria and of no further relevance to animal and human health. Instead, public authorities, veterinarians and farmers should be aware of the risk resulting from introduction of *M. tuberculosis* into populations of susceptible ruminants. The consequences are not restricted to bTB monitoring and eradication programmes, but extend to safety issues concerning animal workers and even consumers. This is particularly relevant for countries regarded free from bTB.

In an attempt to clear the suspected herd breakdown, 55 cows were culled for diagnostic purposes, i.e. nearly a quarter of the whole herd referred to in this report. Those mycobacterial species known to interfere with bTB laboratory diagnosis [39], particularly *M. avium* and *M. kansasii*, were not found in this case. MTC members other than *M. bovis* and *M. caprae*, especially *M. tuberculosis*, are rarely considered as a differential diagnosis for bTB, particularly in countries with low prevalence of both bTB and human TB [31]. Results from the current investigation imply, however, that exposure of bovines to *M. tuberculosis* may result in high numbers of animals immunoconverting and thus responding to bTB test antigens. In skin testing campaigns, competent authorities may then face severe difficulties in interpreting test results and, depending on the wording of the legal requirements that apply, in decision making on the status of the affected herd. The German Regulation on bTB prescribes confirmation of cases involving positive or inconclusive SICCT or positive IGRA reactors, on the basis of bacterial culture of *M. bovis* or *M. caprae*, or detection of specific gene sequences. A positive result of intradermal tuberculin testing alone does not fulfil the case definition. According to the German Regulation on Reportable Animal Diseases (as of 17 April 2014), detection of TB in pigs is classified as reportable only, irrespective of the causative MTC species. In the present study,

pathological signs of mycobacteriosis and molecular detection of MTC found in one pig even on a farm with a significant number of reactor cattle did not suffice to justify the declaration of a bTB breakdown. Consequently, grading of the farm as a suspected outbreak had to be reversed after tissue samples from culled cattle were scored negative by PCR and culture and the remaining cattle proved SICCT negative at re-testing in March 2014. More importantly, epidemiological trace-back and trace-on investigations in bovine contact herds lacked a legal basis and, despite strong evidence for bTB at first, these investigations were – in the light of all-negative cattle samples and the proof of *M. tuberculosis* – merely focused on the human sector. Appropriate handling of such cases may require adaptations of national and international regulations to facilitate eradication measures and compensation payments. Considering the zoonotic potential of all MTC species and the relatively low likelihood of bovine infections with MTC members other than *M. bovis* and *M. caprae* being revealed, the spectrum of notifiable bovine infections should be extended to any MTC member. Furthermore, direct detection of MTC in other livestock species with epidemiological links to herds harbouring reactor cattle should be accepted as sufficient evidence to declare a bTB outbreak.

ACKNOWLEDGEMENTS

The authors thank Barbara Gamb for unsurpassed literature service. Matthias Contzen (CVUA Stuttgart) is acknowledged for *gyrB* analysis. We are grateful to Konrad Sachse for critically reading the manuscript.

The Hessian State Laboratory is supported by the Hessian Ministry for the Environment, Climate Change, Agriculture and Consumer Protection (HMUKLV).

DECLARATION OF INTEREST

None.

REFERENCES

1. Alexander KA, et al. Novel *Mycobacterium tuberculosis* complex pathogen, *M. mungi*. *Emerging Infectious Diseases* 2010; **16**: 1296–1299.
2. Singh SK, Verma R, Shah DH. Molecular fingerprinting of clinical isolates of *Mycobacterium bovis* and *Mycobacterium tuberculosis* from India by restriction

- fragment length polymorphism (RFLP). *Journal of Veterinary Science* 2004; **5**: 331–335.
3. **Miller M, Olea-Popelka F.** One health in the shrinking world: experiences with tuberculosis at the human-livestock-wildlife interface. *Comparative Immunology, Microbiology and Infectious Diseases* 2013; **36**: 263–268.
 4. **Panteix G, et al.** Pulmonary tuberculosis due to *Mycobacterium microti*: a study of six recent cases in France. *Journal of Medical Microbiology* 2010; **59**: 984–989.
 5. **van Ingen J, et al.** Characterization of *Mycobacterium orygis* as *M. tuberculosis* complex subspecies. *Emerging Infectious Diseases* 2012; **18**: 653–655.
 6. **Parsons SD, et al.** Novel cause of tuberculosis in meerkats, South Africa. *Emerging Infectious Diseases* 2013; **19**: 2004–2007.
 7. **Parsons S, et al.** Pulmonary infection due to the dassie bacillus (*Mycobacterium tuberculosis* complex sp.) in a free-living dassie (rock hyrax – *Procavia capensis*) from South Africa. *Tuberculosis (Edinburgh, Scotland)* 2008; **88**: 80–83.
 8. **Jahans K, et al.** Isolation of *Mycobacterium microti* from a male Charolais-Hereford cross. *Veterinary Record* 2004; **155**: 373–374.
 9. **Lesslie IW.** Tuberculosis in attested herds caused by the human type tubercle bacillus. *Veterinary Record* 1960; **72**: 218–224.
 10. **Krishnaswami KV, Mani KR.** *Mycobacterium tuberculosis* humanis causing zoonotic tuberculosis among cattle. *Indian Journal of Public Health* 1983; **27**: 60–63.
 11. **Smith IGN.** A herd breakdown due to *Mycobacterium tuberculosis*. *State Veterinary Journal* 1984; **38**: 40–44.
 12. **Steele JH.** Human tuberculosis in animals. In: Steele JH, ed. *CRC Handbook Series in Zoonoses, Section A: Bacterial, Rickettsial and Mycotic Diseases, vol 2*. Boca Raton, FL: CRC Press, Inc., 1980.
 13. **Ocepek M, et al.** Transmission of *Mycobacterium tuberculosis* from human to cattle. *Journal of Clinical Microbiology* 2005; **43**: 3555–3557.
 14. **Kaneene JB, Pfeiffer D.** Epidemiology of *Mycobacterium bovis*. In: Thoen CO, Steele JH, Gilsdorf MJ, eds. *Mycobacterium bovis Infection in Animals and Humans*. Oxford: Blackwell Publishing, 2006, pp. 34–48.
 15. **Meyn A.** Fighting bovine tuberculosis in the Federal Republic of Germany [in German]. *Monatshefte für Tierheilkunde* 1952; **4**: 510–526.
 16. **Domogalla J, et al.** Region of difference 4 in alpine *Mycobacterium caprae* isolates indicates three variants. *Journal of Clinical Microbiology* 2013; **51**: 1381–1388.
 17. **Probst C, et al.** Bovine tuberculosis: making a case for effective surveillance. *Epidemiology and Infection* 2011; **139**: 105–112.
 18. **Anon.** *Official Collection of Diagnostic Methods* [in German]. January 2014 edn. Riems: Friedrich-Loeffler-Institut, Bundesforschungsinstitut für Tiergesundheit, 2014.
 19. **Kamerbeek J, et al.** Simultaneous detection and strain differentiation of *Mycobacterium tuberculosis* for diagnosis and epidemiology. *Journal of Clinical Microbiology* 1997; **35**: 907–914.
 20. **Ruettger A, et al.** Rapid spoligotyping of *Mycobacterium tuberculosis* complex bacteria by use of a microarray system with automatic data processing and assignment. *Journal of Clinical Microbiology* 2012; **50**: 2492–2495.
 21. **Weisburg WG, et al.** 16S ribosomal DNA amplification for phylogenetic study. *Journal of Bacteriology* 1991; **173**: 697–703.
 22. **Kasai H, Ezaki T, Harayama S.** Differentiation of phylogenetically related slowly growing mycobacteria by their *gyrB* sequences. *Journal of Clinical Microbiology* 2000; **38**: 301–308.
 23. **de la Rua-Domenech R, et al.** Ante mortem diagnosis of tuberculosis in cattle: a review of the tuberculin tests, gamma-interferon assay and other ancillary diagnostic techniques. *Research in Veterinary Science* 2006; **81**: 190–210.
 24. **Berg S, et al.** The burden of mycobacterial disease in Ethiopian cattle: implications for public health. *PLoS ONE* 2009; **4**: e5068.
 25. **Srivastava K, et al.** Isolation of *Mycobacterium bovis* & *M. tuberculosis* from cattle of some farms in north India – possible relevance in human health. *Indian Journal of Medical Research* 2008; **128**: 26–31.
 26. **Thakur A, et al.** Detection of *Mycobacterium bovis* and *Mycobacterium tuberculosis* from cattle: possible public health relevance. *Indian Journal of Microbiology* 2012; **52**: 289–291.
 27. **Ameni G, Erkihun A.** Bovine tuberculosis on small-scale dairy farms in Adama Town, central Ethiopia, and farmer awareness of the disease. *Revue Scientifique et Technique (International Office of Epizootics)* 2007; **26**: 711–719.
 28. **Chen Y, et al.** Potential challenges to the stop TB plan for humans in China; cattle maintain *M. bovis* and *M. tuberculosis*. *Tuberculosis (Edinburgh, Scotland)* 2009; **89**: 95–100.
 29. **Fetene T, Kebede N, Alem G.** Tuberculosis infection in animal and human populations in three districts of Western Gojam, Ethiopia. *Zoonoses and Public Health* 2011; **58**: 47–53.
 30. **Mittal M, et al.** Evidence of presence of *Mycobacterium tuberculosis* in bovine tissue samples by multiplex PCR: possible relevance to reverse zoonosis. *Transboundary and Emerging Diseases* 2014; **61**: 97–104.
 31. **Romero B, et al.** Humans as source of *Mycobacterium tuberculosis* infection in cattle, Spain. *Emerging Infectious Diseases* 2011; **17**: 2393–2395.
 32. **McIlroy SG, Neill SD, McCracken RM.** Pulmonary lesions and *Mycobacterium bovis* excretion from the respiratory tract of tuberculin reacting cattle. *Veterinary Record* 1986; **118**: 718–721.
 33. **LoBue PA, Enarson DA, Thoen CO.** Tuberculosis in humans and animals: an overview. *International Journal of Tuberculosis and Lung Disease* 2010; **14**: 1075–1078.
 34. **Müller M, et al.** Detection of *Mycobacterium tuberculosis* complex in wild boars by PCR [in German]. *Tierärztliche Umschau* 2007; **62**: 140–143.
 35. **Capellmann C.** Epidemiological study for the detection of selected mycobacteria species among wild boars in Germany [in German]. Giessen: Justus-Liebig-University; 2011. 145 pp.

36. **Schliesser T.** *Mycobacterium*. In: Blobel H, Schliesser T, eds. *Handbook on Bacterial Infections in Animals* [in German], 1st edn. Stuttgart: Gustav Fischer Verlag, 1985, pp. 155–313
37. **Schliesser T.** Contribution to the epidemiology and allergic diagnosis of human tuberculosis in cattle [in German]. *Tierärztliche Umschau* 1958; **13**: 328–332.
38. **Smith RM, et al.** *Mycobacterium bovis* infection, United Kingdom. *Emerging Infectious Diseases* 2004; **10**: 539–541.
39. **Thacker TC, et al.** Isolation of mycobacteria from clinical samples collected in the United States from 2004 to 2011. *BMC Veterinary Research* 2013; **9**: 100.