

unsuccessful and can lead to the development of serious psychiatric conditions and diseases, most often to depression.

**Key words:** crisis, depression, crisis intervention, prevention.

## P0280

Analysis of suicides committed in Podgorica

L.R. Injac Stevovic<sup>1,2,3</sup>, O.S. Vukovic<sup>1,2,3</sup>, Z.V. Barac Otasevic<sup>1,2,3</sup>.  
<sup>1</sup> *Psychiatric Department, Medical Center of Ministry of Internal, Podgorica, Serbia and Montenegro* <sup>2</sup> *Institute of Psychiatry, Clinical Centre of Serbia, Belgrade, Serbia and Montenegro*  
<sup>3</sup> *Psychiatric Clinic, Clinical Hospital Center, Podgorica, Serbia and Montenegro*

This paper analyzes suicides committed in Podgorica during the 2000-2005 period divided according to gender, age and the way these have been performed. In 2000 we have 25 suicides on 100.000 inhabitants, and in next five years there is a growth in trend thus in 1999, there are 17.3 suicides on 100.000. Comparing to 1989 when suicide rate was 7.89, and 2003 rate which was 35.74, we can notice that significant social stresses, lead to a rise in rate of suicides so average rate for period of 2000-2005 is 28.29 which puts Podgorica into area with high suicide rate. The male/female ratio was 2.8:1. The largest number of people who killed themselves are between 35 and 44 years old. The way that people use to perform suicide is mostly by firearm (40.78%) and hanging themselves (38.03%) which is near to average numbers in the world. This research shows that there is significant rise in suicide rate. As a conclusion, on the basis of cited literature, the authors lined up all moments which can explain why have happen significance rise in suicide rate.

**Key words:** suicide, social stress, depression.

## P0281

Psychological distress among migrant patients admitted in acute psychiatric ward

V.C. Iversen<sup>1</sup>, A.E. Vaaler<sup>1</sup>, J.E. Berg<sup>2</sup>. <sup>1</sup> *St. Olavs Hospital, Østmarka Psychiatric Department, Trondheim, Norway*  
<sup>2</sup> *Lovisenberg Diakonal Hospital, Oslo, Norway*

The adaptation process of migrants has been considered to arise from the differences that may exist between the migrant's traditional cultural background and the cultural norms and values of the society of settlement. These differences are thought to be stressful, and may induce socio-cultural and psychological changes on the part of the migrant, some of which may be maladaptive.

The main purpose of this study is to determine the prevalence of psychological distress and socio-cultural adaptation among migrants who attend acute psychiatric wards. A second purpose is to examine whether or not acute psychiatric wards are used by refugees in situations of crises, like when they asylum application is turned down by the Norwegian authorities.

All immigrants, irrespective of nationality and ethnicity, admitted to acute psychiatric ward by St. Olav's hospital, were eligible to participate in the study. The sample studied included all patients admitted to the hospital from 2005-2008.

The instrument used was a self-administered questionnaire and a structured interview. The questionnaire includes questions on demographic variables, social, psychological and psychosomatic symptoms. In addition, the Beck Hopelessness Scale and Harvard Trauma Questionnaire (HTQ) were used. The data were also collected from the patients' records and from the hospital protocols.

The admission rates, including admission by coercion, length of hospital stay, diagnosis and type of treatment were also compared among the different migrants groups.

## P0282

Some aspects of mental health of population of Kazakhstan

B. Jarbussynova. *Kazak National Medical University, Almaty, Kazakhstan*

**Objective:** To find out distribution of levels of mental and psychologic health within a certain age group and to determine main parameters of correlation of the above mentioned health categories among the population of Kazakhstan. A large number of researchers make precise distinctions between definitions of mental and psychologic health: the first is recognized as absence of mental disorders (biologic criteria); the latter – as a whole complex of personal qualities ensuring satisfactory adaptation of a person (psychologic criteria).

**Methods:** 7968 respondents, aged 12 -55, underwent epidemiologic study with an aim to find out how levels of mental and psychologic health are correlated. Clinical-epidemiologic and clinical-sociologic methods were applied. There was used a special map of multilevel monitoring, which includes indicators and parameters, defining levels of mental and psychologic health.

**Results:** In general, the research clarified all the difference between conceptions of mental and psychologic health; besides analysis revealed relevantly deeper crisis condition of psychologic health of population of Kazakhstan as compared to mental, as well as more dynamic reaction of a mobile level of a psychologic functional activity (psychologic health) to social and economic changes of the last decades. It is necessary to elaborate differentiated psychohygienic and psychoprophylactic approaches to the analysed health categories with more attention to psychologic.

## P0283

Descriptive study of an inpatient sample from the brief hospitalization psychiatry unit at the University Hospital of Valladolid, Spain

M.D. Ortega<sup>1</sup>, M.L. Vargas<sup>2</sup>, N. Jimeno<sup>3</sup>. <sup>1</sup> *Department of Psychiatry, University Hospital, Valladolid, Spain* <sup>2</sup> *Department of Psychiatry, Hospital Santa Elena, Zamora, Spain* <sup>3</sup> *Department of Psychiatry, School of Medicine, Valladolid, Spain*

**Background and Aims:** To establish the main sociodemographic and clinical characteristics of a sample of inpatients admitted at the Brief Hospitalization Psychiatry Unit from the University Hospital of Valladolid, Spain during 2005.

**Methods:** Descriptive study of a sample of inpatients (n=160) admitted in 2005 by means of the discharge report. Sociodemographic variables include age, sex, civil status and both socioeconomic and education levels; clinical variables include diagnosis according ICD and DSM-IV criteria, substance use, duration of admission and pharmacological treatment.

**Results:** The type patient is a male between 31 and 40 years or a woman between 51 and 60 years, single, with a middle socioeconomic and education levels. Substance use is frequently found in male patients, mainly concerning several substances. Mean duration of admission was about 13 days. Women usually are diagnosed of affective disorders and men of adaptive disorders associated with substance use; schizophrenia also appears more frequently in men.