

## Some observations on the faecal carriage of mesophilic *Aeromonas* species in cows and pigs

S. J. GRAY<sup>1</sup> AND D. J. STICKLER<sup>2</sup>

<sup>1</sup> Public Health Laboratory Service, University Hospital of Wales, Heath Park, Cardiff CF4 4XW.

<sup>2</sup> School of Pure and Applied Biology, University of Wales College of Cardiff, P.O. Box 13, Cardiff CF1 3XF

(Accepted 22 June 1989)

### SUMMARY

Replicate faecal samples from healthy individual pigs and cows were examined for the presence of *Aeromonas* sp. over a 12-month period. Aeromonads were found to be minor components of the faecal flora, only 8.8% of 520 samples from pigs and 4.6% of 481 samples from cows proving positive. Isolation rates in both groups of animals were seasonal. *A. hydrophila* (62% of the isolates) was the predominant species in cows, followed by *A. caviae* (32%) and *A. sobria* (15%). This pattern was also recorded in the natural waters that the animals drank from during the period when the faecal carriage rate was at its highest. In pigs, *A. caviae* (59%) was more common than *A. hydrophila* (41%). *A. sobria* was not found in any of the pig-associated samples. It seems that cattle acquire their faecal aeromonads from drinking water. The source of the organisms in pigs is less clear.

### INTRODUCTION

The mesophilic species of the genus *Aeromonas* can be isolated from a wide range of aquatic habitats (1-4). They have been long recognized as primary pathogens of cold-blooded animals such as, amphibians, reptiles and fish (5-7). In humans they can act as opportunistic pathogens in debilitated or immunologically compromised patients (8-11). There have also been many reports of the isolation of *Aeromonas* spp. from cases of human gastro-enteritis when no other recognized enteric pathogen has been found (12-15). Although an attempt to produce enteric disease by administration of aeromonads to human volunteers failed (16) many strains of *Aeromonas* spp. have been shown to produce enterotoxins and other factors which could endow these organisms with the capability for enteropathogenicity (17-22).

The lack of information on the incidence of *Aeromonas* spp. in mammals other than man, prompted Gray (23) to search for them in the intestinal flora of cows, pigs, horses and sheep. Analysis of single samples from 459 animals revealed the presence of aeromonads in 21%, 9.6, 6.4 and 9.0% of the faecal samples respectively. As the reservoir of aeromonads in healthy livestock may represent a potential source of human infection it was decided to carry out a more detailed

study of the ecology of these organisms in the agricultural environment. The specific objectives were (a) to observe the faecal carriage rates in cohorts of cows and pigs over a 12-month period; (b) to establish the incidence of the organisms in the environment and the feed and water supplies of the animals over the same period and (c) to examine the relationship between the faecal and environmental isolates.

#### MATERIALS AND METHODS

##### *Sampling sites*

Two adjacent farms situated in the county of South Glamorgan provided the livestock that were sampled during this study, farm A, the pigs and farm B, the cows.

Pig breeding was the main activity on farm A, up to 2000 pigs and piglets being held at any one time. Sampling was facilitated by the farming practice of keeping gilts and sows (Large White-cross-Landrace) in individual Ministry of Agriculture approved metal cages during gestation and farrowing. The gestation or stall house contained a maximum of 112 sows and was separated from the farrowing houses both of which had a capacity of 36 sows. Once weaned in the farrowing house the piglets were put into rearing pens. The sows were then moved into communal pens in the serving area, containing up to five animals, where they were mated with the resident boars. There was thus a continual three house cycle; serving area – 4 weeks; stall house – 3 months; farrowing house – 4 weeks.

In the farrowing house the only water available to the pigs was a chlorinated supply, operated via snout valves in feeding troughs. The diet consisted of feed pellets. No separate water supply was available in the stall and serving areas, food and liquid being obtained from the piped supply of pig swill, the owner being a Ministry of Agriculture approved producer.

Farm B was a mixed farm which maintained a dairy herd of some 250 Fresian–Holstein cows. From November 1985 to the end of April 1986 the herd was kept indoors and fed on a mixed diet of silage and feed concentrates. The cows drank from galvanized steel water troughs supplied with chlorinated mains water. From 1 May 1986 cows were turned out to graze on pasture by day and housed at night. During the day the animals had access to water in drainage channels or 'reens' and by night to the chlorinated supply. From the end of May 1986 to the time when sampling ceased in October 1986 cows were permanently out to pasture and had access only to natural ree waters. Cows were fed concentrated feed nuts during milking throughout the year.

##### *Sampling procedures*

Faecal samples from pigs and cows were collected at monthly intervals from November 1985 to October 1986 inclusive, although pigs were sampled twice during November 1985. Forty sows and gilts and 40 cows were selected for sampling, pigs being identified by a numbered ear tag and cows by a number branded on their hind quarters. Whenever possible those animals originally selected were sampled at each visit, although this was not always possible. Faeces were collected from the individual pigs when they were located in the stall and farrowing houses, while rectal swabs were obtained from individuals in the

communal pens of the serving area. Rectal swabs were obtained from cows at the time of milking.

Water, feedstuffs, bedding materials and a variety of other environmental samples were collected throughout the study or as they became available to the animals. All samples were collected in appropriate sterile containers and examined within 6 h of sampling. At the beginning of the study a small number of milk samples were obtained from cows found excreting aeromonads at the previous sampling.

#### *Isolation protocol*

*Faeces.* Isolation and enrichment techniques were as previously described (23), using Xylose-Deoxycholate Citrate Agar (XDCA) and alkaline peptone water (APW).

*Waters: enumeration.* Serial tenfold dilutions of chlorinated and unchlorinated waters (100 ml) were examined by a membrane filtration technique. After filtration the membrane being applied to the surface of XDCA and incubated at 37 °C for 18 h. Membranes from undiluted waters were also enriched in APW.

*Straw and pasture washings.* Each sample of material (25 g) was washed in sterile distilled water (200 ml) by violent agitation. Washings were then treated as for water samples.

*Other samples.* To each sample (20 g) of concentrated cattle and piglet feed concentrate, silage, silage-feed concentrate mix, stall bedding, sawdust and soil was added 100 ml APW. One 20 ml volume of pig swill was also treated in this way, while a second was pre-treated with Tween 80 (20 ml, 1% v/v) in an attempt to disperse the fat in the swill, which could possibly have interfered with enrichment procedures. Small samples of dust and sludge which built up on window sills in the pig houses were enriched in 25 ml APW. Trapped flies were immersed in APW (25 ml) and crushed against the side of the container with a sterile swab. Milk samples (1 ml) were enriched in APW (25 ml) at 20 °C for 48 h. Enrichment broths were subcultured after 24 and 48 h onto XDCA and incubated at 37 °C overnight.

#### *Identification of isolates*

All morphologically distinct non-xylose fermenting colonies were subcultured onto horse blood agar for further testing. Presumptive identification of isolates as *Aeromonas* sp. was as described previously (23). At least 10% of the total number of non-xylose fermenting colonies on the membranes were always examined. APW enrichments from waters and washings shown to contain aeromonads by membrane enumeration were not processed further.

Confirmation of presumptive identification was by API 20E (API Laboratory Products, Basingstoke, Hants), a system capable of identification to the generic level. The identification strips were incubated at 30 °C for 5 days and examined daily. Indole and V-P production were tested after 48 h.

Isolates identified as *Aeromonas* spp. were speciated by the criteria of Popoff & Véron (24) as modified by Lee & Donovan (25). Additionally three enzymes assayed by the API ZYM kit (API Laboratory Products, Basingstoke, Hants.), chymotrypsin,  $\beta$ -glucuronidase and  $\beta$ -glucosidase, were used in speciation as these

Table 1. *The isolation of mesophilic Aeromonas sp. from the faeces of pigs and cows*

	Pigs	Cows
Number of animals sampled	71	86
Total number of samples	520	481
Number of animals colonized with <i>Aeromonas</i> sp.	35	20
Number (%) of samples producing <i>Aeromonas</i> sp.	46 (8.8)*	22 (4.6)

\* Difference in isolation rates from pigs and cows was significant ( $\chi^2 = 7.2$ , D.F. = 1,  $P < 0.01$ ).

Table 2. *A comparison of direct culture and enrichment for the isolation of Aeromonas sp.*

Number positive faeces	Number (%) of samples producing <i>Aeromonas</i> sp.					
	Direct culture	Direct culture only	Enrichment		Enrichment 24 + 48 h	Enrichment only
			24 h only	48 h only		
68	1 (1.5)	0 (0)	11 (16)	31 (46)	26 (38)	67 (98.5)

had been shown to be useful adjuncts to the more frequently used biochemical tests (26).

## RESULTS

### *Faecal isolation rates*

Over the 12 month-period faecal samples from 71 pigs and 86 cows were examined for mesophilic aeromonads. A summary of the isolation results is shown in Table 1. All stools examined exhibited the consistency associated with the healthy non-diarrhoeal animal. Aeromonads were isolated more frequently from pig (8.8%) than cow (4.6%) faeces and the difference in isolation rates was significant ( $\chi^2 = 7.2$ , D.F. = 1,  $P < 0.01$ ). Some of the 68 positive faecal samples produced more than one biotype of aeromonad and a total of 97 faecal isolates were collected.

The majority (98.5%) of faecal samples produced aeromonads only after enrichment (Table 2). In 46% of the positive samples the organism was only detected after 48 h enrichment. However, in 11 faecal samples which were positive after 24 h enrichment, aeromonads were not isolated on subsequent subculture, confirming the value of examining the enrichment broth both at 24 h and 48 h.

### *Seasonal variation in faecal isolation rates*

The seasonal variation in the isolation rates from faeces is presented in Figure 1. In cows the isolation rate was low throughout the year except for a summer peak in August. Isolation rates from pigs were generally higher than from cows, and reached a peak in December. Of the 35 pigs and 20 cows that were positive at some stage of the investigation, aeromonads were recovered in consecutive monthly samples from only 2 of the pigs and none of the cows.




Fig. 1. Seasonal variation in the faecal isolation rate of *Aeromonas* spp. from pigs and cows.

#### *Distribution of Aeromonas sp. in environmental samples*

The feed supplies of the pigs on farm A consistently failed to produce any aeromonads (Table 3). The main sources were the slurry of faeces diluted with urine and water and the chlorinated water supply. The ever present population of flies also harboured the organism. Aeromonads were recovered after enrichment from a single sample taken from the outside of a bale of straw, a further nine samples taken from the middle of bales were all negative. The 14 samples of sawdust used as bedding material in the farrowing houses were all negative, as were the nine dust swabs taken from windowsills and the bars of pens.

The results from farm B (Table 4) indicate that cows were similarly exposed to aeromonads present in the chlorinated water supply. When cows were turned out to graze they had access to natural drainage water held in drainage channels or reens, all samples of which contained *Aeromonas* spp. The organisms were also isolated from the pasture they ate and the soil beneath the pasture. *Aeromonas* spp. were present in the slurry found in the milking parlour, overwintering shed and farm yard. Stall bedding for the cows in the overwintering quarters comprised sawdust and straw, and was positive on a single occasion. However, the individual components, sampled before use, were always negative. One sample of freshly cut grass from the silage clamp proved positive.

Many of the 57 positive environmental samples produced more than one biotype of *Aeromonas* sp., and a total of 113 environmental isolates were collected.

#### *Enumeration of Aeromonas spp. in chlorinated water supplies serving farms A and B*

The results of the enumeration of *Aeromonas* spp. on the chlorinated water supplies to the animals are presented in Table 5. The mains supply to the farrowing houses were negative for aeromonads on 12 out of 15 samples. The three positive samples were all from the same farrowing house tap. Counts were either

Table 3. Incidence of *Aeromonas* spp. in environmental samples from farm A

	Type of sample										Totals
	Chlorinated water	Straw	Slurry	Flies	Swill	Concentrated feed	Piglet feed concentrate	Environmental dust swabs	Sawdust		
Number sampled	33	9	17	12	15	16	10	9	14		135
Number positive	6	1	8	3	0	0	0	0	0		18
Number of isolates	6	3	10	3	0	0	0	0	0		22
Isolation rate (%)	18	11	47	25	0	0	0	0	0		13

Table 4. Incidence of *Aeromonas* spp. in environmental samples from farm B

Type of sample	Number sampled	Number positive	Number of isolates	Isolation rate (%)
Chlorinated water	30	16	31	53
Reen water	12	12	42	100
Milk	4	1	3	25
Bedding	6	1	2	17
Slurry	5	4	5	80
Silage	4	1	1	25
Pasture	6	2	3	33
Soil from pasture	4	2	4	50
Flies	3	0	0	0
Straw	6	0	0	0
Teat water spray	1	0	0	0
Teat antibiotic spray	1	0	0	0
Concentrate feed	9	0	0	0
Feed mix (silage + nuts)	9	0	0	0
Sawdust	8	0	0	0
Totals	108	39	91	36

very low or aeromonads were only detected upon enrichment. The supply to the pens was negative on 15 of 18 occasions sampled. In two samples aeromonads were isolated only after enrichment of the membranes. The third positive sample, from which *Aeromonas* sp. was isolated in relatively high numbers, was taken from a supply to a feed trough which contained a broth of water and concentrated feed nuts. There was no indication that *Aeromonas* sp. contamination of the water supply to farm A was seasonal.

Three water samples were collected from farm B at each visit. One sample from each of two troughs in the building where the cattle were overwintered together with a sample direct from the mains supply of trough 1. Aeromonads were recovered by direct isolation from the main supply serving trough 1 during July and August. Enrichment was necessary in order to recover organisms from samples taken in February and April (Table 5). Water from trough 1 contained aeromonads throughout the sampling period, while the organisms were only recovered from 2 of the 10 samples from trough 2.

#### *Aeromonas* spp. in drainage water on farm B

The enumeration of aeromonads in natural drainage water held in reens (Fig. 2) clearly shows a seasonal relationship in the incidence of these organisms. There was a rise in numbers as the weather became warmer, culminating in the August peak. Thereafter, their numbers declined.

#### *Ecological relationship between faecal and environmental aeromonads*

Attempts were made to determine the source of the faecal isolates of aeromonads. The management of the pigs on farm A allowed an examination of the relationship of the faecal carriage at different stages of the 20-week breeding cycle. The distribution of positive faecal samples from sows in each of the three

Table 5. Colony-forming units of aeromonads in 100 ml chlorinated water samples over an 11-month period

Month sampled	Colony-forming units/100 ml water							
	Farm A (pigs)				Farm B (cows)			
	Mains tap FH1	Pen valve FH1	Mains tap FH2	Pen valve FH2	Mains outlet trough 1	Trough 1	Trough 2	
December	NS	ND(-)	NS	NS	NS	ND(+)	NS	
January	NS	0(-)	NS	NS	NS	242(+)	17(+)	
February	NS	0(+)	NS	NS	0(+)	16(+)	0(-)	
March	0(-)	NS	0(-)	0(-)	0(-)	9(+)	0(-)	
April	NS	0(-)	0(-)	0(-)	0(+)	775(+)	0(-)	
May	0(-)	0(-)	0(-)	0(-)	0(-)	86(+)	0(-)	
June	0(-)	0(-)	0(+)	0(-)	0(-)	0(+)	0(-)	
July	0(-)	0(-)	0(-)	0(-)	1(+)	0(+)	0(-)	
August	0(-)	0(-)	0(-)	0(+)	14(+)	0(+)	0(-)	
September	0(-)	0(-)	1(-)	0(-)	0(-)	0(+)	0(+)	
October	0(-)	346(+)	2(-)	0(-)	0(-)	0(-)	0(-)	

FH1, farrowing house 1; FH2, farrowing house 2; ND, not done; NS, not sampled; Figures, colony-forming units in 100 ml water; (+), (-), indicates success or failure of isolation by enrichment from material on the filter.




Fig. 2. Enumeration of aeromonads in drainage (reen) waters on farm B.

Table 6. Relationship of 46 positive pig faecal samples to site of sampling

Location of animals	Total faeces sampled	Number positive	Percentage positive	Proportion (%) of total positive samples
Stall house	278	10	3.6	22
Serving area	81	1	1.2	2
Farrowing houses	161	35	22	76




Fig. 3. Monthly isolation rates from pig faeces in relation to the three-house cycle.

sites in which they were housed is presented in Table 6, and the location of the positive pigs over the 12-month sampling period illustrated in Fig. 3. It is clear that pigs were more likely to excrete aeromonads in their faeces whilst in the farrowing houses than when situated in either of the other two locations. It was also noted that when the sows were in the farrowing houses their faeces was much softer in texture than normal.

From November 1985 to April 1986 (period A) cows were kept indoors with access to chlorinated water only. Their feed comprised a mixture of silage and feed concentrates. During May (period B) cows were turned out to graze by day but were housed at night. Water consumption was thus a mixture of chlorinated and natural drainage water and feed a mixture of grass, concentrates and silage. From June to the end of the study in October 1986 (period C) the cows were at pasture with feed nuts being offered during milking. The animals had access to natural drainage water only.

During May till October, aeromonads were enumerated in washings from grass cut from fields where cows were grazing, and results related to the total viable counts obtained on the DCA agar. During the 6 months of sampling aeromonads were only recovered after enrichment, and then only when total colony counts exceeded  $1.28 \times 10^6$  per 100 ml of washings. The majority (78%) of organisms




Fig. 4. Monthly isolation of aeromonads from cow faeces and cow drinking water.

recovered from the pasture washings were oxidative, oxidase-positive organisms, pseudomonad-like in nature.

Figure 4 attempts to relate water sources shown to contain aeromonads to faecal carriage rates in cows in each of the three different periods of farming practice. It can be seen that the numbers of aeromonads in the trough water never exceeded  $8 \times 10^2$  c.f.u.  $100 \text{ ml}^{-1}$  and this degree of contamination did not result in high rates of faecal carriage during period A. In the summer months however (period C) there was a strong association between the increase in numbers of aeromonads in the reen water and the faecal carriage of the organisms.

#### *Relationship of species to the source of isolation*

Table 7 shows the relationship of the three species of *Aeromonas* to the source of isolation. Of the 93 pig-associated isolates 41 (44%) proved to be *A. hydrophila* and 52 (56%) *A. caviae*. *A. sobria* was not isolated. Of the 117 cow-associated isolates *A. hydrophila* was more frequently found (74/117, 63%) than *A. caviae* (32/117, 27%) and *A. sobria* (11/117, 9.4%). The predominance of *A. hydrophila*

Table 7. Relationship of Aeromonas species to the source of isolation

Source of isolate	Number (%) of isolates from each source			Totals
	<i>A. hydrophila</i>	<i>A. sobria</i>	<i>A. caviae</i>	
Pig faeces	29 (41)	0 (0)	42 (59)	71
Cow faeces	16 (62)	4 (15)	6 (23)	26
Chlorinated water on farm A	3 (50)	0 (0)	3 (50)	6
Chlorinated water on farm B	24 (77)	0 (0)	7 (23)	31
Environmental samples from farm A	9 (56)	0 (0)	7 (44)	16
Environmental samples from farm B	10 (67)	1 (7)	4 (27)	15
Cow's milk	2 (67)	0 (0)	1 (33)	3
Reen water from farm B	22 (52)	6 (14)	14 (33)	42
Totals	115 (55)	11 (5)	84 (40)	210

(62%) over *A. caviae* (23%) and *A. sobria* (15%) in cow faecal isolates and the ratios of the three species seems to be reflected in the species make-up of cow environmental isolates. *A. sobria* was found only in cow faeces and the reen water they drank. *A. hydrophila* was the predominant isolate from both chlorinated (73%) and reen water supplies (52%), *A. caviae* being isolated less frequently (27% and 33% respectively). Overall, *A. hydrophila* was the most frequent isolate (54%) followed by *A. caviae* (40%) and *A. sobria* (5.2%).

#### DISCUSSION

In an earlier investigation of the incidence of the *A. hydrophila* group in the faeces of livestock, single samples from 123 cows and 115 pigs produced isolation rates of 21 and 9.6% respectively (23). The results from the present study (Table 1) show that over a 12-month period, aeromonads were isolated more frequently from pig (8.8%) than cow (4.6%) faeces. Invariably the aeromonads were only isolated from the faecal samples after enrichment in APW for 24 or 48 h (Table 2) confirming the value of this procedure (27, 28). Millership & Chattopadhyay (29) have reported that extending the enrichment beyond 18 h reduced the isolation rate of aeromonads from artificially seeded faeces. In the current study while it was observed that enrichment broths found to be positive at 24 h were not always positive at 48 h, overall, enrichment beyond 24 h increased the isolation rate by 82%.

The isolation of aeromonads from both animal groups exhibited a seasonal pattern (Fig. 1). The frequency of aeromonad isolation from cows was greatest in August (28%). Similar seasonal patterns have been noted in human faeces (13, 22, 30, 31). In contrast in pigs the frequency of isolation was greatest in December.

In the present study 35 of the 71 pigs and 20 of the 86 cows provided positive samples at some stage of the 12-month period. Examination of the replicated monthly samples from the same animals revealed that repeated isolation of aeromonads from consecutive samples was rare. In the case of pigs the organism

was isolated from just single samples from 26 of the animals. In the remaining nine animals where the organisms were recovered from more than one sample, in only two cases were isolations in consecutive months. All 20 of the cows produced aeromonads from the faeces in just single monthly samples. These results suggest that *Aeromonas* spp. are minor components of the faecal flora of normal cows and pigs. While faecal carriage of these organisms by individual animals seems to be intermittent, aeromonads could of course be present in such small numbers that they were detected only irregularly by the isolation procedures.

George and colleagues (32) reported that in humans, aeromonads persist in the faeces of individuals during diarrhoea, but disappear soon after its resolution. Some authors have suggested that while aeromonads compete poorly with the normal intestinal flora in a healthy gut, the changes in the bacterial populations in the diarrhoeic state allows them to compete successfully and survive (33–35). In the present study none of the animals seemed to be suffering from gastroenteritis at the time of sampling.

Examination of a variety of environmental samples from the farms revealed that the only identifiable major sources of the aeromonads were the chlorinated water supplies to the animals, the natural waters and materials that had been contaminated with animal faeces. On farm A aeromonads were not isolated from any of the samples of pig-swill or concentrated feed-stuffs (Table 3). The results from farm B (Table 4) again indicate the absence of the organism from the prepared feedstuffs. The single sample of silage that proved to be positive for aeromonads was taken when freshly cut grass had been added to the old silage stack. Aeromonads were not isolated from mature silage presumably because it is acidic. Pasture used for the grazing of cows was positive on 2 of 6 occasions that it was sampled, perhaps as a result of faecal contamination.

The presence of aeromonads in milk (Table 4) has been noted by other workers (33, 36, 37). This might result from faecal contamination of the milk whilst obtaining the sample by hand, particularly as later samples from the same cow and two others, obtained using aseptic precautions, proved to be negative.

The water provided for the pigs, a chlorinated supply to the farrowing houses, was only occasionally a source of aeromonads (Table 5). No seasonal pattern in isolations from the chlorinated supply was noted and numbers did not increase with an increase in ambient temperature. This is in contrast to previous studies which had demonstrated such relationships in the aquatic habitat (38, 39). It is thus possible that the presence of the organism in the water samples on farm A was a direct result of environmental contamination or contamination of the trough outlets by the pigs themselves. In this connection Figura & Marri (40) have shown carriage of aeromonads in buccal cavities of pigs and suggested that these organisms may contaminate food by this route. It is thus possible that water from the snout-operated valves was contaminated by oral secretions from the pigs.

Similarly on farm B, it may be that the cows were a source of the aeromonads found in the drinking water troughs. Trough 1, located close to the animal stalls and the more intensively used was also the more heavily contaminated (Table 5). In addition the numbers of aeromonads recovered from trough 1 were highest during the time the cows had access to this water source and declined sharply when the animals were put out to pasture in the summer.

Over the period April to October water samples were examined from the drainage channels (reens) on farm B, which were used by the cows as a source of drinking water when they were turned out into the fields for grazing. The results (Fig. 2) show clearly that the population of aeromonads rises sharply to a maximum in August and confirms the results of other studies on natural waters (2, 41).

Analyses of the faecal carriage rates in the animals were carried out in an attempt to explain them in terms of the local farming practices. On farm A, the pigs were put through a three location cycle of serving area, gestation pens and farrowing houses. Analysis of the data presented in Table 6 and Fig. 3 by  $\chi^2$  tests confirms a highly significant association between the frequency of faecal isolations and residence in the farrowing houses which was independent of the time of year ( $P < 0.001$ ). It is interesting to note the increased frequency of aeromonads during the farrowing period, a time of apparent stress for the sow. The necessity of stress related factors for systemic infection with aeromonads has been noted for fish (41, 42), cows (43), and rabbits (44).

In the case of cows, the maximum rate of faecal carriage occurred during the period in which the animals were continuously outdoors grazing on pasture and obtaining their water from the natural sources of the reens. This maximum isolation rate also coincided with highest recovery of aeromonads from the reen water (Fig. 4).

There is little published information regarding the incidence of the various species of aeromonads in animal stools. While Figura & Marri (40) found *A. hydrophila* (75%) to predominate over *A. caviae* (25%) and failed to isolate *A. sobria* from tongue swabs and caecal contents of slaughtered swine, other studies have collectively referred to isolates as *A. hydrophila* (23, 45-47).

The results presented in Table 7 show that the distribution of the three aeromonad species was clearly different in the two groups of animals. In the case of the pigs *A. caviae* was more frequently isolated from the faeces than *A. hydrophila* and *A. sobria* could not be found. In cow faeces *A. hydrophila* was the predominant species followed by *A. caviae* and *A. sobria*. This pattern of species distribution was also observed in the water from the land drainage channels on farm B, the source of drinking water for the animals when the faecal carriage rate was at its highest.

In general, the results of this study are consistent with the hypothesis that cattle acquire their faecal aeromonads from their drinking water. The source of the faecal aeromonads in pigs however was less obvious. The highest rate of aeromonad isolation from faeces occurred when the pigs were in the farrowing houses being fed on concentrates rather than swill and when their only source of water was a chlorinated supply which was rarely contaminated with aeromonads.

#### ACKNOWLEDGEMENTS

We would like to thank Dr C. D. Ribeiro for his constructive criticism in the preparation of this manuscript and Dr R. Newcombe for statistical analysis.

## REFERENCES

1. Hazen TC, Fliermans CB, Hirsch RP, Esch GW. Prevalence and distribution of *Aeromonas hydrophila* in the United States. *Appl Environ Microbiol* 1978; **36**: 731–8.
2. Seidler RJ, Allen DA, Lockman H, Colwell RR, Joseph SW, Daily OP. Isolation, enumeration, and characterization of *Aeromonas* from polluted waters encountered in diving operations. *Appl Environ Microbiol* 1980; **39**: 1010–8.
3. Kaper JB, Lockman H, Colwell RR. *Aeromonas hydrophila*: ecology and toxigenicity of isolates from an estuary. *J Appl Bacteriol* 1981; **50**: 359–77.
4. Millership SE, Chattopadhyay B. *Aeromonas hydrophila* in chlorinated water supplies. *J Hosp Infect* 1985; **6**: 75–80.
5. Emerson H, Norris C. 'Red-leg' – an infectious disease of frogs. *J Exp Med* 1905; **7**: 32–60.
6. Shotts, Jr. EB, Gaines JL, Martin L, Prestwood AK. *Aeromonas* induced deaths among fish and reptiles in an eutrophic inland lake. *J Am Vet Med Assoc* 1972; **161**: 603–7.
7. Hazen TC, Raker ML, Esch GW, Fliermans CB. Ultrastructure of red-sore lesion on largemouth bass (*Micropterus salmoides*): association of the ciliate *Epistylis* sp. and the bacterium *Aeromonas hydrophila*. *J Protozool* 1978; **25**: 351–5.
8. Davies WA II, Kane JG, Garagsi VF. Human *Aeromonas* infections: a review of the literature and a case report of endocarditis. *Medicine (Baltimore)* 1978; **57**: 267–77.
9. Wolff RL, Wiseman SL, Kitchens CS. *Aeromonas hydrophila* bacteraemia in ambulatory immunocompromised hosts. *Am J Med* 1980; **68**: 238–42.
10. Ampel P, Peter G. *Aeromonas* bacteraemia in a burn patient. *Lancet* 1981; **ii**: 987.
11. Cookson BD, Houang ET, Lee JV. The use of a biotyping system to investigate an unusual clustering of bacteraemias caused by *Aeromonas* species. *J Hosp Infect* 1984; **5**: 205–9.
12. Rahman AFMS, Willoughby JMT. Dysentery-like syndrome associated with *Aeromonas hydrophila*. *Br Med J* 1980; **281**: 976.
13. Burke V, Gracey M, Robinson J, Peck D, Beaman J, Bundell C. The microbiology of childhood gastroenteritis: *Aeromonas* species and other infective agents. *J Infect Dis* 1983; **148**: 68–74.
14. Janda JM, Bottone EJ, Skinner CV, Calcaterra D. Phenotypic markers associated with gastrointestinal *Aeromonas hydrophila* isolates from symptomatic children. *J Clin Microbiol* 1983; **17**: 588–91.
15. Holmberg SD, Schell WL, Fanning GR, et al. *Aeromonas* intestinal infections in the United States. *Ann Intern Med* 1986; **105**: 683–9.
16. Morgan DR, Johnson PC, Dupont HL, Satterwhite TK, Wood LV. Lack of correlation between known virulence properties of *Aeromonas hydrophila* and enteropathogenicity for humans. *Infect Immun* 1985; **50**: 62–5.
17. Sanyal SC, Singh SJ, Sen PC. Enteropathogenicity of *Aeromonas hydrophila* and *Plesiomonas shigelloides*. *J Med Microbiol* 1975; **8**: 195–8.
18. Ljungh A, Popoff M, Wadström T. *Aeromonas hydrophila* in acute diarrhoeal disease: Detection of enterotoxin and biotyping of strains. *J Clin Microbiol* 1977; **6**: 96–100.
19. Cumberbatch N, Gurwith MJ, Langston C, Sack RB, Bruton JL. Cytotoxic enterotoxin produced by *Aeromonas hydrophila*: Relationship of toxigenic isolates to diarrhoeal disease. *Infect Immun* 1979; **23**: 829–37.
20. Atkinson HM, Trust TJ. Haemagglutination properties and adherence ability of *Aeromonas hydrophila*. *Infect Immun* 1980; **27**: 938–46.
21. Daily OP, Joseph SW, Coolbough JC, et al. Association of *Aeromonas sobria* with human infection. *J Clin Microbiol* 1981; **13**: 769–77.
22. Agger WA, McCormick JD, Gurwith MJ. Clinical and microbiological features of *Aeromonas hydrophila*-associated diarrhoea. *J Clin Microbiol* 1985; **21**: 909–13.
23. Gray SJ. *Aeromonas hydrophila* in livestock: incidence, biochemical characteristics and antibiotic susceptibility. *J Hyg* 1984; **92**: 365–75.
24. Popoff MY, Véron MA. A taxonomic study of *Aeromonas hydrophila*–*Aeromonas punctata* group. *J Gen Microbiol* 1976; **94**: 11–22.
25. Lee JV, Donovan TJ. *Vibrio, Aeromonas and Plesiomonas*. In: Collins CH, Grange JH, eds. Isolation and identification of micro-organisms of medical and veterinary importance. Society for Applied Bacteriology, London: Academic Press, 1985: 13–33.

26. Gray SJ. Characterisation of *Aeromonas* sp. in the API ZYM system. *Med Lab Sci* 1987; **44**: 287–9.
27. Millership SE, Curnow SR, Chattopadhyay B. Faecal carriage rate of *Aeromonas hydrophila*. *J Clin Pathol* 1983; **36**: 920–3.
28. Robinson J, Beaman J, Wagener L, Burke V. Comparison of direct plating with the use of enrichment culture for isolation of *Aeromonas* spp. from faeces. *J Med Microbiol* 1986; **22**: 315–7.
29. Millership SE, Chattopadhyay B. Methods for the isolation of *Aeromonas hydrophila* and *Plesiomonas shigelloides* from faeces. *J Hyg* 1984; **92**: 145–52.
30. Janda JM, Bottone EJ, Reitano M. *Aeromonas* species in clinical microbiology: significance, epidemiology and speciation. *Diagn Microbiol Infect Dis* 1983; **1**: 221–8.
31. Dosso Par M, Duchassin M, Koua-Amian P, Kouakon K, Gnanagne P, Kotchi R. *Aeromonas hydrophila* in the Ivory Coast. Epidemiological aspects of biological and enteropathogenic role. *Bull Soc Pathol Exot* 1984; **77**: 17–27.
32. George WL, Nakata MM, Thompson J, White ML. *Aeromonas*-related diarrhoea in adults. *Arch Intern Med* 1985; **145**: 2207–11.
33. Turnbull PCB, Lee JV, Miliotis MD, et al. Enterotoxin production in relation to taxonomic grouping and source of isolation of *Aeromonas* species. *J Clin Microbiol* 1984; **19**: 175–80.
34. Agger WA. Diarrhoea associated with *Aeromonas hydrophila*. *Pediatr Infect Dis* 1986; **5**: 106–8.
35. Figura N, Marri L, Verdiani S, Ceccherini C, Barberi A. Prevalence, species differentiation, and toxigenicity of *Aeromonas* strains in cases of childhood gastro-enteritis and in controls. *J Clin Microbiol* 1986; **23**: 595–9.
36. Driessen FM, Stadhouders J. Suitability of four different media for the enumeration of pseudomonads in milk. *Neth Milk Dairy J* 1972; **26**: 91–9.
37. Denis F, Veillet-Pancet L. Essai de mise en évidence d'une activité protéolytique d'origine intracellulaire chez *Aeromonas hydrophila* LP50. *Can J Microbiol* 1984; **30**: 1190–2.
38. Rouf MA, Rigney MM. Growth temperatures and temperature characteristics of *Aeromonas*. *Appl Microbiol* 1971; **22**: 503–6.
39. Picard B, Gouillet Ph. Seasonal prevalence of nosocomial *Aeromonas hydrophila* infection related to aeromonas in hospital water. *J Hosp Infect* 1987; **10**: 152–5.
40. Figura N, Marri L. Isolation of *Aeromonas* species from animals. *Eur J Clin Microbiol* 1985; **4**: 354–5.
41. Hazen TC. Ecology of *Aeromonas hydrophila* in a South Carolina cooling reservoir. *Microbial Ecol* 1979; **5**: 179–95.
42. Haley R, Davis SP, Hyde JM. Environmental stress and *Aeromonas liquefaciens* in American and threadfin shad mortalities. *Prog Fish Cult* 1967; **29**: 193.
43. Wohlgemuth K, Pierce RL, Kirkbride CA. Bovine abortion associated with *Aeromonas hydrophila*. *J Am Vet Med Assoc* 1972; **160**: 1001–2.
44. Hibbs CM, Merker JW, Kruckeberg SM. Experimental *Aeromonas hydrophila* infection in rabbits. *Cornell Vet* 1971; **61**: 380–6.
45. Annapurna E, Sanyal SC. Enterotoxicity of *Aeromonas hydrophila*. *J Med Microbiol* 1977; **10**: 317–23.
46. Dubey RS, Sanyal SC. Enterotoxicity of *Aeromonas hydrophila*: skin responses and *in vivo* neutralization. *Zentralbl Bakteriol (Orig A)* 1978; **242**: 487–99.
47. Pitarangsi C, Echeverria P, Whitmire R et al. Entero-pathogenicity of *Aeromonas hydrophila* and *Plesiomonas shigelloides*: Prevalence among individuals with and without diarrhoea in Thailand. *Infect Immun* 1982; **35**: 666–73.