

6th International Workshop on Immunonutrition, 15th–17th October 2012, Palma de Mallorca

Human intestinal dendritic cells decrease cytokine release against *Salmonella* infection in the presence of *Lactobacillus paracasei* CNCM I-4034, a novel strain isolated from breast-fed newborns

M. Bermúdez-Brito¹, S. Muñoz-Quezada¹, C. Gómez-Llorente¹, E. Matencio², M. J. Bernal²,
F. Romero² and A. Gil¹

¹Department of Biochemistry and Molecular Biology II, Institute of Nutrition and Food Technology “José Mataix”, Biomedical Research Centre, University of Granada, Armilla, Granada, Spain and ²Hero Institute for Infant Nutrition, Hero Spain, Alcantarilla, Murcia, Spain

Dendritic cells (DCs) are antigen-presenting cells that are involved in immunity and tolerance. Innate pattern-recognition receptors, such as Toll-like receptors, play a crucial role in the host recognition of probiotics. Signalling via these receptors influences the chemokine and cytokine production, proteins that are essential in regulating immune response. Monocyte-derived dendritic cells (MoDCs) and murine DCs are different from human gut DCs; therefore, in this study, we used human DCs harvested from umbilical cord blood CD34+ progenitor cells, which are similar to the lamina propria DCs in the gut. In the present work, we investigated the immunomodulatory properties of a novel strain, *Lactobacillus (L.) paracasei* CNCM I-4034, isolated from exclusively breast-feeding infant feces, on human DCs against *Salmonella (S.) typhimurium* CECT 4594. Human intestinal DCs were directly challenged by addition of *L. paracasei* CNCM I-4034, *S. typhimurium* CECT 4594 or both. After 4 hours incubation, the medium was replaced with a new one, containing antibiotics and cytokines. After 20 hours, culture supernatants were collected for cytokine analysis. IFN- γ , IL-1 β , IL-6, IL-8, IL-10, IL-12p40, IL-12p70, TNF- α , MCP-1(CCL2), MIP-1 α (CCL3), RANTES (CCL5), MDC (CCL22), IP-10 (CXCL10) and TGF- β were measured by immunoassay, with a MILLIplex kit using the Luminex 200 system based in the xMap technology. Differences between treatments were assessed by the U Mann Whitney test. We found that DCs decreased the secretion of the pro-inflammatory cytokines, such as IL-6 and IL-8, and chemokines, like MIP-1 α and RANTES, in response to stimulation with the probiotic and *S. typhimurium*. Interestingly, *L. paracasei* was a potent inducer of TGF- β 2 secretion. In conclusion, *L. paracasei* CNCM I-4034 appears to be a promising modulator of immune system which is able to activate dendritic cells and reduce pathogen-induced inflammation.