

The origin of the Galactic center diffuse X-ray emission investigated by near-infrared imaging and polarimetric observations

Shogo Nishiyama¹, Kazuki Yasui², Tetsuya Nagata²,
Tatsuhito Yoshikawa², Hideki Uchiyama³ and Motohide Tamura¹

¹National Astronomical Observatory of Japan, Mitaka, Tokyo 181-8588, Japan
email: shogo.nishiyama@nao.ac.jp

²Department of Astronomy, Kyoto University, Kyoto 606-8502, Japan

³Science Education, Faculty of Education, Shizuoka University, Shizuoka 422-8529, Japan

Abstract. The origin of the Galactic center diffuse X-ray emission (GCDX) is still under intense investigation. We have found a clear excess in a longitudinal GCDX profile over a stellar number density profile in the nuclear bulge region, suggesting a significant contribution of diffuse, interstellar hot plasma to the GCDX. We have estimated that contributions of an old stellar population to the GCDX are $\sim 50\%$ and $\sim 20\%$ in the nuclear stellar disk and nuclear star cluster, respectively. Our near-infrared polarimetric observations show that the GCDX region is permeated by a large scale, toroidal magnetic field. Together with observed magnetic field strengths in nearly energy equipartition, the interstellar hot plasma could be confined by the toroidal magnetic field.

Keywords. Galaxy: center, magnetic fields, polarization

1. Introduction

Past X-ray observations have revealed the presence of a diffuse 6.7 keV emission from highly ionized, He-like Fe ions toward the Galactic center (e.g., Koyama *et al.* 1989). The line emission and an associated continuum component, called the Galactic center diffuse X-ray emission (GCDX), resembles the Galactic ridge diffuse X-ray emission (GRXE). For the GRXE, more than 80% of the diffuse emission has been claimed to be resolved into point sources (Revnivtsev *et al.* 2009), suggesting faint X-ray point sources in origin.

However, the origin of the GCDX, in particular its very hot component with a temperature of $kT \sim 7$ keV, is more puzzling. Only 10 – 40% of the GCDX has been claimed to be resolved into faint point sources so far (e.g., Muno *et al.* 2004). The plasma temperature of the GCDX is likely to be systematically higher than the GRXE (Yamauchi *et al.* 2009). These results indicate a different origin of the GCDX from the GRXE.

Another idea for the origin of the GCDX is a diffuse, interstellar hot plasma (e.g., Koyama *et al.* 1989). In this case, the spatial distribution of the GCDX is expected to be different from that of stellar populations. Hence we have constructed a stellar number density map of the Galactic center region from new near-infrared (NIR) imaging observations. Here we summarize results of the NIR observations, and provide additional evidence for the hypothesis that the GCDX arises from a truly diffuse hot plasma.

We also show results of our NIR polarimetric survey to reveal magnetic field configuration in the Galactic center region. The results provide strong evidence for a large-scale toroidal magnetic field configuration in the GCDX region, implying a magnetic field confinement of the interstellar hot plasma.

2. Stellar number density profile

The central region of our Galaxy, $|l| \lesssim 3.0^\circ$ and $|b| \lesssim 1.0^\circ$ was observed from 2002 to 2004 using the NIR camera SIRIUS on the 1.4 m telescope IRSF. After removing foreground/background sources, we carry out an extinction correction for detected stars (Nishiyama *et al.* 2006), and obtain an extinction-corrected $K_{S,0}$ -band magnitude, $K_{S,0}$ for each star. Taking into account completeness limits, we construct a stellar number density map using stars with $K_{S,0} < 10.5$. For more detail, see Nishiyama *et al.* (2013).

The profile of the 6.7 keV line emission along the Galactic plane measured by *Suzaku* (Koyama *et al.* 2007, Uchiyama *et al.* 2011) clearly show an excess at the nuclear bulge (NB) region ($|l| \lesssim 1^\circ$) over the stellar number density profile (Figure 1, top panel). The two profiles are over-plotted and scaled to have the same values at $1.5^\circ < |l_*| < 2.8^\circ$ (l_* denotes the angular distance from Sgr A* along the Galactic longitude).

The majority of faint X-ray ($L_{2-10\text{ keV}} < 10^{30} \text{ erg s}^{-1}$) sources which have not been resolved but are expected to contribute to the GCDX are old binary systems such as cataclysmic variables. To investigate the contribution of the point sources, especially of the old stellar population detectable in our observations, we construct a longitudinal profile for the ratio of the stellar number density to the 6.7 keV emission (Figure 1, bottom panel). When the profile is scaled to be unity at the positions of normalization, $1.5^\circ < |l_*| < 2.8$, the ratios are $\sim 0.6 - 0.7$ and ~ 0.3 in the nuclear stellar disk (NSD) and the nuclear star cluster (NSC), respectively. Considering different star formation histories in the NSD, NSC, and the Galactic bulge (GB), the contributions of the old stellar population to the GCDX are estimated to be ~ 0.5 and ~ 0.2 for the NSD and NSC, respectively, in the assumption that the contribution of point sources in the GB is 100%.

The difference of the longitudinal profiles (Figure 1, top panel) is difficult to be explained by a variation of an emissivity of the point sources. In Nishiyama *et al.* (2013), we show that different initial mass functions, binary fractions, and star formation histories in the NSD and NSC from the GB cannot explain the difference in the longitudinal profiles.

Another factor which can change the emissivity is a Fe abundance. A higher Fe abundance of stars in the NB region than the GB leads to a higher X-ray emissivity. However, the 6.7 keV emission is ~ 1.6 times larger than the stellar number density in the NSD when they are shifted to have the same value at $|l_*| \sim 2^\circ$. It means that the required gradient of the Fe abundance is $\sim 60\%/2^\circ \sim 20\%/100 \text{ pc}$. This gradient is much larger than that in the inner Galaxy; an observed gradient of the Fe abundance at the inner Galaxy is $\sim 0.06 \text{ dex/kpc}$, which corresponds to only a few $\%/100 \text{ pc}$ (González-Fernández *et al.* 2008). In addition, the stellar Fe abundance in the Galactic center is not so high. Recent observations show a nearly solar Fe abundance even for young stars (e.g., Davies *et al.* 2009, Najarro *et al.* 2009). Thus the abundance for old stars which could be the origin of the GCDX is expected to be lower than the solar Fe abundance.

3. Magnetic field configuration

The most puzzling aspect of the GCDX is its high temperature. Since the $kT \approx 7 \text{ keV}$ plasma is too hot to be gravitationally bound, it requires a huge energy source without the confinement of the plasma. One idea to address this energetic issue is a confinement by magnetic fields (MFs; Makishima 1994). If a large-scale toroidal MF is developed and sustained, and the MF is strong enough for nearly energy equipartition with the plasma of $kT \sim 7 \text{ keV}$, the GCDX could be almost confined within the NB.




Figure 1. Top: Longitudinal profiles of the stellar number density (red) and the 6.7 keV emission (blue; Koyama *et al.* 2007; Uchiyama *et al.* 2011). The region in the Galactic bulge, $1.5^\circ < |l_*| < 2.8^\circ$, is used to scale the the 6.7 keV emission profile to have the same value as the stellar number density. **Bottom:** Longitudinal profile of the ratio of the stellar number density to the 6.7 keV emission, scaled to be unity at the position for normalization. This profile represents a contribution of point sources, traced by our NIR observations, to the GCDX in the assumption that the contribution of truly diffuse hot plasma is negligible at the position for normalization (i.e., the Galactic ridge region). [A COLOR VERSION IS AVAILABLE ONLINE.]

To determine the large-scale interstellar MF configuration, we have carried out NIR polarimetric observations for $|l| \lesssim 1.5^\circ$ and $|b| \lesssim 1.0^\circ$ using IRSF and a NIR polarimetric imager SIRPOL from 2006 to 2010. Comparing the polarization between stars distributed further and closer side in the Galactic center, we obtain polarization originating from magnetically aligned dust grains in the Galactic center (for more detail, see Nishiyama *et al.* 2009, 2010). The polarized angle traces the MF direction in the Galactic center projected onto the sky (Figure 2).




Figure 2. Near-infrared mosaic image of the Galactic center region covering $3.0^\circ \times 2.0^\circ$ in the Galactic coordinate, taken with the IRSF telescope and NIR camera SIRIUS. The three NIR bands are J (blue, $1.25\mu\text{m}$), H (green, $1.63\mu\text{m}$), and K_S (red, $2.14\mu\text{m}$). Observed polarization angles for the Galactic center components in the K_S band are also plotted with white bars whose length indicates the degree of polarization. [A COLOR VERSION IS AVAILABLE ONLINE.]

The obtained polarization map (Figure 2) suggests a large-scale toroidal MF configuration in the NB. The histogram of the MF directions at $|b| < 0.4^\circ$ has a clear peak at 90° (see Figure 5 in Nishiyama *et al.* 2013) which is the direction parallel to the Galactic plane. At high Galactic latitude ($|b| \gtrsim 0.4^\circ$), the fields are nearly perpendicular to the plane, i.e., poloidal configuration. This suggests a transition of the large-scale configuration, and such a transition can be naturally explained by the time evolution of MFs. The transition region, $|b| \sim 0.3^\circ$ to 0.4° , is in good agreement with the scale height of the 6.7 keV emission, 0.27° (Uchiyama *et al.* 2013).

If the plasma were not confined, it would be rushing out of the Galactic plane vertically as a Galactic wind. The escape velocity, typically several hundred km s^{-1} , is smaller than the sound speed of the 7 keV hot plasma of $> 1000 \text{ km s}^{-1}$. Assuming the gas flows out from the X-ray emitting region at the sound speed, the escape timescale is $\sim 4 \times 10^4 \text{ yr}$ (Belmont *et al.* 2005), requiring a huge energy input to sustain the hot plasma. If the plasma is confined magnetically, and there is no other cooling mechanism, the hot plasma only cools by radiation with a timescale of $10^7 - 10^8 \text{ yr}$ (Muno *et al.* 2004), several orders of magnitude longer than the escape timescale. This would reduce the required energy input and thus relax the energetic problem.

References

- Belmont, R., *et al.* 2005, *ApJ* 631, L53
 Davies, B., *et al.* 2009, *ApJ* 694, 46
 González-Fernández, C., *et al.* 2008, *A&A* 479, 131

- Koyama, K., *et al.* 1989, *PASJ* 59, 245
Koyama, K., *et al.* 2007, *Nature* 339, 603
Makishima, K., 2006, ASCA Observations of Galaxies. In F. Makino and T. Ohashi (Eds.), *New Horizon of X-Ray Astronomy. First Results from ASCA*, pp. 171.
Muno, M. P., *et al.* 2004, *ApJ* 613, 326
Najarro, F., *et al.* 2009, *ApJ* 691, 1816
Nishiyama, S., *et al.* 2006, *ApJ* 638, 839
Nishiyama, S., *et al.* 2009, *ApJ* 690, 1648
Nishiyama, S., *et al.* 2010, *ApJ*, 722, L23
Nishiyama, S., *et al.* 2013, *ApJ* 769, L28
Revnivtsev, M., *et al.* 2009, *Nature* 458, 1142
Uchiyama, H., *et al.* 2011, *PASJ* 63, 903
Uchiyama, H., *et al.* 2013, *PASJ* 65, 19
Yamauchi, S., *et al.* 2009, *PASJ* 61, 225