



COMPOSITIO MATHEMATICA

The homotopy theory of polyhedral products associated with flag complexes

Taras Panov and Stephen Theriault

Compositio Math. **155** (2019), 206–228.

[doi:10.1112/S0010437X18007613](https://doi.org/10.1112/S0010437X18007613)



FOUNDATION
COMPOSITIO
MATHEMATICA



LONDON
MATHEMATICAL
SOCIETY
EST. 1865



The homotopy theory of polyhedral products associated with flag complexes

Taras Panov and Stephen Theriault

ABSTRACT

If K is a simplicial complex on m vertices, the flagification of K is the minimal flag complex K^f on the same vertex set that contains K . Letting L be the set of vertices, there is a sequence of simplicial inclusions $L \rightarrow K \rightarrow K^f$. This induces a sequence of maps of polyhedral products $(\underline{X}, \underline{A})^L \xrightarrow{g} (\underline{X}, \underline{A})^K \xrightarrow{f} (\underline{X}, \underline{A})^{K^f}$. We show that Ωf and $\Omega f \circ \Omega g$ have right homotopy inverses and draw consequences. For a flag complex K the polyhedral product of the form $(\underline{CY}, \underline{Y})^K$ is a co- H -space if and only if the 1-skeleton of K is a chordal graph, and we deduce that the maps f and $f \circ g$ have right homotopy inverses in this case.

1. Introduction

The purpose of this paper is to investigate the homotopy theory of polyhedral products associated with flag complexes. Polyhedral products have received considerable attention recently as they unify diverse constructions from several seemingly separate areas of mathematics: toric topology (moment-angle complexes), combinatorics (complements of complex coordinate subspace arrangements), commutative algebra (the Golod property of monomial rings), complex geometry (intersections of quadrics), and geometric group theory (Bestvina–Brady groups).

To be precise, let K be a simplicial complex on the vertex set $[m] = \{1, 2, \dots, m\}$. For $1 \leq i \leq m$, let (X_i, A_i) be a pair of pointed CW -complexes, where A_i is a pointed CW -subcomplex of X_i . Let $(\underline{X}, \underline{A}) = \{(X_i, A_i)\}_{i=1}^m$ be the sequence of pairs. For each simplex $\sigma \in K$, let $(\underline{X}, \underline{A})^\sigma$ be the subspace of $\prod_{i=1}^m X_i$ defined by

$$(\underline{X}, \underline{A})^\sigma = \prod_{i=1}^m Y_i, \quad \text{where } Y_i = \begin{cases} X_i & \text{if } i \in \sigma, \\ A_i & \text{if } i \notin \sigma. \end{cases}$$

The *polyhedral product* determined by $(\underline{X}, \underline{A})$ and K is

$$(\underline{X}, \underline{A})^K = \bigcup_{\sigma \in K} (\underline{X}, \underline{A})^\sigma \subseteq \prod_{i=1}^m X_i.$$

For example, suppose each A_i is a point. If K is a disjoint union of m points, then $(\underline{X}, *)^K$ is the wedge $X_1 \vee \dots \vee X_m$, and if K is the standard $(m-1)$ -simplex, then $(\underline{X}, *)^K$ is the product $X_1 \times \dots \times X_m$.

The combinatorics of K informs greatly on the homotopy theory of $(\underline{X}, \underline{A})^K$. One notable family of simplicial complexes is the collection of flag complexes. A simplicial complex K is flag

Received 11 October 2017, accepted in final form 14 September 2018, published online 23 November 2018.

2010 Mathematics Subject Classification 55P35 (primary), 05E45 (secondary).

Keywords: polyhedral product, moment-angle complex, flag complex.

This journal is © [Foundation Compositio Mathematica](http://www.compositio-mathematica.org/) 2018.

if any set of vertices of K which are pairwise connected by edges spans a simplex. Flag complexes are important in graph theory, where they are referred to as clique complexes, in the study of metric spaces, where they are referred to as Rips complexes, and in geometric group theory, where they are referred to as Gromov’s no- Δ complexes.

The *flagification* of K , denoted K^f , is the minimal flag complex on the same set $[m]$ that contains K . We therefore have a simplicial inclusion $K \rightarrow K^f$. For example, the $(m - 1)$ -simplex Δ^{m-1} , consisting of all subsets of $[m]$, is flag, while its boundary $\partial\Delta^{m-1}$, consisting of all proper subsets of $[m]$, is flag only for $m = 2$. The flagification of $\partial\Delta^{m-1}$ with $m > 2$ is Δ^{m-1} . An m -*cycle* (the boundary of an m -gon) is flag whenever $m > 3$.

The main result of the paper is the following.

THEOREM 1.1. *Let K be a simplicial complex on the vertex set $[m]$, let K^f be the flagification of K , and let L be the simplicial complex given by m disjoint points. Let $(\underline{X}, \underline{A}) = \{(X_i, A_i)\}_{i=1}^m$ be a sequence of pairs of pointed CW-complexes, where A_i is a pointed CW-subcomplex of X_i . Let $(\underline{X}, \underline{A})^L \xrightarrow{g} (\underline{X}, \underline{A})^K \xrightarrow{f} (\underline{X}, \underline{A})^{K^f}$ be the maps of polyhedral products induced by the maps of simplicial complexes $L \rightarrow K \rightarrow K^f$. Then the following hold:*

- (a) *the map Ωf has a right homotopy inverse;*
- (b) *the composite $\Omega f \circ \Omega g$ has a right homotopy inverse.*

In particular, consider the special case when each A_i is a point. Write $(\underline{X}, *)$ for $(\underline{X}, \underline{A})$ and notice that $(\underline{X}, *)^L = X_1 \vee \cdots \vee X_m$. If K is a flag complex on the vertex set $[m]$, then the simplicial map $L \rightarrow K$ induces a map $f : X_1 \vee \cdots \vee X_m = (\underline{X}, *)^L \rightarrow (\underline{X}, *)^K$. By Theorem 1.1, Ωf has a right homotopy inverse. That is, $\Omega(\underline{X}, *)^K$ is a retract of $\Omega(X_1 \vee \cdots \vee X_m)$. This informs greatly on the homotopy theory of $\Omega(\underline{X}, *)^K$ since the homotopy type of $\Omega(X_1 \vee \cdots \vee X_m)$ has been well studied; in particular, in the special case when each X_i is a suspension the Hilton–Milnor theorem gives an explicit homotopy decomposition of the loops on the wedge. Theorem 1.1 also greatly generalizes [GPTW16, Theorem 5.3], which stated that such a retraction exists in the special case when each $X_i = \mathbb{C}P^\infty$ provided spaces and maps have been localized at a prime $p \neq 2$.

Theorem 1.1 can be improved in certain cases. In §6 we consider polyhedral products of the form $(\underline{CY}, \underline{Y})^K$, where CY is the cone on Y , and identify the class of flag complexes K for which $(\underline{CY}, \underline{Y})^K$ is a co- H -space. As a corollary, we obtain conditions that allow for a delooping of the statement of Theorem 1.1. In §7 we relate Theorem 1.1 to Whitehead products. First, we consider polyhedral products of the form $(\underline{X}, *)^K$ with flag K whose 1-skeleton is a chordal graph, and obtain a generalization of Porter’s description of the homotopy fiber of the inclusion of an m -fold wedge into a product in terms of Whitehead brackets. Second, we consider the loop space $\Omega(\underline{S}, *)^K$ on a polyhedral product formed from spheres for an arbitrary flag complex K , and obtain a generalization of the Hilton–Milnor theorem.

2. Combinatorial preparation

This section records the combinatorial information that will be needed. We begin with some definitions. Let K be an abstract *simplicial complex* on the set $[m] = \{1, 2, \dots, m\}$, i.e., K is a collection of subsets $\sigma \subseteq [m]$ such that for any $\sigma \in K$ all subsets of σ also belong to K . We refer to $\sigma \in K$ as a *simplex* (or a *face*) of K and denote by $|\sigma|$ the number of elements in σ . We always assume that the empty set \emptyset belongs to K . We do not assume that K contains all one-element

subsets $\{i\} \subseteq [m]$. We refer to $\{i\} \in K$ as a *vertex* of K , and refer to $\{i\} \notin K$ as a *ghost vertex*. We say that K is a simplicial complex *on the vertex set* $[m]$ when there are no ghost vertices.

Let K be a simplicial complex on the set $[m]$. For a vertex $v \in K$, the *star*, *restriction* (or *deletion*) and *link* of v are the subcomplexes:

$$\begin{aligned} \text{star}_K(v) &= \{\tau \in K \mid \{v\} \cup \tau \in K\}; \\ K \setminus v &= \{\tau \in K \mid \{v\} \cap \tau = \emptyset\}; \\ \text{link}_K(v) &= \text{star}_K(v) \cap K \setminus v. \end{aligned}$$

Throughout the paper we follow the convention of regarding $\text{star}_K(v)$ as a simplicial complex on the same set $[m]$ as K , while regarding $K \setminus v$ and $\text{link}_K v$ as simplicial complexes on the set $[m] \setminus v$. This implies that $\text{star}_K(v)$ and $\text{link}_K v$ may have ghost vertices even if K does not.

The *join* of two simplicial complexes K_1, K_2 on disjoint sets is the simplicial complex

$$K_1 * K_2 = \{\sigma_1 \cup \sigma_2 \mid \sigma_i \in K_i\}.$$

From the definitions, it follows that $\text{star}_K(v)$ is a join,

$$\text{star}_K(v) = \{v\} * \text{link}_K(v),$$

and there is a pushout

$$\begin{array}{ccc} \text{link}_K(v) & \longrightarrow & \text{star}_K(v) \\ \downarrow & & \downarrow \\ K \setminus v & \longrightarrow & K \end{array} \tag{1}$$

A *non-face* of K is a subset $\omega \subseteq [m]$ such that $\omega \notin K$. A *missing face* (a *minimal non-face*) of K is an inclusion-minimal non-face of K , that is, a subset $\omega \subseteq [m]$ such that ω is not a simplex of K , but every proper subset of ω is a simplex of K . A ghost vertex is therefore a missing face consisting of one element. Denote the set of missing faces of K by $\text{MF}(K)$. For a subset $\omega \subseteq [m]$, let $\partial\omega$ denote the collection of proper subsets of ω . Observe that $\omega \in \text{MF}(K)$ if and only if $\omega \notin K$ but $\partial\omega \subseteq K$.

A simplicial complex K on the set $[m]$ is called a *flag complex* if each of its missing faces consists of at most two elements. Equivalently, K is flag if any set of vertices of K which are pairwise connected by edges spans a simplex. Every flag complex K is determined by its 1-skeleton K^1 , and is obtained from the graph K^1 by filling in all cliques (complete subgraphs) by simplices.

LEMMA 2.1. *Let K be a flag complex on the set $[m]$ and let v be a vertex of K . If $\omega \in \text{MF}(\text{link}_K(v))$ and $|\omega| \geq 2$, then $\omega \in \text{MF}(K \setminus \{v\})$.*

Proof. Suppose not. Then there is a missing face ω of $\text{link}_K(v)$ with $\omega \in K \setminus \{v\}$ and $|\omega| \geq 2$. Therefore, $\partial\omega \subseteq \text{link}_K(v)$ but $\omega \notin \text{link}_K(v)$. Since $\omega \in K \setminus \{v\}$, we also have $\omega \in K$. On the other hand, as $\text{star}_K(v) = \text{link}_K(v) * \{v\}$, we have $\partial\omega * \{v\} \subseteq \text{star}_K(v)$, and so $\partial\omega * \{v\} \subseteq K$. Therefore $\partial\omega * \{v\} \cup \omega \subseteq K$.

Observe that $\partial\omega * \{v\} \cup \omega = \partial\tau$ where $\tau = \omega * \{v\}$. Thus $\partial\tau \subseteq K$. As K is flag and $|\omega * \{v\}| > 2$, this implies that $\tau = \omega * \{v\} \in K$. Hence, $\omega \in \text{link}_K v$, which contradicts the supposition. \square

LEMMA 2.2. *Let K be a flag complex on the set $[m]$ and let v be a vertex of K . Then $K \setminus \{v\}$, $\text{star}_K(v)$ and $\text{link}_K(v)$ are all flag complexes.*

Proof. Since $K \setminus \{v\}$ is a full subcomplex of K , any missing face of $K \setminus \{v\}$ is also a missing face of K . So as K is flag, any missing face has at most two elements, implying that any missing face of $K \setminus \{v\}$ also has at most two elements. Thus $K \setminus \{v\}$ is flag.

Let $\omega \in \text{MF}(\text{star}_K(v))$ and $|\omega| \geq 2$. We claim that $\omega \in \text{MF}(K)$ as well. As $\partial\omega \subseteq \text{star}_K(v)$, we also have $\partial\omega \subseteq K$, so if the claim does not hold then it must be the case that $\omega \in K$. Then $v \notin \omega$, as otherwise $\omega \in \text{star}_K(v)$. For $\tau = \omega * v$ we have $\partial\tau = \partial\omega * v \cup \omega \in K$. As K is flag and $|\omega * \{v\}| > 2$, we obtain $\tau = \omega * \{v\} \in K$. This implies that $\omega \in \text{star}_K(v)$, which contradicts the supposition. Hence, $\omega \in \text{MF}(K)$ and so $|\omega| = 2$ since K is flag. Thus $\text{star}_K(v)$ is flag.

Let $\omega \in \text{MF}(\text{link}_K(v))$ and $|\omega| \geq 2$. By Lemma 2.1, $\omega \in \text{MF}(K \setminus \{v\})$ as well. It has already been established that $K \setminus \{v\}$ is flag, so we have $|\omega| = 2$. Thus $\text{link}_K(v)$ is also flag. \square

Given a subset $\omega \subseteq [m]$, the full subcomplex of K on ω is

$$K_\omega = \{\sigma \in K \mid \sigma \subseteq \omega\}.$$

Note that $K \setminus \{v\} = K_{[m] \setminus \{v\}}$. A key property that will be important subsequently is the following.


LEMMA 2.3. *Let K be a flag complex on the set $[m]$ and let v be a vertex of K . Then $\text{link}_K(v)$ is a full subcomplex of $K \setminus \{v\}$.*

Proof. Let ω be the vertex set of $\text{link}_K(v)$. Suppose that $\text{link}_K(v)$ is not a full subcomplex of $K \setminus \{v\}$. Then there is a face $\sigma \in K \setminus \{v\}$ such that $\sigma \subseteq \omega$ and $\sigma \notin \text{link}_K(v)$. By selecting a proper face of σ if necessary, we may assume that σ is a missing face of $\text{link}_K(v)$ with $|\sigma| \geq 2$. But then as K is flag, Lemma 2.1 implies that σ is also a missing face of $K \setminus \{v\}$. In particular, $\sigma \notin K \setminus \{v\}$, which contradicts the supposition. \square

3. Homotopy theoretic preparation

3.1 The Cube Lemma

Assume that all spaces are pointed and have nondegenerate basepoints, implying that the inclusion of the basepoint is a cofibration. This holds, for example, for pointed CW-complexes, and hence for polyhedral products. One part of Mather’s Cube Lemma [Mat76] states that if there is a diagram of spaces and maps



where the bottom face is a homotopy pushout and the four sides are obtained by pulling back with $H \rightarrow D$, then the top face is also a homotopy pushout. In what follows this will be used to identify the homotopy type of the pushout H in a certain context. However, we need this identification to have a naturality property, which is not immediate from the statement of the Cube Lemma. To obtain this, we prove a special case of the Cube Lemma from first principles.

In what follows, we work with strictly commutative pushouts and pullbacks rather than homotopy commutative ones. For a space Y let 1_Y be the identity map on Y . Suppose that there

is a strictly commutative diagram

$$\begin{array}{ccc}
 B \times A & \xrightarrow{1_B \times i} & B \times X \\
 \downarrow j \times 1_A & & \downarrow \\
 C \times A & \xrightarrow{\quad} & D \\
 & \searrow 1_C \times i & \swarrow f \\
 & & C \times X
 \end{array}
 \quad (2)$$

where the square is a pushout, and the maps i, j and f are pointed inclusions of subspaces. We will turn the maps $f, 1_C \times i, j \times 1_X$ and $j \times i$ from the four corners of the pushout to $C \times X$ into fibrations, up to homotopy, and examine their fibres.

There is a standard way of turning a pointed, continuous map $g : Y \rightarrow Z$ between locally compact, Hausdorff spaces into a fibration, up to homotopy. Let I be the unit interval and let $\text{Map}(I, Z)$ be the space of continuous (not necessarily pointed) maps from I to Z . Let $d : \text{Map}(I, Z) \rightarrow Z \times Z$ be defined by evaluating a map $\omega : I \rightarrow Z$ at the two endpoints, explicitly, $d(\omega) = (\omega(0), \omega(1))$. Define the space \tilde{P}_g by the pullback

$$\begin{array}{ccc}
 \tilde{P}_g & \longrightarrow & \text{Map}(I, Z) \\
 \downarrow & & \downarrow \text{ev}_0 \\
 Y & \xrightarrow{g} & Z
 \end{array}$$

where $\text{ev}_0(\omega) = \omega(0)$. As a set,

$$\tilde{P}_g = \{(y, \omega) \in Y \times \text{Map}(I, Z) \mid \omega(0) = g(y)\}. \quad (3)$$

Then, as in [Sel97, p. 59] for example, there is an inclusion $Y \rightarrow \tilde{P}_g$ which is a homotopy equivalence and the composite

$$q : \tilde{P}_g \rightarrow \text{Map}(I, Z) \xrightarrow{\text{ev}_1} Z$$

is a fibration, where $\text{ev}_1(\omega) = \omega(1)$. Moreover, if 1 is the basepoint of I and PZ is the path space of Z (with paths at time 1 ending at the basepoint of Z), then the fibre of q is homeomorphic to the mapping path space of g ,

$$P_g = \{(y, \omega) \in Y \times PZ \mid \omega(0) = g(y)\}, \quad (4)$$

which is obtained by the pullback

$$\begin{array}{ccc}
 P_g & \longrightarrow & PZ \\
 \downarrow & & \downarrow \text{ev}_0 \\
 Y & \xrightarrow{g} & Z
 \end{array}$$

Consider how these constructions behave with respect to pointed subspace inclusions. Let $S \xrightarrow{s} Y$ be the inclusion of a pointed subspace. If Q is the pullback of $S \xrightarrow{s} Y$ and $\tilde{P}_g \rightarrow Y$, then the pullback defining \tilde{P}_g implies that Q is also the pullback of $g \circ s$ and ev_0 . But this pullback is the definition of $\tilde{P}_{g \circ s}$, so $Q = \tilde{P}_{g \circ s}$. Similarly for P_g , giving pullbacks

$$\begin{array}{ccc}
 \tilde{P}_{g \circ s} & \longrightarrow & \tilde{P}_g \\
 \downarrow & & \downarrow \\
 S & \xrightarrow{s} & Y
 \end{array}
 \quad
 \begin{array}{ccc}
 P_{g \circ s} & \longrightarrow & P_g \\
 \downarrow & & \downarrow \\
 S & \xrightarrow{s} & Y
 \end{array}$$

Since P_g and $P_{g \circ s}$ are the respective fibres of \tilde{P}_g and $\tilde{P}_{g \circ s}$ over Z , we obtain a pullback

$$\begin{array}{ccc} P_{g \circ s} & \longrightarrow & P_g \\ \downarrow & & \downarrow \\ \tilde{P}_{g \circ s} & \longrightarrow & \tilde{P}_g \end{array} \tag{5}$$

Next, suppose that Y is the union of pointed, closed subspaces S and T . Let $s : S \rightarrow Y$ and $t : T \rightarrow Y$ be the pointed subspace inclusions, and let u and v be the pointed subspace inclusions $u : S \cap T \rightarrow S$ and $v : S \cap T \rightarrow T$. Since S and T are closed subspaces of Y , the pushout of u and v is Y . (More generally this is true if $(Y; S, T)$ is an excisive triad, but we do not need this level of generality: in our case each of S, T and Y will be certain polyhedral products.)

LEMMA 3.1. *Suppose that $Y \xrightarrow{g} Z$ is a pointed subspace inclusion and that $Y = S \cup T$ where S and T are closed, pointed subspaces of Y . Then there are pushouts*

$$\begin{array}{ccc} \tilde{P}_{g \circ s \circ u} & \longrightarrow & \tilde{P}_{g \circ t} \\ \downarrow & & \downarrow \\ \tilde{P}_{g \circ s} & \longrightarrow & \tilde{P}_g \end{array} \qquad \begin{array}{ccc} P_{g \circ s \circ u} & \longrightarrow & P_{g \circ t} \\ \downarrow & & \downarrow \\ P_{g \circ s} & \longrightarrow & P_g \end{array}$$

Proof. By its definition, \tilde{P}_g is the space of paths on Z that begin in $\text{Im}(g)$ and end in Z . As g is a subspace inclusion, we may regard \tilde{P}_g as the space of paths on Z that begin in Y and end in Z . As $Y = S \cup T$, any such path begins either in S or in T , that is, the path is either in $\tilde{P}_{g \circ s}$ or $\tilde{P}_{g \circ t}$. Moreover, the intersection $\tilde{P}_{g \circ s} \cap \tilde{P}_{g \circ t}$ is all paths on Z that begin in $S \cap T$ and end in Z , that is, the paths in $\tilde{P}_{g \circ s \circ u} = \tilde{P}_{g \circ t \circ v}$. Thus $\tilde{P}_g = \tilde{P}_{g \circ s} \cup \tilde{P}_{g \circ t}$ and $\tilde{P}_{g \circ s \circ u} = \tilde{P}_{g \circ s} \cap \tilde{P}_{g \circ t}$. Further, since S and T are closed subspaces of Y , we have $\tilde{P}_{g \circ s}$ and $\tilde{P}_{g \circ t}$ closed subspaces of \tilde{P}_g . Therefore there is a pushout

$$\begin{array}{ccc} \tilde{P}_{g \circ s \circ u} & \longrightarrow & \tilde{P}_{g \circ t} \\ \downarrow & & \downarrow \\ \tilde{P}_{g \circ s} & \longrightarrow & \tilde{P}_g \end{array} \tag{6}$$

The same argument shows that P_g is the pushout of $P_{g \circ s}$ and $P_{g \circ t}$ over $P_{g \circ s \circ u} = P_{g \circ t \circ v}$. □

Now apply this construction to the maps $f, 1_C \times i, j \times 1_X$ and $j \times i$ from the four corners of the pushout in (2) to $C \times X$.

LEMMA 3.2. *There is a commutative cube*

$$\begin{array}{ccccc} P_{j \times i} & \longrightarrow & P_{j \times 1_X} & & \\ \downarrow & \searrow & \downarrow & \searrow & \\ & P_{1_C \times i} & \longrightarrow & P_f & \\ \downarrow & \downarrow & \downarrow & \downarrow & \\ \tilde{P}_{j \times i} & \longrightarrow & \tilde{P}_{j \times 1_X} & & \\ \downarrow & \searrow & \downarrow & \searrow & \\ & \tilde{P}_{1_C \times i} & \longrightarrow & \tilde{P}_f & \end{array}$$

where the top and bottom faces are pushouts and the four sides are pullbacks. Further, this cube is natural for maps of diagrams of the form (2).

Proof. Since $f, 1_C \times i, j \times 1_X$ and $j \times i$ are all subspace inclusions, the four sides of the cube are pullbacks by (5). Since D is a pushout, it is the union of $C \times A$ and $B \times X$ with intersection $B \times A$. The top and bottom faces of the cube are therefore pushouts by (6). The naturality statement holds since the constructions of \tilde{P}_g and P_g are natural. \square

The top face of the cube in Lemma 3.2 will be more precisely identified. This requires two lemmas.

LEMMA 3.3. A map $g \times h : Y \times M \rightarrow Z \times N$ has $P_{g \times h} = P_g \times P_h$. Further, this decomposition is natural for compositions $s \times t : Z \times N \rightarrow Z' \times N'$.

Proof. First observe that $P(Z \times N) = PZ \times PN$ since any pointed path $\omega : I \rightarrow Z \times N$ is equivalent to the product of the pointed paths $\omega_1 : I \rightarrow Z$ and $\omega_2 : I \rightarrow N$ given by projecting ω to Z and N respectively. Moreover, the evaluation map $P(Z \times N) \xrightarrow{ev_0} Z \times N$ becomes a product of evaluation maps $PZ \times PN \xrightarrow{ev_0 \times ev_0} Z \times N$. Thus the pullback $P_{g \times h}$ is identical to the pullback

$$\begin{array}{ccc} Q & \longrightarrow & PZ \times PN \\ \downarrow & & \downarrow ev_0 \times ev_0 \\ Y \times M & \xrightarrow{s \times t} & Z \times N \end{array}$$

where

$$\begin{aligned} Q &= \{((y, m), (\omega_1, \omega_2)) \in Y \times M \times PZ \times PN \mid s(y) = \omega_1(0), t(m) = \omega_2(0)\} \\ &= \{(y, \omega_1) \in Y \times PZ \mid s(y) = \omega_1(0)\} \times \{(m, \omega_2) \in M \times PN \mid t(m) = \omega_2(0)\} \\ &= P_s \times P_t. \end{aligned}$$

The identification of $P_{s \times t}$ as $P_s \times P_t$ only used the fact that $P(Z \times N) = PZ \times PN$. As the latter decomposition is natural, therefore so is the former. \square

LEMMA 3.4. There is a natural homeomorphism $P_{1_Y} \cong PY$.

Proof. Taking $g = 1_Y$ in (4) gives

$$P_{1_Y} = \{(y, \omega) \in Y \times PY \mid \omega(0) = y\}.$$

Define $\phi : PY \rightarrow P_{1_Y}$ by $\phi(\omega) = (\omega(0), \omega)$ and $\psi : P_{1_Y} \rightarrow PY$ by $\psi(y, \omega) = \omega$. Both ϕ and ψ are continuous, $\psi \circ \phi = \text{id}_{PY}$ and, because for any pair $(y, \omega) \in P_{1_Y}$ there is the condition $y = \omega(0)$, we also have $\phi \circ \psi = 1_{P_{1_Y}}$. Hence ψ is a homeomorphism. As both ϕ and ψ are natural, the homeomorphism is too. \square

Applying Lemmas 3.3 and 3.4 to the top face in Lemma 3.2, the space P_f is homeomorphic to the space Q_f defined by the pushout

$$\begin{array}{ccc} P_j \times P_i & \longrightarrow & P_j \times PX \\ \downarrow & & \downarrow \\ PC \times P_i & \longrightarrow & Q_f \end{array} \tag{7}$$

Moreover, the naturality statements in Lemmas 3.2–3.4 imply that (7) is natural for maps of diagrams of the form (2).

One further modification of (7) is needed. If Y is a pointed space, the reduced cone on Y is the space $CY = Y \wedge I$ (i.e., $CY = (Y \times I)/(Y \vee I)$). If Y and Z are pointed spaces with basepoints y_0 and z_0 respectively, then the reduced join is defined by $Y * Z = (Y \times I \times Z)/\sim$, where $(y, 0, z) = (y, 0, z')$, $(y, 1, z) \sim (y', 1, z)$ and $(y_0, t, z_0) = (y_0, 0, z_0)$ for all $y, y' \in Y$, $z, z' \in Z$ and $t \in I$. Observe that there is a pushout

$$\begin{array}{ccc} Y \times Z & \longrightarrow & Y \times CZ \\ \downarrow & & \downarrow \\ CY \times Z & \longrightarrow & Y * Z \end{array}$$

PROPOSITION 3.5. *Up to homotopy equivalences, the top face in Lemma 3.2 can be identified with the pushout*

$$\begin{array}{ccc} P_j \times P_i & \longrightarrow & P_j \times CP_i \\ \downarrow & & \downarrow \\ CP_j \times P_i & \longrightarrow & P_j * P_i \end{array}$$

*In particular, P_f is homotopy equivalent to $P_j * P_i$. Further, this homotopy equivalence may be chosen to be natural for maps of diagrams of the form (2).*

Proof. In general, suppose that Z is contractible. Then there is a pointed homotopy $Z \times I \rightarrow Z$ which at $t = 0$ is the identity map on Z and at $t = 1$ is the constant map to the basepoint. The homotopy sends $Z \vee I$ to the basepoint, and so factors through a map $CZ = Z \wedge I \rightarrow Z$. That is, the contracting homotopy for Z determines a specific map $CZ \rightarrow Z$. If the contracting homotopy is natural for maps $Z \rightarrow Z'$, then the map $CZ \rightarrow Z$ is also natural. In fact, it is a natural homotopy equivalence. Refining, if $g : Y \rightarrow Z$ is a pointed map with Z being contractible, then we obtain a composite $CY \xrightarrow{Cg} CZ \rightarrow Z$ with the same naturality properties.

In our case, consider (7). Since PC and PX are contractible, we obtain composites $P_j \rightarrow CP_j \rightarrow PC$ and $P_i \rightarrow CP_i \rightarrow PX$ in which the right-hand maps are homotopy equivalences. Thus the pushout Q_f in (7) is homotopy equivalent to the space $P_j * P_i$ obtained from the pushout

$$\begin{array}{ccc} P_j \times P_i & \longrightarrow & P_j \times CP_i \\ \downarrow & & \downarrow \\ CP_j \times P_i & \longrightarrow & P_j * P_i \end{array}$$

Since P_f is homeomorphic to Q_f , we obtain $P_f \simeq P_j * P_i$. Further, since the contracting homotopy for a path space PZ can be chosen to be natural for any map $Z \rightarrow Z'$, this homotopy equivalence for P_f is natural to the same extent as (7) is natural. That is, it is natural for maps of diagrams of the form (2). □

3.2 Two general results on fibrations

Now assume that all spaces have the homotopy type of pointed CW -complexes. If X is such a space, then by [Mil59, Corollary 3] so is ΩX . Also, any weak homotopy equivalence between two such spaces is a homotopy equivalence (see, for example, [Spa66, ch. 7, § 6, Corollary 24]).

LEMMA 3.6. Suppose that $\Omega B \xrightarrow{\partial} F \xrightarrow{f} E \xrightarrow{p} B$ is a homotopy fibration sequence and p has a left homotopy inverse. Then ∂ has a right homotopy inverse.

Proof. Let $s : B \rightarrow E$ be a map such that $s \circ p$ is homotopic to the identity map on E . Then $f \simeq s \circ p \circ f$, implying that f is null homotopic since $p \circ f$ is. If X is any pointed space, then the homotopy fibration $\Omega B \xrightarrow{\partial} F \xrightarrow{f} E$ induces an exact sequence of pointed sets $[X, \Omega B] \xrightarrow{\partial_*} [X, F] \xrightarrow{f_*} [X, B]$, where $[X, Y]$ is the set of pointed homotopy classes of maps from X to Y . Since f is null homotopic, $f_* = 0$, so ∂_* is onto. Taking $X = F$ implies that the (homotopy class of the) identity map on F lifts through ∂_* to a map $t : F \rightarrow \Omega B$. That is, $\partial \circ t$ is homotopic to the identity map on F . \square

In general, if $F \xrightarrow{f} E \xrightarrow{p} B$ is a homotopy fibration, where E is an H -space and p has a right homotopy inverse $s : B \rightarrow E$, then the composite

$$B \times F \xrightarrow{s \times f} E \times E \xrightarrow{\mu} E$$

is a weak homotopy equivalence, and hence a homotopy equivalence. We wish to give a slight variation on this in the case when $B = B_1 \times B_2$ and each factor has a right homotopy inverse. For $i = 1, 2$ let p_i be the composite $p_i : E \xrightarrow{p} B_1 \times B_2 \xrightarrow{\pi_i} B_i$ where π_i is the projection. As maps into a product are determined by their projection onto each factor, we have $p = (p_1, p_2)$.

LEMMA 3.7. Let $F \xrightarrow{f} E \xrightarrow{p} B_1 \times B_2$ be a homotopy fibration where p is an H -map. Suppose that for $i = 1, 2$ there are maps $s_i : B_i \rightarrow E$ such that $p_i \circ s_i$ is homotopic to the identity map on B_i , and $p_i \circ s_j$ is null homotopic for $i \neq j$. Then the composite

$$B_1 \times B_2 \times F \xrightarrow{s_1 \times s_2 \times f} E \times E \times E \xrightarrow{\mu \circ (\mu \times 1)} E$$

is a homotopy equivalence, where μ is the multiplication on E .

Proof. From the general result stated before the lemma, it suffices to show that $s_1 \times s_2$ is a right homotopy inverse for p . Consider the diagram

$$\begin{array}{ccccc} B_1 \times B_2 & \xrightarrow{s_1 \times s_2} & E \times E & \xrightarrow{\mu} & E \\ & \searrow^{i_1 \times i_2} & \downarrow p \times p & & \downarrow p \\ & & (B_1 \times B_2) \times (B_1 \times B_2) & \xrightarrow{\mu'} & B_1 \times B_2 \end{array}$$

where i_1 and i_2 are the inclusions into the first and second factors respectively and μ' is the multiplication on $B_1 \times B_2$. The left triangle homotopy commutes since $p_i \circ s_i$ is homotopic to the identity map on B_i and $p_i \circ s_j \simeq *$ if $i \neq j$. The right square homotopy commutes since p is an H -map. Observe that the lower direction around the diagram is homotopic to the identity map on $B_1 \times B_2$. Therefore the upper direction around the diagram implies that $\mu \circ (s_1 \times s_2)$ is a right homotopy inverse for p . \square

4. Polyhedral products and the proof of Theorem 1.1

Let K be a simplicial complex on the set $[m]$ and let v be a vertex of K . Following Félix and Tanré [FT09], define a new simplicial complex \overline{K} on $[m]$ by

$$\overline{K} = K \setminus \{v\} * \{v\}.$$

Observe that there is an inclusion of simplicial complexes $K \setminus \{v\} \longrightarrow \overline{K}$ given by including the join factor, so as $\text{star}_K(v) = \text{link}_K(v) * \{v\}$, there is a pushout map

$$K \longrightarrow \overline{K}.$$

Observe also that $K \setminus \{v\}$ is the full subcomplex of \overline{K} . That is, $\overline{K} \setminus \{v\} = K \setminus \{v\}$.

By [GT13], the pushout of simplicial complexes in (1) induces a pushout of polyhedral products

$$\begin{array}{ccc} (\underline{X}, \underline{A})^{\text{link}_K(v)} \times A_v & \xrightarrow{1 \times i_v} & (\underline{X}, \underline{A})^{\text{link}_K(v)} \times X_v \\ \downarrow j \times 1 & & \downarrow \\ (\underline{X}, \underline{A})^{K \setminus \{v\}} \times A_v & \longrightarrow & (\underline{X}, \underline{A})^K \end{array} \tag{8}$$

where i_v is the inclusion. (Here we regard $\text{link}_K(v)$ and $K \setminus \{v\}$ as simplicial complexes on the set $[m] \setminus \{v\}$.) To relate this to $(\underline{X}, \underline{A})^{\overline{K}}$, observe that the definition of the join of two simplicial complexes implies that if $K = K_1 * K_2$, then there is a homeomorphism

$$(\underline{X}, \underline{A})^K \cong (\underline{X}, \underline{A})^{K_1} \times (\underline{X}, \underline{A})^{K_2}.$$

In particular, as $\overline{K} = K \setminus \{v\} * \{v\}$ there is a homeomorphism

$$(\underline{X}, \underline{A})^{\overline{K}} \cong (\underline{X}, \underline{A})^{K \setminus \{v\}} \times X_v$$

and a strictly commutative diagram

$$\begin{array}{ccc} (\underline{X}, \underline{A})^{\text{link}_K(v)} \times A_v & \xrightarrow{1 \times i_v} & (\underline{X}, \underline{A})^{\text{link}_K(v)} \times X_v \\ \downarrow j \times 1 & & \downarrow \\ (\underline{X}, \underline{A})^{K \setminus \{v\}} \times A_v & \longrightarrow & (\underline{X}, \underline{A})^K \end{array} \begin{array}{l} \searrow j \times 1 \\ \searrow f \\ \searrow 1 \times i_v \end{array} \begin{array}{l} \\ \\ \end{array} \begin{array}{l} \\ \\ (\underline{X}, \underline{A})^{K \setminus \{v\}} \times X_v \end{array} \tag{9}$$

where f is the map induced by the simplicial map $K \longrightarrow \overline{K}$ and all maps are inclusions of subspaces.

Let B_v^K be the fibre P_j obtained by turning the map $(\underline{X}, \underline{A})^{\text{link}_K(v)} \xrightarrow{j} (\underline{X}, \underline{A})^{K \setminus \{v\}}$ into a fibration and let Y_v be the fibre P_i obtained by turning the inclusion $A_v \xrightarrow{i_v} X_v$ into a fibration.

LEMMA 4.1. *If F_v^K is the fibre P_f obtained by turning the map $(\underline{X}, \underline{A})^K \xrightarrow{f} (\underline{X}, \underline{A})^{K \setminus \{v\}} \times X_v$ into a fibration, then there is a homotopy equivalence*

$$F_v^K \simeq B_v^K * Y_v.$$

Further, this homotopy equivalence is natural for inclusions of simplicial complexes $K \longrightarrow K'$ on the set $[m]$.

Proof. Proposition 3.5 immediately implies the asserted homotopy equivalence for F_v^K and states that it is natural for maps of diagrams of the form (9). Now observe that any inclusion of simplicial complexes $K \rightarrow K'$ on the vertex set $[m]$ induces such a map of diagrams. \square

To take this further we need a general result about polyhedral products.

LEMMA 4.2. *Suppose that K_ω is a full subcomplex of a simplicial complex K . Then the map of polyhedral products $(\underline{X}, \underline{A})^{K_\omega} \rightarrow (\underline{X}, \underline{A})^K$ induced by the simplicial inclusion $K_\omega \rightarrow K$ has a left inverse, that is, there is a retraction $(\underline{X}, \underline{A})^K \rightarrow (\underline{X}, \underline{A})^{K_\omega}$. Further, the construction of the left inverse is natural for simplicial inclusions $K \rightarrow K'$.*

Proof. We have

$$(\underline{X}, \underline{A})^K = \bigcup_{\sigma \in K} \left(\prod_{i \in \sigma} X_i \times \prod_{i \in [m] \setminus \sigma} A_i \right), \quad (\underline{X}, \underline{A})^{K_\omega} = \bigcup_{\sigma \in K, \sigma \subseteq \omega} \left(\prod_{i \in \sigma} X_i \times \prod_{i \in \omega \setminus \sigma} A_i \right).$$

Since each A_i is a pointed space, there is a canonical inclusion $(\underline{X}, \underline{A})^{K_\omega} \rightarrow (\underline{X}, \underline{A})^K$. Furthermore, for each $\sigma \in K$ there is a projection

$$r_\sigma : \prod_{i \in \sigma} X_i \times \prod_{i \in [m] \setminus \sigma} A_i \rightarrow \prod_{i \in \sigma \cap \omega} X_i \times \prod_{i \in \omega \setminus \sigma} A_i.$$

Since K_ω is a full subcomplex, the image of r_σ belongs to $(\underline{X}, \underline{A})^{K_\omega}$. The projections r_σ patch together to give a retraction $r = \bigcup_{\sigma \in K} r_\sigma : (\underline{X}, \underline{A})^K \rightarrow (\underline{X}, \underline{A})^{K_\omega}$. The naturality assertion follows from the naturality of inclusions and projections. \square

PROPOSITION 4.3. *Let K be a simplicial complex on the index set $[m]$ and let v be a vertex of K . Then there is a homotopy equivalence*

$$\Omega(\underline{X}, \underline{A})^K \simeq \Omega X_v \times \Omega(\underline{X}, \underline{A})^{K \setminus \{v\}} \times \Omega(B_v^K * Y_v)$$

which is natural for inclusions of simplicial complexes $K \rightarrow K'$ on the set $[m]$.

Proof. Consider the homotopy fibration

$$F_v^K \rightarrow (\underline{X}, \underline{A})^K \xrightarrow{f} (\underline{X}, \underline{A})^{K \setminus \{v\}} \times X_v \tag{10}$$

from Lemma 4.1. Observe that $K \setminus \{v\}$ and $\{v\}$ are the full subcomplexes of K on the sets $[m] - \{v\}$ and $\{v\}$ respectively. So by Lemma 4.2, the maps $s_1 : (\underline{X}, \underline{A})^{K \setminus \{v\}} \rightarrow (\underline{X}, \underline{A})^K$ and $s_2 : X_v \rightarrow (\underline{X}, \underline{A})^K$ have left inverses $(\underline{X}, \underline{A})^K \xrightarrow{f_1} (\underline{X}, \underline{A})^{K \setminus \{v\}}$ and $(\underline{X}, \underline{A})^K \xrightarrow{f_2} X_v = (\underline{X}, \underline{A})^{\{v\}}$ respectively. Since the vertex sets for $K \setminus \{v\}$ and $\{v\}$ are disjoint, the left inverses have the property that $f_1 \circ s_2$ and $f_2 \circ s_1$ are trivial. Lemma 3.7 cannot be applied immediately since f is usually not an H -map, but after looping the homotopy fibration (10) it can be applied, and this gives the asserted homotopy equivalence.

The naturality property follows from the naturality properties of the simplicial map $K \rightarrow K \setminus \{v\} * \{v\}$, the polyhedral product and Lemma 4.2, together with the fact that $\Omega(\underline{X}, \underline{A})^K \rightarrow \Omega(\underline{X}, \underline{A})^{K'}$ is an H -map. \square

One more preliminary result is needed before the proof of Theorem 1.1. Let K be a simplicial complex on the vertex set $[m]$, let K^f be the flagification of K , and let L be the simplicial complex consisting of the vertices of K . Let M be either L or K . If v is a vertex of K , then the simplicial map $M \rightarrow K^f$ implies that there is commutative diagram of simplicial complexes

$$\begin{array}{ccc} \text{link}_M(v) & \longrightarrow & M \setminus \{v\} \\ \downarrow & & \downarrow \\ \text{link}_{K^f}(v) & \longrightarrow & K^f \setminus \{v\} \end{array}$$

Taking polyhedral products and then taking homotopy fibres gives a homotopy fibration diagram

$$\begin{array}{ccccccc} \Omega(\underline{X}, \underline{A})^{M \setminus \{v\}} & \longrightarrow & B_v^M & \longrightarrow & (\underline{X}, \underline{A})^{\text{link}_M(v)} & \longrightarrow & (\underline{X}, \underline{A})^{M \setminus \{v\}} \\ \downarrow & & \downarrow b_v & & \downarrow & & \downarrow \\ \Omega(\underline{X}, \underline{A})^{K^f \setminus \{v\}} & \longrightarrow & B_v^{K^f} & \longrightarrow & (\underline{X}, \underline{A})^{\text{link}_{K^f}(v)} & \longrightarrow & (\underline{X}, \underline{A})^{K^f \setminus \{v\}} \end{array} \tag{11}$$

for some induced map of fibres b_v .

LEMMA 4.4. *Let M be either L or K . Suppose that in (11) the map $\Omega(\underline{X}, \underline{A})^{M \setminus \{v\}} \rightarrow \Omega(\underline{X}, \underline{A})^{K^f \setminus \{v\}}$ has a right homotopy inverse. Then b_v has a right homotopy inverse $s_v : B_v^{K^f} \rightarrow B_v^M$. Moreover, s_v can be chosen so that it factors through the map $\Omega(\underline{X}, \underline{A})^{M \setminus \{v\}} \rightarrow B_v^M$.*

Proof. Consider the homotopy fibration along the bottom row of (11). Since K^f is flag, by Lemma 2.3, $\text{link}_{K^f}(v)$ is a full subcomplex of $K^f \setminus \{v\}$. Thus $(\underline{X}, \underline{A})^{\text{link}_{K^f}(v)}$ is a retract of $(\underline{X}, \underline{A})^{K^f \setminus \{v\}}$. Therefore, by Lemma 3.6, the map $\Omega(\underline{X}, \underline{A})^{K^f \setminus \{v\}} \rightarrow B_v^{K^f}$ has a right homotopy inverse $t : B_v^{K^f} \rightarrow \Omega(\underline{X}, \underline{A})^{K^f \setminus \{v\}}$. By hypothesis, the map $\Omega(\underline{X}, \underline{A})^{M \setminus \{v\}} \rightarrow \Omega(\underline{X}, \underline{A})^{K^f \setminus \{v\}}$ has a right homotopy inverse $s : \Omega(\underline{X}, \underline{A})^{K^f \setminus \{v\}} \rightarrow \Omega(\underline{X}, \underline{A})^{M \setminus \{v\}}$. Thus there is a homotopy commutative diagram

$$\begin{array}{ccccccc} B_v^{K^f} & \xrightarrow{t} & \Omega(\underline{X}, \underline{A})^{K^f \setminus \{v\}} & \xrightarrow{s} & \Omega(\underline{X}, \underline{A})^{M \setminus \{v\}} & \longrightarrow & B_v^M \\ & & \searrow & & \downarrow & & \downarrow b_v \\ & & & & \Omega(\underline{X}, \underline{A})^{K^f \setminus \{v\}} & \longrightarrow & B_v^{K^f} \end{array}$$

As the lower direction around the diagram is homotopic to the identity map on $B_v^{K^f}$, the upper direction around the diagram implies that b_v has a right homotopy inverse. \square

Proof of Theorem 1.1. Let K be a simplicial complex on the vertex set $[m]$, let K^f be its flagification, and let L be m disjoint points. Then there is a sequence of inclusions of simplicial complexes $L \rightarrow K \rightarrow K^f$. Taking polyhedral products with respect to $(\underline{X}, \underline{A})$ gives a sequence of maps $h : (\underline{X}, \underline{A})^L \xrightarrow{g} (\underline{X}, \underline{A})^K \xrightarrow{f} (\underline{X}, \underline{A})^{K^f}$. We will show that Ωh has a right homotopy inverse, implying that the map $\Omega f : \Omega(\underline{X}, \underline{A})^K \rightarrow \Omega(\underline{X}, \underline{A})^{K^f}$ also has a right homotopy inverse. This would prove both parts of the statement of the theorem.

The proof is by induction on the number of vertices. If $m = 1$, then L, K and K^f all equal the single vertex $\{1\}$, implying that h is the identity map, and so Ωh has a right homotopy inverse.

Assume that the statement of the theorem holds for all simplicial complexes with strictly less than m vertices. The decomposition and naturality statements in Proposition 4.3 imply that there is a homotopy commutative diagram of homotopy equivalences

$$\begin{CD}
 (\Omega(\underline{X}, \underline{A})^{L \setminus \{v\}} \times \Omega X_v) \times \Omega(B_v^L * Y_v) @>\simeq>> \Omega(\underline{X}, \underline{A})^L \\
 @VV(\Omega a \times 1) \times \Omega(b_v * 1)V @VVV \\
 (\Omega(\underline{X}, \underline{A})^{K^f \setminus \{v\}} \times \Omega X_v) \times \Omega(B_v^{K^f} * Y_v) @>\simeq>> \Omega(\underline{X}, \underline{A})^{K^f}
 \end{CD} \tag{12}$$

Observe that $K^f \setminus \{v\}$ has $m - 1$ vertices and $L \setminus \{v\} \rightarrow K^f \setminus \{v\}$ is the inclusion of these vertices. Since L and K^f are flag complexes, by Lemma 2.2 so are $L \setminus \{v\}$ and $K^f \setminus \{v\}$. Therefore, by inductive hypothesis, the map Ωa has a right homotopy inverse $s : \Omega(\underline{X}, \underline{A})^{K^f \setminus \{v\}} \rightarrow \Omega(\underline{X}, \underline{A})^{L \setminus \{v\}}$. As L and K^f are flag complexes and Ωa has a right homotopy inverse, by Lemma 4.4 the map b_v also has a right homotopy inverse $t : B_v^{K^f} \rightarrow B_v^L$. Therefore $t' = \Omega(t * 1)$ is a right homotopy inverse for $\Omega(b_v * 1)$. Putting s and t' together we obtain a map

$$\Omega(\underline{X}, \underline{A})^{K^f \setminus \{v\}} \times \Omega X_v \times \Omega(B_v^{K^f} * Y_v) \xrightarrow{s \times 1 \times t'} \Omega(\underline{X}, \underline{A})^{L \setminus \{v\}} \times \Omega X_v \times \Omega(B_v^L * Y_v)$$

which is a right homotopy inverse of $(\Omega a \times 1) \times \Omega(b_v * 1)$. The homotopy equivalences in (12) therefore imply that the map $h : \Omega(\underline{X}, \underline{A})^L \rightarrow \Omega(\underline{X}, \underline{A})^{K^f}$ has a right homotopy inverse. This completes the induction. \square

5. Refinements

This section gives two refinements describing the homotopy type of the space B_v^K under certain conditions. First consider the homotopy fibration diagram (11) in the case when $M = K$. Define the space D_v^K and the map d_v by the homotopy fibration

$$D_v^K \xrightarrow{d_v} B_v^K \xrightarrow{b_v} B_v^{K^f}. \tag{13}$$

LEMMA 5.1. *Given the hypotheses of Lemma 4.4, there is a homotopy equivalence*

$$B_v^K \simeq B_v^{K^f} \times D_v^K.$$

Proof. By Lemma 4.4, b_v has a right homotopy inverse $B_v^{K^f} \xrightarrow{s_v} B_v^K$. As B_v^K need not be an H -space this does not immediately imply that it is homotopy equivalent to $B_v^{K^f} \times D_v^K$. However, Lemma 4.4 also says that s_v can be chosen to factor through the homotopy fibration connecting map $\Omega(\underline{X}, \underline{A})^{K \setminus \{v\}} \rightarrow B_v^K$. That is, s_v can be chosen to be a composite $B_v^{K^f} \xrightarrow{s'_v} \Omega(\underline{X}, \underline{A})^{K \setminus \{v\}} \rightarrow B_v^K$ for some map s'_v . For any homotopy fibration sequence $\Omega B \xrightarrow{\delta} F \rightarrow E \rightarrow B$ the connecting map δ satisfies a homotopy action $\theta : \Omega B \times F \rightarrow F$ which restricts to the identity map on F and δ on ΩB . In our case, we obtain a composite

$$\psi : B_v^{K^f} \times D_v^K \xrightarrow{s'_v \times d_v} \Omega(\underline{X}, \underline{A})^{K \setminus \{v\}} \times B_v^K \xrightarrow{\theta} B_v^K.$$

Observe that the restriction of ψ to $B_v^{K^f}$ is s_v and the restriction to D_v^K is d_v . Thus ψ is a trivialization of the homotopy fibration (13), implying that it is a homotopy equivalence. \square

Second, suppose that K is a flag complex. By Lemma 2.3, $\text{link}_K(v)$ is a full subcomplex of $K \setminus \{v\}$. So by Lemma 4.2, the inclusion $(\underline{X}, \underline{A})^{\text{link}_K(v)} \rightarrow (\underline{X}, \underline{A})^{K \setminus \{v\}}$ has a left inverse. Define C_v^K by the homotopy fibration

$$C_v^K \rightarrow (\underline{X}, \underline{A})^{K \setminus \{v\}} \rightarrow (\underline{X}, \underline{A})^{\text{link}_K(v)}. \tag{14}$$

From the retraction of $(\underline{X}, \underline{A})^{\text{link}_K(v)}$ off $(\underline{X}, \underline{A})^{K \setminus \{v\}}$ and the definitions of B_v^K and C_v^K we obtain a homotopy pullback diagram

$$\begin{array}{ccccc} B_v^K & \xlongequal{\quad} & B_v^K & & \\ \downarrow & & \downarrow & & \\ * & \longrightarrow & (\underline{X}, \underline{A})^{\text{link}_K(v)} & \xlongequal{\quad} & (\underline{X}, \underline{A})^{\text{link}_K(v)} \\ \downarrow & & \downarrow & & \parallel \\ C_v^K & \longrightarrow & (\underline{X}, \underline{A})^{K \setminus \{v\}} & \longrightarrow & (\underline{X}, \underline{A})^{\text{link}_K(v)} \end{array}$$

Thus $B_v^K \simeq \Omega C_v^K$.

LEMMA 5.2. *Let K be a flag complex on the vertex set $[m]$ and let v be a vertex of K . Then there are homotopy equivalences*

$$\begin{aligned} \Omega(\underline{X}, \underline{A})^K &\simeq \Omega X_v \times \Omega(\underline{X}, \underline{A})^{K \setminus \{v\}} \times \Omega(\Omega C_v^K * Y_v), \\ \Omega(\underline{X}, \underline{A})^{K \setminus \{v\}} &\simeq \Omega(\underline{X}, \underline{A})^{\text{link}_K(v)} \times \Omega C_v^K. \end{aligned}$$

Proof. The first homotopy equivalence follows immediately from Proposition 4.3, while the second is an immediate consequence of the homotopy fibration (14) and the retraction of $(\underline{X}, \underline{A})^{\text{link}_K(v)}$ off $(\underline{X}, \underline{A})^{K \setminus \{v\}}$. □

6. Co- H -space properties

In this section we consider polyhedral products of the form $(\underline{CY}, \underline{Y})^K$ and identify the class of flag complexes K for which $(\underline{CY}, \underline{Y})^K$ is a co- H -space. As a corollary, we obtain conditions that allow for a delooping of the statement of Theorem 1.1. This begins with an abstract lemma.

LEMMA 6.1. *Let A and B be pointed spaces with the homotopy types of CW-complexes. Suppose that there is a pointed map $f : A \rightarrow B$ and B is a co- H -space. If Ωf has a right homotopy inverse, then f has a right homotopy inverse.*

Proof. Since B is a co- H -space, by [Gan70] there is a map $s : B \rightarrow \Sigma\Omega B$ which is a right homotopy inverse to the canonical evaluation map $ev : \Sigma\Omega B \rightarrow B$. Let $t : \Omega B \rightarrow \Omega A$ be a right homotopy inverse of Ωf . Consider the diagram

$$\begin{array}{ccc} B \xrightarrow{s} \Sigma\Omega B & & \\ \downarrow \Sigma t & \searrow & \\ \Sigma\Omega A & \xrightarrow{\Sigma\Omega f} & \Sigma\Omega B \\ \downarrow ev & & \downarrow ev \\ A & \xrightarrow{f} & B \end{array}$$

The upper triangle homotopy commutes since t is a right homotopy inverse of Ωf . The lower square homotopy commutes by the naturality of the evaluation map. The upper direction around the diagram is homotopic to $ev \circ s$, which is the identity map on B . The lower direction around the diagram therefore implies that $ev \circ \Sigma t \circ s$ is a right homotopy inverse of f . \square

PROPOSITION 6.2. *Let K be a simplicial complex on the vertex set $[m]$, let K^f be the flagification of K , and let Y_1, \dots, Y_m be pointed CW-complexes. If $(\underline{CY}, \underline{Y})^{K^f}$ is homotopy equivalent to a co- H -space, then the map $f : (\underline{CY}, \underline{Y})^K \rightarrow (\underline{CY}, \underline{Y})^{K^f}$ induced by the simplicial inclusion $K \rightarrow K^f$ has a right homotopy inverse.*

Proof. Taking $(\underline{X}, \underline{A}) = (\underline{CY}, \underline{Y})$, by Theorem 1.1, $\Omega f : \Omega(\underline{CY}, \underline{Y})^K \rightarrow \Omega(\underline{CY}, \underline{Y})^{K^f}$ has a right homotopy inverse. Since $(\underline{CY}, \underline{Y})^{K^f}$ is a co- H -space, Lemma 6.1 implies that f has a right homotopy inverse. \square

Remark 6.3. Note that in Proposition 6.2 we do not need to assume that Y_1, \dots, Y_m are path-connected. Since we assume that every singleton of $[m]$ is a vertex (K is on the vertex set $[m]$), $(\underline{CY}, \underline{Y})^K$ is path-connected even if Y is not.

Next we obtain a characterization of those flag complexes K for which $(\underline{CY}, \underline{Y})^K$ is a co- H -space. In terms of notation, when all pairs in the sequence $\{(X_i, A_i)\}_{i=1}^m$ are the same, $(X_i, A_i) = (X, A)$, we use the notation $(X, A)^K$ for $(\underline{X}, \underline{A})^K$. Special cases are the Davis–Januszkiewicz space $DJ(K) = (\mathbb{C}P^\infty, *)^K$ and the moment-angle complex $\mathcal{Z}_K = (D^2, S^1)^K$.

A graph Γ is called *chordal* if each of its cycles with at least four vertices has a chord (an edge joining two vertices that are not adjacent in the cycle). Equivalently, a chordal graph is a graph with no induced cycles of length more than three. By the result of Fulkerson and Gross [FG65] a graph is chordal if and only if its vertices can be ordered in such a way that, for each vertex i , the lesser neighbours of i form a clique. Such an order of vertices is called a *perfect elimination ordering*.

By [GPTW16], $\mathcal{Z}_{K^f} = (D^2, S^1)^{K^f}$ is homotopy equivalent to a wedge of spheres if and only if the 1-skeleton of K^f is a chordal graph. In particular, if the 1-skeleton of K^f is a chordal graph, then \mathcal{Z}_{K^f} is a co- H -space. This result is readily extended to general polyhedral products of the form $(\underline{CY}, \underline{Y})^K$, where CY denotes the cone over Y . Let $X^{\vee k}$ be the k -fold wedge of X .

THEOREM 6.4. *Assume that K is a flag complex on the vertex set $[m]$ and $\tilde{H}^*(Y_i; \mathbb{Z}) \neq 0$ for $1 \leq i \leq m$. The following conditions are equivalent:*

- (a) *the 1-skeleton K^1 is a chordal graph;*
- (b) *$(\underline{CY}, \underline{Y})^K$ is a co- H -space.*

Furthermore, if K^1 is chordal, there is a homotopy equivalence

$$(\underline{CY}, \underline{Y})^K \simeq \bigvee_{k=2}^m \bigvee_{1 \leq i_1 < \dots < i_k \leq m} (\Sigma Y_{i_1} \wedge \dots \wedge Y_{i_k})^{\vee c(i_1, \dots, i_k)}, \tag{15}$$

where $c(i_1, \dots, i_k) = \text{rank } \tilde{H}^0(K_{\{i_1, \dots, i_k\}})$ is one less than the number of connected components of the full subcomplex $K_{\{i_1, \dots, i_k\}}$.

Proof. The argument is similar to [GPTW16, Theorem 4.6] or [PV16, Theorem 4.3], but this time we keep track of the wedge summands. Assume that K^1 is chordal. Choose a perfect elimination ordering of vertices, and for each vertex $i = 1, \dots, m$ denote by σ_i the face of K corresponding to the clique of K^1 consisting of i and its lesser neighbours. All maximal faces of K are among $\sigma_1, \dots, \sigma_m$, so we have $\bigcup_{i=1}^m \sigma_i = K$. Furthermore, for each $k = 1, \dots, m$ the perfect elimination ordering on K induces such an ordering on the full subcomplex $K_{\{1, \dots, k-1\}}$, so we have $\bigcup_{i=1}^{k-1} \sigma_i = K_{\{1, \dots, k-1\}}$. In particular, the simplicial complex $\bigcup_{i=1}^{k-1} \sigma_i$ is flag as a full subcomplex in a flag complex. The intersection $\sigma_k \cap \bigcup_{i=1}^{k-1} \sigma_i$ is a clique $\sigma_k \setminus \{k\}$, so it is a face of $\bigcup_{i=1}^{k-1} \sigma_i$. Therefore, K is obtained by iteratively attaching σ_k to $\bigcup_{i=1}^{k-1} \sigma_i$ along the common face $\sigma_k \setminus \{k\}$.

We use induction on m to prove the decomposition (15). When $m = 1$, both sides of (15) are trivial. Now assume that (15) holds for K with $< m$ vertices. The pushout square (1) for $v = \{m\}$ becomes

$$\begin{array}{ccc} \sigma_m \setminus \{m\} & \longrightarrow & \sigma_m \\ \downarrow & & \downarrow \\ K \setminus \{m\} & \longrightarrow & K \end{array}$$

According to our convention, $\sigma_m \setminus \{m\}$ and $K \setminus \{m\}$ are regarded as simplicial complexes on $[m] \setminus \{m\} = [m - 1]$, while σ_m is regarded as a complex on $[m]$. The corresponding pushout square (8) of the polyhedral products becomes

$$\begin{array}{ccc} (\underline{CY}, \underline{Y})^{\sigma_m \setminus \{m\}} \times Y_m & \longrightarrow & (\underline{CY}, \underline{Y})^{\sigma_m} \\ \downarrow j \times 1 & & \downarrow \\ (\underline{CY}, \underline{Y})^{K \setminus \{m\}} \times Y_m & \longrightarrow & (\underline{CY}, \underline{Y})^K \end{array} \tag{16}$$

As $\sigma_m \setminus \{m\}$ is a face of $K \setminus \{m\}$ and σ_m is a face of K , we have

$$(\underline{CY}, \underline{Y})^{\sigma_m \setminus \{m\}} = \prod_{i \in \sigma_m \setminus \{m\}} CY_i \times \prod_{i \notin \sigma_m} Y_i, \quad (\underline{CY}, \underline{Y})^{\sigma_m} = \prod_{i \in \sigma_m} CY_i \times \prod_{i \notin \sigma_m} Y_i.$$

Since each $\{i\}$ is a vertex of K , the inclusion $\prod_{i \in \omega} Y_i \rightarrow (\underline{CY}, \underline{Y})^K$ is null-homotopic for any subset $\omega \subseteq [m]$, and the same holds with K replaced by $K \setminus \{m\}$. Hence, the map $j \times 1$ in (16) decomposes into the composition $i_2 \circ \pi_2$ of the projection onto the second factor and the inclusion. It follows that the pushout square (16) decomposes as

$$\begin{array}{ccc} \prod_{i \notin \sigma_m} Y_i \times Y_m & \xrightarrow{\pi_1} & \prod_{i \notin \sigma_m} Y_i \\ \downarrow \pi_2 & & \downarrow \\ Y_m & \xrightarrow{\epsilon} & (\prod_{i \notin \sigma_m} Y_i) * Y_m \\ \downarrow i_2 & & \downarrow \\ (\underline{CY}, \underline{Y})^{K \setminus \{m\}} \times Y_m & \longrightarrow & (\underline{CY}, \underline{Y})^K \end{array}$$

where the map ϵ is null-homotopic. From the bottom pushout square we obtain

$$(\underline{CY}, \underline{Y})^K \simeq ((\underline{CY}, \underline{Y})^{K \setminus \{m\}} \times Y_m) \vee \left(\left(\prod_{i \notin \sigma_m} Y_i \right) * Y_m \right),$$

where $X \rtimes Y = X \times Y / (* \times Y)$ is the right half-smash product, which is homotopy equivalent to $X \vee (X \wedge Y)$ when X is a suspension. By the inductive hypothesis, $(\underline{CY}, \underline{Y})^{K \setminus \{m\}}$ is a suspension, so we can rewrite the identity above as

$$(\underline{CY}, \underline{Y})^K \simeq (\underline{CY}, \underline{Y})^{K \setminus \{m\}} \vee ((\underline{CY}, \underline{Y})^{K \setminus \{m\}} \wedge Y_m) \vee \left(\bigvee_{\substack{1 \leq i_1 < \dots < i_k \leq m-1 \\ \{i_j, m\} \notin K}} \Sigma Y_{i_1} \wedge \dots \wedge Y_{i_k} \wedge Y_m \right).$$

Now a simple counting argument together with the inductive hypothesis gives (15). This also proves the implication (a)⇒(b).

To prove the implication (b)⇒(a), assume that K^1 is not chordal. Choose an induced chordless cycle K_ω with $|\omega| \geq 4$ (i.e., a full subcomplex isomorphic to the boundary of an $|\omega|$ -gon). Then there is a nontrivial product in the cohomology ring $H^*((\underline{CY}, \underline{Y})^{K_\omega}; \mathbb{Z})$. (When $(\underline{CY}, \underline{Y}) = (D^1, S^0)$, the polyhedral product $(D^1, S^0)^{K_\omega}$ is an orientable surface of positive genus [BP02, Example 6.40]; the general case then follows from [BBCG12, Theorem 1.9].) By Lemma 4.2, the same nontrivial product appears in $H^*((\underline{CY}, \underline{Y})^K; \mathbb{Z})$. Thus, $(\underline{CY}, \underline{Y})^K$ is not a co- H -space. □

Remark 6.5. Theorem 6.4 implies that the wedge decomposition of $\Sigma(\underline{CY}, \underline{Y})^K$ of [BBCG10] desuspends when K is flag and K^1 is chordal; this also follows from the results of Iriye and Kishimoto [IK18, Theorem 1.2, Proposition 3.2]. Other classes of simplicial complexes K with this property are described in [IK18] and [GT16]. The novelty of Theorem 6.4 compared to [IK18] is the description of the wedge decomposition of $(\underline{CY}, \underline{Y})^K$ in terms of the degree-zero cohomology of full subcomplexes of K , which does not follow readily from desuspending the decomposition in [BBCG10].

When K is not flag, the implication (b)⇒(a) of Theorem 6.4 still holds, but (a)⇒(b) fails. Indeed one can take K to be the boundary of a *cyclic polytope* [BP15, Example 1.1.17] of dimension $n \geq 4$ with $m > n + 1$ vertices. Then K^1 is a complete graph on m vertices, so it is chordal. On the other hand, $\mathcal{Z}_K = (D^2, S^1)^K$ is an $(m + n)$ -manifold with nontrivial cohomology product, so it cannot be a co- H -space.

Finally, we give conditions that allow for a delooping of the maps in Theorem 1.1.

COROLLARY 6.6. *Let K be a simplicial complex on the vertex set $[m]$ whose 1-skeleton is a chordal graph. If K^f is the flagification of K , then the map $f : (\underline{CY}, \underline{Y})^K \rightarrow (\underline{CY}, \underline{Y})^{K^f}$ has a right homotopy inverse.*

Proof. As K and K^f have the same 1-skeleton, Theorem 6.4 implies that $(\underline{CY}, \underline{Y})^{K^f}$ is a co- H -space (and even a suspension). The result follows from Proposition 6.2. □

COROLLARY 6.7. *Let K be a flag simplicial complex on the vertex set $[m]$, and let L be the simplicial complex given by m disjoint points. The map $h : (\underline{CY}, \underline{Y})^L \rightarrow (\underline{CY}, \underline{Y})^K$ has a right homotopy inverse if and only if the 1-skeleton of K is a chordal graph.*

Proof. Assume that K^1 is a chordal graph. As K is flag, Theorem 1.1 implies that Ωh has a right homotopy inverse, and Theorem 6.4 implies that $(\underline{CY}, \underline{Y})^K$ is a co- H -space. Then h has a right homotopy inverse by Lemma 6.1.

Now assume that h has a right homotopy inverse. Then $(\underline{CY}, \underline{Y})^K$ is a co- H -space, being a retract of the co- H -space $(\underline{CY}, \underline{Y})^L$. Theorem 6.4 implies that K^1 is a chordal graph. □

Remark 6.8. Given $(\underline{CY}, \underline{Y})^L \xrightarrow{g} (\underline{CY}, \underline{Y})^K \xrightarrow{f} (\underline{CY}, \underline{Y})^{K^f}$, Theorem 1.1 states that each of the two maps Ωf and $\Omega h = \Omega f \circ \Omega g$ has a right homotopy inverse. Corollary 6.6 gives a sufficient condition for a delooping of the first map, and Corollary 6.7 gives a necessary and sufficient condition for a delooping of the second map. In both cases the condition is that K^1 is a chordal graph. However, this condition is obviously not necessary for a delooping of Ωf . Indeed, f has a right inverse for any flag K , not only for those with chordal K^1 , because in this case $K^f = K$ and f is the identity map.

7. Whitehead products

In this section we describe two ways of relating the results of Theorems 1.1 and 6.4 to the classical iterated Whitehead products. First, we consider polyhedral products of the form $(\underline{X}, \ast)^K$ with flag K whose 1-skeleton is a chordal graph, and obtain a generalization (Proposition 7.1) of Porter’s description of the homotopy fiber of the inclusion of an m -fold wedge into a product in terms of Whitehead brackets. Second, we consider the loop space $\Omega(\underline{S}, \ast)^K$ on a polyhedral product of spheres for an arbitrary flag complex K , and obtain a generalization (Proposition 7.2) of the Hilton–Milnor theorem.

First, specialize to the case when each pair (X_i, A_i) is of the form (X_i, \ast) and write (\underline{X}, \ast) for $(\underline{X}, \underline{A})$. By [GT07], for example, there is a homotopy fibration

$$(\underline{C}\Omega\underline{X}, \underline{\Omega}X)^K \xrightarrow{\gamma_K} (\underline{X}, \ast)^K \longrightarrow \prod_{i=1}^m X_i$$

for any simplicial complex K . This is natural for simplicial inclusions, so if K is a flag complex on the vertex set $[m]$ and $L \rightarrow K$ is the inclusion of the vertex set, then there is a homotopy fibration diagram

$$\begin{CD} (\underline{C}\Omega\underline{X}, \underline{\Omega}X)^L @>\gamma_L>> (\underline{X}, \ast)^L @>>> \prod_{i=1}^m X_i \\ @V h' VV @VV h V @| \\ (\underline{C}\Omega\underline{X}, \underline{\Omega}X)^K @>\gamma_K>> (\underline{X}, \ast)^K @>>> \prod_{i=1}^m X_i \end{CD} \tag{17}$$

where both h and h' are induced maps of polyhedral products. By Theorem 1.1, $\Omega h'$ has a right homotopy inverse. Further, if K^1 is a chordal graph then Proposition 6.2 and Theorem 6.4 imply that h' has a right homotopy inverse.

Observe that as L is m disjoint points we have $(\underline{X}, \ast)^L = X_1 \vee \dots \vee X_m$, implying that $(\underline{C}\Omega\underline{X}, \underline{\Omega}X)^L$ is the homotopy fibre of the inclusion of the wedge into the product. Porter [Por66] identified the homotopy type of this fibre, from which we obtain a homotopy equivalence

$$(\underline{C}\Omega\underline{X}, \underline{\Omega}X)^L \simeq \bigvee_{k=2}^m \bigvee_{1 \leq i_1 < \dots < i_k \leq m} (\Sigma\Omega X_{i_1} \wedge \dots \wedge \Omega X_{i_k})^{\vee(k-1)}. \tag{18}$$

Notice that L^1 is a chordal graph and the decomposition in (18) exactly matches that of $(\underline{C}\Omega\underline{X}, \underline{\Omega}X)^L$ in (15). Moreover, by [The18, Theorem 6.2], Porter’s homotopy type identification can be chosen so that the composite

$$\varphi_L : \bigvee_{k=2}^m \bigvee_{1 \leq i_1 < \dots < i_k \leq m} (\Sigma\Omega X_{i_1} \wedge \dots \wedge \Omega X_{i_k})^{\vee(k-1)} \xrightarrow{\simeq} (\underline{C}\Omega\underline{X}, \underline{\Omega}X)^L \xrightarrow{\gamma_L} (\underline{X}, \ast)^L$$

is a wedge sum of iterated Whitehead products of the maps

$$ev_i : \Sigma\Omega X_i \xrightarrow{ev} X_i \hookrightarrow X_1 \vee \cdots \vee X_m = (\underline{X}, \ast)^L.$$

Returning to (17), the naturality of the Whitehead product implies that $h \circ \varphi_L$ is a wedge sum of Whitehead products mapping into $(\underline{X}, \ast)^K$. The right homotopy inverse for h' when K^1 is a chordal graph therefore implies the following.

PROPOSITION 7.1. *Let K be a flag complex such that K^1 is a chordal graph. Then the map $(\underline{C}\Omega X, \underline{\Omega}X)^K \xrightarrow{\gamma_K} (\underline{X}, \ast)^K$ factors through a wedge sum of Whitehead products.*

In the case when $(\underline{X}, \ast)^K = (\mathbb{C}P^\infty, \ast)^K = DJ(K)$ and $(\underline{C}\Omega X, \underline{\Omega}X)^K \simeq (D^2, S^1)^K = \mathcal{Z}_K$ the result above follows from [GPTW16, Theorem 4.3], where the Whitehead products were explicitly specified as iterated brackets of the canonical generators.

Theorem 1.1 also leads to a generalization of the Hilton–Milnor theorem. In this case we specialize to pairs $(\Sigma X_i, \ast)$, giving $(\underline{\Sigma}X, \ast)^L = \Sigma X_1 \vee \cdots \vee \Sigma X_m$. The Hilton–Milnor theorem states that there is a homotopy equivalence

$$\Omega(\Sigma X_1 \vee \cdots \vee \Sigma X_m) \simeq \prod_{\alpha \in L\langle V \rangle} \Omega(\Sigma X_1^{\wedge \alpha_1} \wedge \cdots \wedge X_m^{\wedge \alpha_m}), \tag{19}$$

where: V is a free \mathbb{Z} -module on m elements x_1, \dots, x_m ; $L\langle V \rangle$ is the free Lie algebra on V ; α runs over a \mathbb{Z} -module basis of $L\langle V \rangle$; and α_i is the number of occurrences of x_i in the bracket α . Here, if $\alpha_i = 0$, we interpret X_i as being omitted from the smash product rather than as being trivial. For example, $X_1^{\wedge 2} \wedge X_2^0 = X_1^{\wedge 2}$. The Hilton–Milnor theorem also describes the maps from the factors on the right side of (19) into $\Omega(\Sigma X_1 \vee \cdots \vee \Sigma X_m)$. If the length of α is 1, then the relevant factor is $\Omega \Sigma X_i$ for some i and the map $\Omega \Sigma X_i \rightarrow \Omega(\Sigma X_1 \vee \cdots \vee \Sigma X_m)$ is the loops on the inclusion into the wedge. If the length of α is larger than 1, then the map $\Omega(\Sigma X_1^{\wedge \alpha_1} \wedge \cdots \wedge X_m^{\wedge \alpha_m}) \rightarrow \Omega(\Sigma X_1 \vee \cdots \vee \Sigma X_m)$ is the loops on the Whitehead product corresponding to the bracket α .

By Theorem 1.1, if K is a flag complex on the vertex set $[m]$, then the map $\Omega(\underline{\Sigma}X, \ast)^L \xrightarrow{h} \Omega(\underline{\Sigma}X, \ast)^K$ has a right homotopy inverse. In particular, $\Omega(\underline{\Sigma}X, \ast)^K$ is a retract of the product on the right side of (19). It is probably the case that the retraction consists of selecting an appropriate subproduct, but this is not immediately clear. That is, simply knowing that Ωh has a right homotopy inverse leaves open the possibility that some of the factors $\Omega(\Sigma X_1^{\wedge \alpha_1} \wedge \cdots \wedge X_m^{\wedge \alpha_m})$ split as $A \times B$ where A retracts off $\Omega(\underline{\Sigma}X, \ast)^K$ while B does not. However, if we specialize a bit more, then this possibility is essentially eliminated.

Suppose that each X_i is a connected sphere S^{n_i-1} and write (\underline{S}, \ast) for $(\underline{\Sigma}X, \ast)$. Since each X_i is a sphere, the space $\Sigma X_1^{\wedge \alpha_1} \wedge \cdots \wedge X_m^{\wedge \alpha_m}$ is homotopy equivalent to a sphere, so the right side of (19) becomes a product of looped spheres. The space ΩS^n is indecomposable unless $n \in \{2, 4, 8\}$. In the latter case, we have a homotopy equivalence $\Omega H \times E : \Omega S^{2n-1} \times S^{n-1} \xrightarrow{\simeq} \Omega S^n$, which is a product of the looped Hopf map H and the suspension map E . The retraction of $\Omega(\underline{S}, \ast)^K$ off $\Omega(\underline{S}, \ast)^L$ implies the following.

PROPOSITION 7.2. *Let K be a flag complex. Then*

$$\Omega(\underline{S}, \ast)^K \simeq \left(\prod_{i=1}^m \Omega S^{n_i} \right) \times M,$$

where M is homotopy equivalent to a product of spheres and loops on spheres. Further,

- (a) a factor ΩS^n of M with $n \notin \{3, 7, 15\}$ maps to $\Omega(\underline{S}, *)^K$ by a looped Whitehead product $\Omega S^n \xrightarrow{\Omega w} \Omega(\underline{S}, *)^K$;
- (b) a factor ΩS^{2n-1} of M with $n \in \{2, 4, 8\}$ maps to $\Omega(\underline{S}, *)^K$ by a looped Whitehead product $\Omega S^{2n-1} \xrightarrow{\Omega w} \Omega(\underline{S}, *)^K$ or by a composite $\Omega S^{2n-1} \xrightarrow{\Omega H} \Omega S^n \xrightarrow{\Omega w} \Omega(\underline{S}, *)^K$, where H is the Hopf map;
- (c) a factor S^{n-1} of M has $n \in \{2, 4, 8\}$ and maps to $\Omega(\underline{S}, *)^K$ by a composite $S^{n-1} \xrightarrow{E} \Omega S^n \xrightarrow{\Omega w} \Omega(\underline{S}, *)^K$, where E is the suspension map and w is a Whitehead product.

Refining a bit, by [GT07] the homotopy fibration $(\underline{C}\Omega S, \underline{\Omega}S)^K \xrightarrow{\gamma_K} (\underline{S}, *)^K \rightarrow \prod_{i=1}^m S^{n_i}$ splits after looping to give a homotopy equivalence

$$\Omega(\underline{S}, *)^K \simeq \left(\prod_{i=1}^m \Omega S^{n_i} \right) \times \Omega(\underline{C}\Omega S, \underline{\Omega}S)^K.$$

Therefore, Proposition 7.2 implies that if K is a flag complex then $\Omega(\underline{C}\Omega S, \underline{\Omega}S)^K$ is homotopy equivalent to a product of spheres and loops on spheres, and under this homotopy equivalence $\Omega\gamma_K$ becomes a product of maps of the form $\Omega w, \Omega w \circ \Omega H$ or $\Omega w \circ E$.

This has implications for moment-angle complexes and Davis–Januszkiewicz spaces. Recall that $DJ(K) \simeq (\mathbb{C}P^\infty, *)^K$ and $\mathcal{Z}_K \simeq (D^2, S^1)^K$. There is a homotopy fibration

$$\mathcal{Z}_K \xrightarrow{\psi_K} DJ(K) \rightarrow \prod_{i=1}^m \mathbb{C}P^\infty$$

which splits after looping to give a homotopy equivalence

$$\Omega DJ(K) \simeq \left(\prod_{i=1}^m S^1 \right) \times \Omega \mathcal{Z}_K.$$

The inclusion $S^2 \rightarrow \mathbb{C}P^\infty$ induces maps of pairs $(S^2, *) \rightarrow (\mathbb{C}P^\infty, *)$ and $(C\Omega S^2, \Omega S^2) \rightarrow (C\Omega \mathbb{C}P^\infty, \Omega \mathbb{C}P^\infty) \xrightarrow{\simeq} (D^2, S^1)$. These then induce a commutative diagram of polyhedral products

$$\begin{CD} (C\Omega S^2, \Omega S^2)^K @>\gamma_K>> (S^2, *)^K \\ @V G VV @VV F V \\ \mathcal{Z}_K @>\psi_K>> DJ(K) \end{CD} \tag{20}$$

Observe that the suspension map $S^1 \xrightarrow{E} \Omega S^2$ induces a map of pairs $(CS^1, S^1) \rightarrow (C\Omega S^2, \Omega S^2)$ with the property that the composite $(CS^1, S^1) \rightarrow (C\Omega S^2, \Omega S^2) \rightarrow (D^2, S^1)$ is a homotopy equivalence. This implies that the map G in (20) has a right homotopy inverse. If K is a flag complex, then Proposition 7.2 says that $\Omega(C\Omega S^2, \Omega S^2)^K$ is homotopy equivalent to a product of spheres and loops on spheres, and the factors map to $\Omega(S^2, *)^K$ by maps of the form $\Omega w, \Omega w \circ \Omega H$ or $\Omega w \circ E$. Thus from the map G in (20) having a right homotopy inverse, and F being natural with respect to Whitehead products, we obtain the following.

COROLLARY 7.3. *Let K be a flag complex. Then $\Omega \mathcal{Z}_K$ is homotopy equivalent to a product of spheres and loops on spheres, and under this equivalence the map $\Omega \mathcal{Z}_K \xrightarrow{\Omega \psi_K} \Omega DJ(K)$ becomes a product of maps of the form $\Omega w, \Omega w \circ \Omega H$ or $\Omega w \circ E$ where w is a Whitehead product.*

Notice that \mathcal{Z}_K itself is often not a product or a wedge of spheres. For example, if K is the boundary of an n -gon for $n \geq 5$, then K is flag and \mathcal{Z}_K is diffeomorphic to a connected sum of products of two spheres. Nevertheless, $\Omega\mathcal{Z}_K$ is homotopy equivalent to a product of spheres and loops on spheres.

8. Homotopy theoretic consequences

We restrict attention to Davis–Januszkiewicz spaces $DJ(K) = (\mathbb{C}P^\infty, *)^K$ and moment-angle complexes $\mathcal{Z}_K = (D^2, S^1)^K$. Let $S^2 \rightarrow \mathbb{C}P^\infty$ be the inclusion of $S^2 \cong \mathbb{C}P^1$ into $\mathbb{C}P^\infty$. Then there is an induced map of polyhedral products

$$i_K : (S^2, *)^K \rightarrow (\mathbb{C}P^\infty, *)^K.$$

Building on the fact that the map G in (20) has a right homotopy inverse, in [GT16] the following was proved.

LEMMA 8.1. *The map Ωi_K has a right homotopy inverse.*

LEMMA 8.2. *Let K be a flag complex. Suppose that there is a map $h : (\mathbb{C}P^\infty, *)^K \rightarrow Y$ where Y is 2-connected. Then Ωh is null homotopic. Consequently, h induces the zero map on homotopy groups.*

Proof. Let L be the simplicial complex on m disjoint points. The simplicial map $L \rightarrow K$ induces a map of polyhedral products $g : (S^2, *)^L \rightarrow (S^2, *)^K$. Consider the composite

$$(S^2, *)^L \xrightarrow{g} (S^2, *)^K \xrightarrow{i_K} (\mathbb{C}P^\infty, *)^K \xrightarrow{h} Y.$$

Observe that by the definition of the polyhedral product, $(S^2, *)^L \simeq \bigvee_{i=1}^m S^2$. Since Y is 2-connected, the composite $h \circ i_K \circ g$ is therefore null homotopic. Since K is a flag complex, by Theorem 1.1, Ωg has a right homotopy inverse. Therefore $\Omega h \circ \Omega i_K$ is null homotopic. By Lemma 8.1, Ωi_K also has a right homotopy inverse. Therefore Ωh is null homotopic. \square

For example, let C be the homotopy cofibre of the composite

$$\psi : \bigvee_{i=1}^m S^2 \rightarrow \bigvee_{i=1}^m \mathbb{C}P^\infty \rightarrow DJ(K),$$

where the left map is the wedge of inclusions of the bottom cells and the right map is the map of polyhedral products induced by including the vertices into K . The description of $H^*(DJ(K); \mathbb{Z})$ (see, for example, [BP02]) implies that C is 3-connected. Therefore Lemma 8.2 implies that if K is a flag complex, then the quotient map

$$f : DJ(K) \rightarrow C = DJ(K) / \left(\bigvee_{i=1}^m S^2 \right)$$

induces the trivial map on homotopy groups.

Lemma 8.2 says that if K is a flag complex, then the bottom 2-spheres in $DJ(K)$ have a great impact on its homotopy theory. The next lemma says this much more dramatically in the case of \mathcal{Z}_K when K^1 is a chordal graph.

LEMMA 8.3. *Let K be a flag complex such that K^1 is a chordal graph. Then there is a homotopy commutative diagram*

$$\begin{array}{ccc} & \bigvee_{i=1}^m S^2 & \\ \lambda \nearrow & \downarrow \psi & \\ \mathcal{Z}_K & \longrightarrow & DJ(K) \end{array}$$

for some map λ .

Proof. As usual, let L be the vertex set of K . Consider the diagram

$$\begin{array}{ccc} (C\Omega S^2, \Omega S^2)^L & \xrightarrow{\gamma^L} & (S^2, *)^L \\ \downarrow h & & \downarrow \\ (C\Omega S^2, \Omega S^2)^K & \xrightarrow{\gamma^K} & (S^2, *)^K \\ \downarrow G & & \downarrow F \\ \mathcal{Z}_K & \xrightarrow{\psi_K} & DJ(K) \end{array}$$

The upper square is induced by the simplicial inclusion of L into K . The lower square homotopy commutes by (20). Notice that the right column is equal to ψ . As mentioned in the previous section, the map G has a right homotopy inverse. Since K is a flag complex and K^1 is a chordal graph, by Corollary 6.7, the map h has a right homotopy inverse. Thus $G \circ h$ has a right homotopy inverse and the lemma follows. \square

ACKNOWLEDGEMENTS

The research of the first author was carried out at the Institute for Information Transmission Problems of Russian Academy of Sciences and was supported by the Russian Science Foundation (grant no. 14-50-00150). The second author would like to thank the Royal Society for the award of an International Exchanges Grant which helped make this research possible. The authors would also like to thank the referee for making several helpful comments.

REFERENCES

BBCG10 A. Bahri, M. Bendersky, F. R. Cohen and S. Gitler, *The polyhedral product functor: a method of decomposition for moment-angle complexes, arrangements and related spaces*, Adv. Math. **225** (2010), 1634–1668.

BBCG12 A. Bahri, M. Bendersky, F. R. Cohen and S. Gitler, *Cup-products for the polyhedral product functor*, Math. Proc. Cambridge Philos. Soc. **153** (2012), 457–469.

BP02 V. M. Buchstaber and T. E. Panov, *Torus actions and their applications in topology and combinatorics*, University Lecture Series, vol. 24 (American Mathematical Society, Providence, RI, 2002).

BP15 V. M. Buchstaber and T. E. Panov, *Toric topology*, Mathematical Surveys and Monographs, vol. 204 (American Mathematical Society, Providence, RI, 2015).

FT09 Y. Félix and D. Tanré, *Rational homotopy of the polyhedral product functor*, Proc. Amer. Math. Soc. **137** (2009), 891–898.

FG65 D. R. Fulkerson and O. Gross, *Incidence matrices and interval graphs*, Pacific J. Math **15** (1965), 835–855.

- Gan70 T. Ganea, *Cogroups and suspensions*, *Invent. Math.* **9** (1970), 185–197.
- GPTW16 J. Grbić, T. Panov, S. Theriault and J. Wu, *The homotopy types of moment-angle complexes for flag complexes*, *Trans. Amer. Math. Soc.* **368** (2016), 6663–6682.
- GT07 J. Grbić and S. Theriault, *The homotopy type of the complement of a coordinate subspace arrangement*, *Topology* **46** (2007), 357–396.
- GT13 J. Grbić and S. Theriault, *The homotopy type of the polyhedral product for shifted complexes*, *Adv. Math.* **245** (2013), 690–715.
- GT16 J. Grbić and S. Theriault, *Homotopy theory in toric topology*, *Russian Math. Surveys* **71** (2016), 185–251.
- IK18 K. Iriye and D. Kishimoto, *Fat wedge filtration and decomposition of polyhedral products*, *Kyoto J. Math.*, doi:[10.1215/21562261-2017-0038](https://doi.org/10.1215/21562261-2017-0038).
- Mat76 M. Mather, *Pull-backs in homotopy theory*, *Canad. J. Math.* **28** (1976), 225–263.
- Mil59 J. Milnor, *On spaces having the homotopy type of a CW-complex*, *Trans. Amer. Math. Soc.* **90** (1959), 272–280.
- PV16 T. Panov and Y. Veryovkin, *Polyhedral products and commutator subgroups of right-angled Artin and Coxeter groups*, *Mat. Sb.* **207** (2016), 105–126 (in Russian); *Sb. Math.* **207** (2016), 1582–1600 (Engl. transl.).
- Por66 G. J. Porter, *The homotopy groups of wedges of suspensions*, *Amer. J. Math.* **88** (1966), 655–663.
- Sel97 P. Selick, *Introduction to homotopy theory*, *Fields Institute Monographs*, vol. 9 (American Mathematical Society, Providence, RI, 1997).
- Spa66 E. H. Spanier, *Algebraic topology* (McGraw-Hill, New York, Toronto, London, 1966).
- The18 S. Theriault, *The dual polyhedral product, cocategory and nilpotence*, *Adv. Math.* **340** (2018), 138–192.

Taras Panov tpanov@mech.math.msu.su

Department of Mathematics and Mechanics, Moscow State University, Leninskie Gory,
119991 Moscow, Russia

and

Institute for Theoretical and Experimental Physics, Moscow, Russia

and

Institute for Information Transmission Problems, Russian Academy of Sciences, Russia

Stephen Theriault S.D.Theriault@soton.ac.uk

Mathematical Sciences, University of Southampton, Southampton SO17 1BJ, UK