

in the synthesis of prostaglandins from arachidonic acid (AA), is associated with schizophrenia.

SCHIZOPHRENIC SYMPTOMS AND DIETARY INTAKE OF N-3 FATTY ACIDS

Jan E. Mellor, Jonathan D.E. Laugharne, Malcolm Peet. *University Department of Psychiatry, Northern General Hospital, Herries Road, Sheffield S5 7AU, UK*

There is evidence that certain n3 and n6 essential fatty acids (EFAs) are depleted in cell membranes from red blood cells (RBC) and brains of patients suffering from schizophrenia. If these findings are of primary significance then the possibility is raised of modifying schizophrenic symptomatology by dietary supplementation with fatty acids. We have carried out detailed analysis of dietary fatty acid intake of 20 schizophrenic patients. It was found that a greater intake of n3 fatty acids and particularly eicosapentenoic (EPA) in the normal diet, was associated with less severe schizophrenic symptoms and particularly less positive symptoms, as well as less tardive dyskinesia (TD). Furthermore, supplementation of the diet for 6 weeks with 10 g/day of concentrated fish oil (MaxEPA) resulted in significant amelioration of both schizophrenic symptoms and TD.

THE RELATIONSHIP BETWEEN PLASMA FATTY ACID CONCENTRATIONS AND MOVEMENT DISORDERS

A. Nilsson. *Department of Clinical Neurosciences, Sahlgrenska University Hospital, Sweden; Karsudden Hospital, Pl 4000, S-641 96 Katrineholm, Sweden*

It has been postulated that dyskinesia may be attributable to the schizophrenic disease process rather than to its treatment. This is in line with the hypothesis that a putative organic vulnerability may cause dyskinesia to emerge even without exposure to neuroleptics. In the present general population survey of dyskinetic phenomena, no 'a priori' assumption was made of their cause.

Spontaneous and tardive dyskinesias were studied in a random population sample of 446 men, aged 59 years. Dyskinesia (AIMS-score of ≥ 2 in any body part) was seen in 15.1% (n = 74). Dyskinetic men had higher cigarette consumption, more psychiatric morbidity and they were more often exposed to neuroleptics. *Dyskinesia was associated with several abnormalities in Essential Fatty Acid (EFA) concentrations in plasma, but the most consistent finding was low arachidonic acid (AA) levels in phospholipids, triglycerides and cholesterol esters. In a logistic regression model, cigarette consumption (p < 0.02), exposure to neuroleptics (p < 0.01), and low AA levels in the phospholipid fraction (p < 0.0001) were independently associated with dyskinesia. Further analyses of our data indicated that impaired conversion of linoleic acid (LA) to AA is a problem in dyskinesia.*

Conclusion: Dyskinesia is associated with EFA abnormalities. These abnormalities are present also in individuals who have not been exposed to neuroleptics and who have no psychiatric disorder. The results are compatible with the free radical hypothesis of dyskinesia, but they also indicate that impaired conversion of LA to AA contributes to the low EFA levels seen in dyskinetic men.

S59. Psychiatry and the media

Chairman: A Clare

'MINDERS' — A TV YEAR WITH A CMHT

Tom Burns. *St. George's Hospital Medical Lung, London*

Minders was a series of six programmes broadcast at prime time on BBC2 during May and June 1995 to an estimated audience of just under 2,000,000. Most of the programmes were for half an hour although two were extended. Most episodes focused on the experience of an individual patient centering on their contact with the mental health team. The programmes were ground-breaking in the way in which they obtained access to acute mental health crises, including police involvement in compulsory admissions and even the filming of an appeal tribunal against detention.

The reasons for agreeing to take part and the details of agreements between managers, staff and patients will be outlined. The mechanics and ethics of obtaining informed consent were complex. There were a whole series of complications which could not have been foreseen (e.g. dependency on the TV crew, family involvement and rights, differing staff responses and involvement of other patients).

Response to the series varied very much between different groups. It was also sharply divided between that to the first programme 'Whose Mind is it anyway? John's story' and the other five. 'Whose mind is it anyway?' charted the compulsory admission and care of a young black patient suffering from schizophrenia in such a way that most viewers were left confused about why he was admitted in the first place and outraged at the perceived effects of his treatment. As staff we felt betrayed and grossly misrepresented. The issues which led to this state of affairs has many lessons for those considering co-operating with the media.

Overall, however, we felt that the programmes gave a sympathetic and honest portrayal of the untidy reality of suffering from a mental illness. This may act as an antidote to the oversimplified sound-bite approach so common in this field. Hopefully it will help generate a more realistic debate about the problems inherent in modern mental health care. It has undoubtedly removed some of the barriers to further such programmes which hopefully will build upon its achievements and not repeat its failures.

THE PSYCHIATRIST AS MEDIA INTERVIEWER — FINDING THE LIMITS

Anthony Clare. *Clinical Professor of Psychiatry, Trinity College, Dublin*

The growing participation of professional psychiatrists within radio and television raises a number of important ethical questions concerning the limits of enquiry and the exploitation of subjects interviewed. Concerns regarding the abuse of psychiatric power and the devaluation of the image of psychiatry and psychiatrists are considered in this paper as are the similarities and differences between the roles and responsibilities of professional journalists and professional psychiatrists participating in the media. Questions considered include: Are psychiatrists justified in acting as interviewers in the public media? Is a degree of unacceptable manipulation involved in such interviews? How are appropriate limits set to both the content and the method of such interviews? In the light of this experience, the author, himself a psychiatrist who has participated within the British and Irish media for more than twenty years, puts forward certain guidelines which should govern the behaviour of psychiatrists