

- #2 (Green) Minimal;
 #3 (Yellow): Moderate;
 #4 (Red) Severe, but require no surgical management;
 #5 (Red. Op) Severe and require surgical management; or
 #6 (Black) Dead.

Finally, patients will be transferred to each medical field tent according to the triage categories.

III. In the third step, #1 and/or #2 patients will be transferred to their own houses or their relative's houses and #3, #4, and #5 will be transferred to the other general hospitals. The triage officer should be the most experienced medical officer who is in complete control of the reception and categorization, and will be appointed by the director of each department.

068. Avoiding Trocar-Related Injuries

Duncan Turner, MD, Mark W. Surrey, MD
 Center for Reproductive Medicine and Surgery, Beverly Hills, California USA

In the past, trocar insertions have been associated with a significant incidence of discus perforation as well as vascular injury. A new, radial-expanding, blunt trocar has been designed that holds promise in decreasing the incidence of these injuries. Postoperative complications most common to the defect caused by trocar insertions are incision hernia and small bowel obstruction. Because this device stretches the tissues of the anterior abdominal wall instead of cutting, the abdominal wall defects produced are significantly smaller, and, in theory, can be associated with a lower incidence of these postoperative complications.

In light of this, a study was performed at the University of California, San Francisco, Department of Surgery whereby 12 pigs were separated into two equal groups using the new expandable trocar in one-half and the conventional 10 mm, disposable trocar in the other half. Observations were made as to the shape of the muscular fascial defects.

The results indicate radial-expanding devices produced slit-like defects arranged in a gridiron fashion, whereas, the conventional trocars produced clover-leaf-shaped larger defects. There was visually less tissue ecchymoses associated with the radially expanding devices, as well as the smaller size.

122. An Interactive CD-ROM on Triage

Wim Van der Heyden, (MAA), Surgeon
 Surgeon Dijkzigt Hospital, Rotterdam, Traumatologic Department
 Scientific co-worker Vrije Universiteit Amsterdam Department of Disaster Medicine

An interactive, multimedia education and training CD-ROM on disaster medicine will be produced. Since many disasters are crossing national borders, generally accepted guidelines

for education will promote international collaboration and unity of doctrine to achieve uniformity standardization, and interoperability in medical plans, procedures, and protocols.

The interactive CD-ROM on disaster medicine can be used in all medicine or nursing schools and training centres for ambulance crews and other rescue workers. It is based on an educational curriculum.

A. Medical Care

1. Surgery and Traumatology
2. Anesthesiology and Resuscitation
3. Chemical Injuries
4. Nuclear Injuries
5. Infectious Diseases
6. Psychological Care

B. Public Health

1. Epidemiology
2. Sanitation Health and Environment
3. Nutrition and Alimentation

C. Disaster Management

1. Risk Management
2. Definition Classification and Evaluation of a Disaster
3. Medical Relief Organizations
4. Regulations
5. Planning
6. Coordination, Communication, Command
7. Logistics
8. Protection and Security
9. Triage
10. Prehospital Phase Management
11. Transportation
12. Hospital Phase Management
13. Information and News Media
14. Rehabilitation
15. Management of Displaced Persons and Refugees

Salient among the training methods is simulation. Computer simulation can be quite educational, and is relatively inexpensive to conduct. The triage program is completed and provides a lot of randomized casualties to the participating sorter.

081. Analysis of Fire Disasters in Taiwan

Lee-Min Wang, MD, Sheng-Chuan Hu, MD, Chen-Hsen Lee, MD
 Emergency Department, Veterans General Hospital-Taipei, and the Society of Emergency and Critical Care Medicine, Taiwan, Republic of China

Taiwan is located in a subtropical area, and has an area of 13,803 square miles. It has one of the world's densest population: the total population is about 21 million.

In recent years, because the environment has been changing from agriculture to industry, fire-disaster has resulted in major damage to peoples' lives and properties. From 1978 to 1993, the analysis of fire-disasters that occurred in Taiwan included:

- 1) Incidence: the two peak years were 1983 and 1986;

- 2) Season index: the most common months for their occurrence are January, October, November, and December;
- 3) Cause: except unknown origin, electric wires, wildness burned, smoking, cooking, and mechanical abrasion;
- 4) Duration: first: from 1,200 to 1,500; second: from 1,500 to 1,800; and
- 5) Mortality: the mortality depends on the rate of increase in number of people in the city: rate: Taiwan: 2.77%, Taipei: 6.18%, Kaohsiung: 4.42%.
-

NOTES