

## Review

## Experiences of in-patient mental health services: systematic review

Sophie Staniszewska, Carole Mockford, Greg Chadburn, Sarah-Jane Fenton, Kamaldeep Bhui, Michael Larkin, Elizabeth Newton, David Crepez-Keay, Frances Griffiths and Scott Weich

**Background**

In-patients in crisis report poor experiences of mental healthcare not conducive to recovery. Concerns include coercion by staff, fear of assault from other patients, lack of therapeutic opportunities and limited support. There is little high-quality evidence on what is important to patients to inform recovery-focused care.

**Aims**

To conduct a systematic review of published literature, identifying key themes for improving experiences of in-patient mental healthcare.

**Method**

A systematic search of online databases (MEDLINE, PsycINFO and CINAHL) for primary research published between January 2000 and January 2016. All study designs from all countries were eligible. A qualitative analysis was undertaken and study quality was appraised. A patient and public reference group contributed to the review.

**Results**

Studies (72) from 16 countries found four dimensions were consistently related to significantly influencing in-patients' experiences of crisis and recovery-focused care: the importance

of high-quality relationships; averting negative experiences of coercion; a healthy, safe and enabling physical and social environment; and authentic experiences of patient-centred care. Critical elements for patients were trust, respect, safe wards, information and explanation about clinical decisions, therapeutic activities, and family inclusion in care.

**Conclusions**

A number of experiences hinder recovery-focused care and must be addressed with the involvement of staff to provide high-quality in-patient services. Future evaluations of service quality and development of practice guidance should embed these four dimensions.

**Declaration of interest**

K.B. is editor of *British Journal of Psychiatry* and leads a national programme (Synergi Collaborative Centre) on patient experiences driving change in services and inequalities.

**Keywords**

In-patient; mental health services; experiences; systematic review.

**Copyright and usage**

© The Royal College of Psychiatrists 2019.

Patient experience is a vital source of evidence that can drive the provision of high-quality health services.<sup>1,2</sup> Mental health in-patients report a range of experiences including fear of assault, concerns regarding coercion, limited recovery-focused support and lack of therapeutic activities.<sup>3–8</sup> A triennial review of mental health services in England by the Care Quality Commission (2017)<sup>9</sup> highlighted several serious concerns about in-patient care, including wards located in older buildings not designed to meet the needs of acute patients, unsafe staffing levels and overly restrictive care in wards far from patients' homes and families.

The National Health Service (NHS) is under pressure to deliver timely, effective and affordable care with increasingly constrained resources. The National Institute for Health and Care Excellence, the NHS National Quality Board and others have restated core principles of patient-centred care including dignity, compassion, choice and autonomy,<sup>3–5,8</sup> and called for a strengthening of the patient voice. Healthcare providers are now required to collect data to assess patients' experiences of care.<sup>9–12</sup> However, the impact of this data collection on services is unclear<sup>13</sup> because of: the diverse and poor-quality feedback methods,<sup>14</sup> a lack of consensus about which experiences are most salient (and hence should be asked about), and limited evidence about how patient experience data can guide service improvements.<sup>13,15</sup> Such challenges highlight the need for robust evidence to inform best practice, with clarity about the experiences of most importance to patients. In response to this need, this systematic review aimed to identify the most salient experiences of people using in-patient mental healthcare to inform the provision of high-quality services.

**Method**

The review was divided into a scoping review to ascertain the nature and size of the evidence base, and the main systematic review.

**Protocol and registration**

The EURIPIDES (Evaluating the Use of Patient Experience Data to Improve the Quality of Inpatient Mental Health Care) systematic review was registered in 2016 on PROSPERO: CRD42016033556.

**Scoping review**

Before the systematic review, a scoping review was conducted to ascertain the extent, range and nature of studies to map emerging key themes without describing the findings in full or performing a quality check<sup>16</sup> and to inform the main review. Six key authors known to be experts in mental health patient experience were contacted for new or unpublished reports and studies.

**Patient and Public Involvement Reference Group**

The Patient and Public Involvement Reference Group (PIRG) included 10 service users, recruited by the Mental Health Foundation, with experience of in-patient care or caring for someone who had been an in-patient. They were invited to two meetings: first, to obtain their views on the themes identified in the scoping review, with the potential to add further concepts they felt had not been identified; and second, to obtain their opinions on themes identified in the main systematic review and to contribute to the interpretation

**Table 1** Reporting Patient and Public Involvement in the EURIPIDES study using GRIPP2

GRIPP2 <sup>a</sup> Short Form item	Description
Aims: Report the aim of Patient and Public Involvement in the study	(a) Ensure there is a patient voice included at all stages of the EURIPIDES study; (b) to discuss the scoping study themes and to identify additional ones service users feel are important; (c) to discuss the themes and sub-themes identified in the main review to ensure face and content validity.
Methods: Provide a clear description of the methods used for Patient and Public Involvement in the study	The Patient and Public Involvement Reference Group was established by the Mental Health Foundation. Members were varying in background and experience. The Group met regularly and at key points during the study. The group were facilitated by D.C.-K. who ensured they felt able and were supported to contribute and challenge methods.
Study result outcomes: Report the results of Patient and Public Involvement in the study, including both positive and negative outcomes	The Patient and Public Involvement Reference Group provided a strong patient and carer perspective. They critiqued the content of the themes identified in the scoping review, identifying additional areas such as boredom. They provided content and face validity of the themes and sub-themes identified in the main review. They provided real life examples of the themes from their own experiences. The Patient and Public Involvement Reference Group also checked if the themes from international studies resonated in a UK context.
Discussion and conclusion outcomes: Comment on the extent to which Patient and Public Involvement influenced the study overall. Describe positive and negative effects	The Patient and Public Involvement Reference Group was important in confirming the systematic review had identified the themes of importance to patients and carers. This was particularly important because the strength of the patient voice was uncertain in the papers reviewed.
Reflections/critical perspective: Comment critically on the study, reflecting on the things that went well and those that did not, so others can learn from this experience	The Patient and Public Involvement Reference Group worked well in the study. On reflection more embedded forms of involvement, with members of the group working more closely on the analysis, may have embedded the service user voice more strongly into the study and could have created the conditions for the co-production of knowledge and possibly additional sub-themes.

a. Staniszewska S, Brett J, Simera I, Seers K, Mockford C, Goodlad S, Altman DG, Moher D, Barber R, Denegri S, Entwistle A, Littlejohns P, Morris C, Suleman R, Thomas V, Tysall C. GRIPP2 reporting checklist: tools to improve reporting of patient and public involvement in research. *BMJ* 2017; **358**: j3453.

of our findings. A full description of the patient involvement in the study is reported using the GRIPP2 Short Form Checklist in Table 1.

### Identification of studies for the systematic review

Guided by the themes that emerged from the scoping review, search terms and a search strategy were developed and applied to the databases MEDLINE, CINAHL and PsycINFO. An example of search terms and results is reported in Fig. 1. Reference lists of included papers were scanned. The search deviated from the protocol in that only three of five databases were searched due to the large numbers of abstracts retrieved (Web of Science and Embase were not used).

### Inclusion and exclusion criteria

All study designs were considered if papers included experiences of current or former in-patients of mental health institutions. No restrictions were applied based on country. Articles were included if they reported primary research, were peer reviewed and published in English between January 2000 and January 2016. Papers were excluded if they were not primary studies, based on pre-2000 data, included children and adolescents (aged under 18 years) or were not in the English language. Where study participants included both in- and out-patients, only data regarding in-patient experiences were extracted. Reviews (Table A.1) were noted and reference lists scanned, but excluded from the review to avoid bias.

### Study selection

Titles and abstracts were screened (C.M., G.C.), 20% of which were independently cross-checked for agreement before obtaining full-text articles (S.S. and C.M.). Full texts were obtained where the abstract was unclear. Any disagreements could be resolved by consensus (C.M., G.C. and S.S.) but no disagreements occurred.

### Data extraction

The data extracted, using Microsoft Excel (version 2013), included citation details, sample recruitment and research methods,

findings related to key concepts and any other emerging concepts (C.M.).

### Quality and risk of bias in individual studies

The quality of the studies were evaluated by the Critical Appraisal Skills Programme (CASP) Qualitative Checklist,<sup>17</sup> undertaken by C.M. Because of the heterogeneity of the included studies, many of which were descriptive in their approach, this checklist provided an appropriate basis for comparison between studies. The only question change in the CASP checklist was from 'Is the qualitative methodology appropriate for this study?' to 'Is the methodology appropriate for this study?'

### Data analysis

The scoping review informed the development of a thematic framework, which guided but did not restrict the review. A narrative synthesis of the themes was undertaken.<sup>18</sup> As the researcher read each study, an initial preliminary synthesis of the study was undertaken and emerging sub-themes were identified. The researcher was then able to compare themes and sub-themes within and across studies and further develop them into the main themes. Themes were summarised in a descriptive form, allowing for the findings of all review studies, regardless of study design, to be aggregated and summarised. We used the concept of data saturation to help us decide when to complete data extraction. Saturation of data is judged to have happened at a point where no new themes are being identified in the studies when compared with what has already been extracted.<sup>7</sup> It is a useful approach for large reviews where the addition of further papers is unlikely to change key findings.

## Results

### PPIRG

Key themes identified in the scoping review were discussed in detail by group members who critiqued their content and identified additional areas such as boredom. The PPIRG provided content and

Results	Search type	Actions
1	exp Inpatients/ or inpatient*.mp.	73 820
2	service user*.mp.	2556
3	patient/	17 869
4	exp "Commitment of Mentally Ill"/	6286
5	involuntary.mp.	10 996
6	1 or 2 or 3 or 4 or 5	108 766
7	exp Hospitals, Psychiatric/ or psychiatric.mp.	218 311
8	psychiatry.mp. or Psychiatry/	74 187
9	Mental Disorders/	139 896
10	7 or 8 or 9	341 433
11	exp Patient Satisfaction/	67 505
12	(satisf* or experience*).mp. [mp=title, abstract, original title, name of substance word, subject heading word, keyword heading word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier]	930 899
13	11 or 12	933 891
14	6 and 10 and 13	3204
15	limit 14 to yr="2000–Current"	2181
16	limit 15 to english language	1943

Fig. 1 Example of search strategy from MEDLINE.

face validity for the identified themes and provided real-life examples of the themes from their own experiences. The PPIRG also provided an opportunity to check if the themes identified from international studies resonated in a UK context.

### The systematic review

A total of 4979 abstracts were screened and 116 papers fulfilled the inclusion criteria (Fig. 2). Two consecutive sifts were conducted

due to an error in the first search of the PsycINFO database omitting 2980 hits which was identified after the first sift was completed. The first sift of 1999 hits resulted in 72 relevant papers for the review; 11 papers were from same studies.<sup>19–29</sup> Following this, the second sift of 2980 abstracts resulted in an additional 44 studies fitting the criteria (total  $n = 116$ ). Drawing on the principles of data saturation,<sup>30</sup> additional studies that repeated themes already identified were excluded from the main review. In total, eight studies added new themes and were included at this stage.




Fig. 2 PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) 2009 flow diagram.

A total of 16 systematic reviews (Table A.1) which investigated in-patient experience were identified. In total, 72 studies were included in the review, a third of which were from the UK<sup>24–47</sup> ( $n = 24$ )<sup>19–21,25,27,31–49</sup> (Supplementary Table 1 available at <https://doi.org/10.1192/bjp.2019.22>). Although studies using qualitative methods were most common (Table A.1), studies using patient experience questionnaires and patient record data were also included. The CASP checklist identified many of the papers as being of medium to poor quality.

### Timing of data collection in included studies

Little information was provided about the timing of data collection in over a third of papers (37%), other than describing participants as in-patients at the time.<sup>25–27,31,32,35,36,43,44,48–63</sup> Data were mostly collected just before,<sup>28,29,45,64–73</sup> immediately after discharge<sup>20,45,59,74,75</sup> or from former in-patients.<sup>22,23,34,37–39,41,42,46,47,63,76–80</sup> This suggests that patients were recovering when experiences were elicited. In three studies, data collection coincided with a ward event (e.g. refurbishment).<sup>81–83</sup> A number of studies ( $n = 12$ , 17%) collected data shortly after an event such as admission,<sup>19,21,84–86</sup> seclusion, sedation or restraint.<sup>24,33,87–92</sup>

### Identification of key themes

Patient experiences were categorised into four overarching themes or dimensions of experience: the importance of high-quality relationships; averting negative experiences of coercion; a healthy, safe and enabling physical environment and ward milieu; and authentic experiences of patient-centred care. These key themes accompanied by sub-themes are described in detail below.

#### The importance of high-quality relationships

The importance of high-quality relationships was the most consistently reported theme.

Important factors in developing such relationships with staff included being treated with respect, feelings of stability, recognising empathy and high-quality communication<sup>19,23,24,27,28,35,36,38,39,51,55,60,61,63,78,87,90</sup> with staff who patients felt were trustworthy, reliable<sup>35,63,69</sup> or helpful.<sup>27,51,54,62</sup> Good staff–patient relationships facilitated the in-patient care pathway in mental health institutions<sup>28,35,39,51,68</sup> and reduced the use of coercive measures.<sup>35,45,78</sup> Ward rounds were an important setting for staff–patient interaction and patients reported these as helpful and informative.<sup>44</sup>

Potential barriers to therapeutic relationships included: gender-specific problems – male nursing staff were not welcome if the patient had a history of abuse by male perpetrators<sup>36,78</sup> or where gender-specific cultural barriers existed (e.g. a Muslim woman supervised by a male nurse);<sup>68</sup> lack of meaningful communication – where communication was compromised due to differences in culture, language, religion,<sup>34,39,57,68</sup> through use of coercive measures<sup>33,60</sup> or where technical language used by staff was not easily understood;<sup>19</sup> absence of regular ward staff – patients were upset by the absence of regular ward staff due to office duties, shift working, reliance on temporary staff<sup>23,24,27,28,35–37,39,45,46,51,54,55,63,69</sup> and having extended waits to speak to staff<sup>24,36,46,54,77,80,82</sup> particularly at ward rounds;<sup>43</sup> poor staff attitude – where patients complained that staff ignored them,<sup>57,87,88,91</sup> displayed indifference<sup>24</sup> or insufficient understanding of patients;<sup>78</sup> inconsistent staff behaviour – reports of staff interpreting ward rules inconsistently, causing confusion;<sup>19,23,27,31,33,36,46,49,82,91</sup> staff abuse – some patients reported abuse by staff, including provocation, bullying, shouting or belittling of patients.<sup>19,23,27,28,33,39,56,62,78,79,83,87,88</sup>

### Relationships with other patients and with relatives:

Patients relied on other patients for information about ward activities and rules, to share experiences and when debriefing after group sessions.<sup>22,45,77,82,83</sup> However, arguments and violence between patients<sup>36,39,48</sup> generated fear and isolation for some, causing them to retreat to their rooms for safety or to abscond.<sup>23,37,39,49,65,80</sup>

Isolation from family caused distress. Patients reported that having a friend or family member with them would have helped with orientation<sup>79</sup> and they could have helped staff with assessments and treatment plans.<sup>22,38,53</sup> However, family members felt left out of decision-making about care.<sup>92</sup>

### Averting negative experiences of coercion

The second main theme was concerned with experiences of coercion. All patients expected to be treated as ‘normal human beings’<sup>24,29,77</sup> and addressed professionally, including during restraint.<sup>87</sup> Patients wanted the reasons for coercive measures to be communicated so they could understand them as this helped some patients trust staff and feel safe.<sup>46,67,75,79,87</sup> Patients valued persuasion over threats of force<sup>60</sup> and coercion,<sup>78</sup> which could bring back memories of past history of violence and neglect.<sup>33,88,89</sup>

Where coercive measures were discussed in the studies, these included experiences of sedation, seclusion and restraint. It has been reported that Black and minority ethnic patients are more likely to experience coercion than White patients.

**Ethnicity:** Two studies examined the commonly held perception that Black and minority ethnic patients experienced more coercion on admission than other patients.<sup>21,74</sup> The findings were not conclusive: although hospitals in the UK with higher proportions of Black and minority ethnic patients employed more coercive practices, this was independent of individual patient ethnicity.<sup>21,74</sup>

**Sedation:** Some patients recognised that medication was important for the in-patient care pathway.<sup>20,39,41</sup> Some trusted staff to decide on appropriate sedation,<sup>32,52</sup> whereas others felt empowered to decide on timing and dose of medication when administered on an ‘as needed’ basis.<sup>32</sup> However, patients also voiced concerns that included lack of communication about consent, information about medication and advanced wishes;<sup>39,52</sup> lack of confidentiality regarding medication;<sup>32,42</sup> perceived overmedication<sup>32,39,41,46,47,52,69</sup> (including overlooked or ignored reports of side effects);<sup>28,41</sup> and fear of harm

during forced medication,<sup>20,32,39,54,60,78</sup> for example patients in crisis reported a fear of being raped by staff or of dying.<sup>20,41,78,88</sup>

**Seclusion:** Some patients reported seclusion as helpful or necessary<sup>24,57,79,88</sup> and that they felt safe as staff were nearby.<sup>24,57,88,90</sup> Patient concerns included having insufficient information about the reasons for seclusion<sup>23,24,46,57,88</sup> before or after the event.<sup>24,57</sup> Seclusion was perceived as a punishment<sup>79</sup> and associated with limited contact;<sup>57,88</sup> lack of concern by staff;<sup>89</sup> degradation and humiliation, e.g. lack of facilities<sup>24,57,89</sup> or being stripped of clothing in front of staff members;<sup>61,79,89,91</sup> and violation of rights<sup>88</sup> and dignity.<sup>61</sup>

**Restraint:** Restraint was described as forcible manual or mechanical restraint and typically involved several staff, mostly nurses<sup>23,60,78,88,92</sup> but occasionally security staff.<sup>78,92</sup> Restraint was described negatively<sup>25,33,78</sup> and fear of restraint prevented patients from seeking help earlier.<sup>33</sup> There was a risk of harm if mechanical restraints were used,<sup>87</sup> although these were not used in all countries. Talking with staff following restraint or being allowed to examine records of the event was considered helpful.<sup>33</sup>

In addition to the use of coercive measures, patients also described perceived punishment by staff<sup>19,35,41,80,91</sup> in the form of the removal of leave entitlements,<sup>35</sup> removal of furniture and personal items<sup>41,91</sup> and not being able to stay up in the evening.<sup>19,80</sup> Patients described this as a violation of their rights.<sup>23,57,58,88</sup>

### A healthy, safe and enabling physical environment and ward milieu

The third main theme focused on a healthy, safe and enabling environment. This contributed to how relatives felt when visiting,<sup>92</sup> how patients felt about themselves<sup>39</sup> and how they reacted to treatment.<sup>36,39,42</sup> Johansson *et al* (2003)<sup>63</sup> argued that the physical environment was as important to patients as receiving satisfactory care. A number of studies reported that patients saw hospital as a ‘sanctuary’<sup>80</sup> or a ‘safe space’<sup>62</sup> where they could have time to reflect away from day-to-day stressors,<sup>38,50</sup> be kept safe<sup>19,39,48,54</sup> and experience a caring, therapeutic environment.<sup>80</sup>

Patients felt that their in-patient care pathway was aided by connection to the ‘real world’<sup>61</sup> and that being made to feel ‘normal’<sup>24,28,51,77</sup> was important. This included being allowed to walk around hospital grounds.<sup>39,80</sup> Older establishments often had extensive grounds and patients reported that access to these spaces resulted in less need for medication.<sup>32</sup> Access to a place of worship was comforting,<sup>51,68</sup> as was freedom to make small decisions<sup>31,41</sup> such as making snacks<sup>62</sup> or hot drinks.<sup>36</sup> Private bedrooms were important,<sup>80</sup> being near windows enabled ward-bound patients to enjoy the outside and fresh air,<sup>83</sup> and appropriate use of colour was described as conducive to recovery.<sup>80</sup> An environment where staff and patients mixed together reduced feelings of stigma<sup>51</sup> and encouraged favourable interactions.<sup>63</sup>

Patients reported several environmental problems that were not conducive to recovery-focused care. Some of these were associated with arguments and violence between patients.<sup>36,39,48</sup> Other environmental problems included noise from doorbells, alarms and telephones.<sup>82</sup> Poor positioning of the nurses’ stations often created physical divisions between patients and staff, reducing interaction.<sup>61,80,92</sup> Communal spaces sometimes lacked privacy for visiting relatives or opportunities for physical activity,<sup>49</sup> especially for those under close observation.<sup>92</sup>

There were also contradictory reports. In several studies, some patients described hospital as a place of confinement rather than therapy.<sup>19,29,36,37,39,42,80</sup> There were analogies with prison<sup>29,36,39,42,80</sup> and punishment.<sup>37,39</sup> This was particularly so in

secure units with a lack of outside space<sup>39</sup> and where more patients were admitted compulsorily.<sup>29</sup>

**Ward milieu:** Related to environment was the experience of ward milieu, which was shaped by the conduct of staff. Staff provided structure, order and safety<sup>82</sup> and were responsible for creating a congenial atmosphere.<sup>54</sup> Feeling safe was a prime concern to patients<sup>48,65</sup> who perceived wards to be safe when they viewed staff as trustworthy,<sup>35</sup> caring and supportive.<sup>35,38</sup> Wards were sometimes criticised as being too busy<sup>36,49,54</sup> and reactive to events such as restraint,<sup>56,79,92</sup> seclusion<sup>91</sup> or violence.<sup>23,58,80</sup> Patients felt vulnerable to the latter,<sup>23,37,39</sup> fearful of other patients<sup>49,78</sup> and worried about security of belongings.<sup>36,65,80</sup> Fear contributed to withdrawing within the ward<sup>49,81</sup> or leaving hospital.<sup>37,80</sup>

Ward routines also shaped patients' experiences. The day<sup>51</sup> was often structured to include individual and group therapies as well as other activities, e.g. puzzles, conversation or listening to music.<sup>92</sup> Evenings were typically less structured.<sup>51</sup> Some patients relished the leisure time<sup>24,38,50,54</sup> and some took this as a time for personal reflection.<sup>38,51,57</sup> However, others were uneasy<sup>38,51</sup> and reported insufficient<sup>36,49</sup> activity.<sup>23,24,39,49, 68</sup> The location of the hospital – being close to family – was important to patients<sup>79</sup> and they appreciated the inclusion of, and support from, families.<sup>22,38,53</sup>

**Boredom:** 'Boredom' or having little to do was mentioned in several studies.<sup>23,24,27,41,51,54,59,68,80,82,83,91</sup> Patients suggested that inactivity slowed the in-patient care pathway,<sup>59</sup> reduced self-efficacy,<sup>41</sup> exacerbated symptoms<sup>80</sup> and was related to aggression and violence on the ward.<sup>23</sup> Some patients reported that inactivity encouraged poor health outcomes, e.g. saying that they would eat, sleep or smoke but not exercise.<sup>24,59,80,83</sup>

#### Authentic experiences of patient-centred care

The final theme brought together a collection of sub-themes focused on authentic experiences of patient-centred care, which included shared decision-making, sensitivity to gender and culture and the provision of information.

**Shared decision-making:** Two studies reported that patients' involvement in treatment decisions was associated with positive experiences of care.<sup>50,65</sup>

**Gender and cultural differences:** Patients wanted to be understood and seen as individuals, and this was framed in respect of their gender, ethnicity and religion.<sup>33,34,68,78</sup> Some patients described cultural differences in perceptions of privacy, and reported concern that staff had not recognised or responded to their discomfort in accepting care from differently gendered staff,<sup>68</sup> for example during restraint and sedation,<sup>33</sup> or for women with a history of sexual abuse by male perpetrators.<sup>78</sup> More positively, female patients tended to prefer single-gender wards (where they felt safer<sup>36</sup>). Where this was not available, female patients were satisfied on mixed wards if they had access to a quiet room, if their privacy was respected and if they had access to personal hygiene products.<sup>81</sup> Faith also mattered: prayer and rituals (e.g. hand washing) offered comfort to some patients<sup>68</sup> but were not always understood or accommodated by staff.<sup>34</sup>

**Provision of information:** There were several reports in which patients felt they had not received sufficient information about their diagnosis,<sup>23,65,69,87</sup> treatment,<sup>20</sup> treatment plan,<sup>23,32,52,57,60,65,69,87,88,90,91</sup> choices or rights.<sup>20,46,53,64,86</sup> Timing was also important as patients found it difficult to understand or remember this information when unwell.<sup>45,69</sup>

## Discussion

The aim of this review was to identify the most salient aspects of in-patient experience to support improvements in care in ways that are conducive to recovery-focused care. To the best of our knowledge this is the largest review of its type in the UK and internationally, with 72 included studies, of which a third were from the UK. A strength of the review was the involvement of the PPIRG who provided important face and content validity checks and were able to identify additional areas of experience, such as boredom, which could be built into the main review.

The review makes an important contribution to the field of mental health in-patient experiences through the identification of four key, interlinked themes: the importance of high quality relationships; averting negative experiences of coercion; a healthy, safe and enabling physical environment and ward milieu; and authentic experiences of patient-centred care. These themes and their associated sub-themes represent the active ingredients of a high-quality mental health in-patient experience (as well as the common causes of very poor experiences). The identified themes can be used to design and deliver high-quality services, provide content for the development of robust patient experience questionnaires or inform qualitative methods that aim to evaluate salient aspects of patient experience. They provide evidence for the development of practice guidance that supports the implementation of high-quality services.

A consistent thread across all four themes was the key role of staff in facilitating a high-quality patient experience. However, staff operate within the context of a wider system that needs to support the delivery of care. It was not always possible to understand this context from the studies reviewed as many did not provide such wider contextual information. This would have been useful, particularly in understanding why some studies reported very negative experiences and others reported more positive experiences. Future studies might consider reporting contextual information to aid interpretation.

It is important to note that the findings of studies relating to discharge appeared to be influenced by the research design, with questionnaires identifying high levels of satisfaction whereas experiences captured using qualitative methods were described differently. Future studies should pay careful attention to the way in which design might affect the reporting of experiences.




## Limitations

A limitation of this review, common to all secondary research, is that it is reliant on the conduct and content of primary studies which may have included biases that we could not account for. Few studies mentioned the involvement of patients in data collection<sup>20,39,46,79</sup> and research design,<sup>20,27,39,46,79</sup> and the study authors' professional perspective is often unreported. It is therefore unclear to what extent a study finding reflects the patient voice or whether it predominantly reflects the researchers' interpretation of their data. Ensuring greater clarity about whose voice is represented, as a means of minimising bias, represents an important methodological challenge for future research. In future reviews, the case could be made to focus on studies where there is evidence of a strong patient voice in the conduct and interpretation of the study.

Although we used the concept of data saturation to decide when to stop data extraction, it is always possible that other papers contained nuances in themes that were unintentionally omitted. The risk of bias in this review may have been partially mitigated by our scoping review which identified key authors and included a citation search of their papers and other literature reviews. In addition,

the PPIRG provided important assurance of face and content validity.

Our study relies on secondary analysis of qualitative data. The findings we have presented are drawn from the reports from participants in primary studies. Many of these claims (e.g. the perceived role of good relationships in reducing a range of unwanted outcomes, the role of boredom in exacerbating those outcomes) are reported across multiple primary sources. However, an important limitation of secondary research involves the gaps that exist in studies. A key gap in this review was the lack of experiences from people of Black and minority ethnic groups, which appears to be under-researched. Future studies should ensure they build ethnicity into their design.

**Sophie Staniszewska**, DPhil (Oxon), BSc (Hons), Professor of Patient and Public Involvement and Experiences of Care, Division of Health Sciences, Warwick Medical School, University of Warwick, UK; **Carole Mockford** , BSc (Hons), MA, DPhil (Oxon), Senior Research Fellow, Division of Health Sciences, Warwick Medical School, University of Warwick, UK; **Greg Chadburn**, WMS, BSc, MBChB, Researcher, Surrey and Sussex Healthcare National Health Service Trust, UK; **Sarah-Jane Fenton** , BSc (Hons) Geography, MPhil (Oxon), Comparative Social Policy, PGCert Advanced Research Methods and Skills, PhD Social Policy, Research Fellow, Division of Health Sciences, Warwick Medical School, University of Warwick, UK; **Kamaldeep Bhui** , BSc (Pharmacy), MBBS, MSc (Epid), MSc (Mental Health Studies), MD (Res), FRCPsych, Dipl. Clin. Psychotherapy, Dip. LSHTM, Professor of Psychiatry, Centre for Psychiatry, Wolfson Institute of Preventive Medicine, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, East London National Health Service Foundation Trust, UK; **Michael Larkin**, PhD, Reader in Psychology, School of Life and Health Sciences, Aston University, UK; **Elizabeth Newton**, BSc, DClinPsy, Consultant Clinical Psychologist, Research Fellow, School of Psychology, University of Birmingham, UK; **David Crepaz-Keay**, DProf, Head of Empowerment and Social Inclusion, Mental Health

Foundation, UK; **Frances Griffiths**, MB, BS, PhD, Professor of Medicine in Society, Division of Health Sciences, Warwick Medical School, University of Warwick, UK; **Scott Weich**, MBBS, MSc, MD, FRCPsych, Professor of Mental Health, Division of Health Sciences, Warwick Medical School, University of Warwick; and School of Health and Related Research (SchARR), University of Sheffield, UK

**Correspondence:** Sophie Staniszewska, Warwick Research in Nursing, Warwick Medical School, Division of Health Sciences, University of Warwick, Coventry CV4 7AL, UK. Email: [sophie.staniszewska@warwick.ac.uk](mailto:sophie.staniszewska@warwick.ac.uk)

First received 22 Dec 2017, final revision 19 Dec 2018, accepted 6 Jan 2019

## Funding

S.S. is part-funded by NIHR CLAHRC West Midlands. K.B. was (in part) supported by the National Institute for Health Research (NIHR) Collaboration for Leadership in Applied Health Research and Care (CLAHRC) North Thames at Bart's Health NHS Trust. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

## Supplementary material

Supplementary material is available online at <https://doi.org/10.1192/bjp.2019.22>

## Acknowledgements

We thank the Patient and Carer Reference Group members, who have been key to the development of this review. We thank them for all their inputs and insights. The results are stronger through their involvement.

## Appendix

**Table A.1** Other systematic reviews

Author	Date	Focus of review	Number studies	Years searched	Key findings: headings or themes from findings
Binnema <sup>93</sup>	2004	Psychiatric patients and boredom	Not stated but ?18	Appears to be 1994–2003	Boredom is a lack of experience of meaning; many psychiatric patients experience boredom and lack opportunities to experience meaning. This indicates a lack in the therapeutic potential of the hospital environment which needs to change
Cutcliffe <i>et al</i> <sup>94</sup>	2015	Evaluations of in-patient mental health care experiences in six countries	Not stated	Not stated	Convergence and congruence in patient experience evaluations, overall disturbing picture of in-patient mental healthcare, major disconnect between policy and practice, problems caused by a multitude of variables, can learn from therapeutic relationships
Duncan <i>et al</i> <sup>95</sup>	2010	Cochrane review: Shared decision-making interventions for people with mental health conditions	2	All to 2008	Effects of interventions: clinical outcomes; health service-related outcome: rate of readmission to hospital; secondary outcomes: level of consumer involvement in decision-making process, consumer satisfaction with information provided, provider satisfaction, consumer concordance with treatment plan, consultation time
Ford <i>et al</i> <sup>96</sup>	2015	Experience of compulsory treatment and implications for recovery-orientated practice	5	2000 onwards	Views of the justification of compulsory detention, power imbalance, lack of information or choice
Gerolamo <sup>97</sup>	2004	Patient outcomes after treatment in acute care psychiatric hospitals and wards	47	1991–2004	Readmission, rehospitalisation, recidivism; symptom and function improvement; client satisfaction; suicide and self-injury
Hopkins <i>et al</i> <sup>98</sup>	2009	Responsiveness as context to understand patient perceptions and expectations to in-patient mental healthcare	10	1998–2008	Respect for dignity, confidentiality, autonomy, prompt attention, amenities, access to social networks, choice of provider
Katsakou and Priebe <sup>99</sup>	2007	Patient experiences of involuntary hospital admission and treatment	5	Selected papers are from 2001–2003	Lack of autonomy and not included in decision-making, quality of care and not being cared for, emotional impact of involuntary treatment and feeling devalued, respect and autonomy, being cared for and treatment benefits, being a human being like other people

(Continued)

Table A.1 (Continued)

Author	Date	Focus of review	Number studies	Years searched	Key findings: headings or themes from findings
McHale and Felton <sup>100</sup>	2010	Factors affecting attitudes towards self-harm	19	Papers from 1998–2009	Lacking education/ training, role expectations and clinical culture, perception of health needs, knowledge of self-harm, education and training use, dissatisfaction with care
Maatta <sup>101</sup> (abstract only available)	2009	Exploring male and female patients' experiences of psychiatric hospital care: a critical analysis of the literature	5	Not in abstract	Treatment specifically related to women, to keep a facade, and single-gender or mixed ward
Newman et al <sup>102</sup>	2015	Mental health patients' experience of mental healthcare	34	2008–2012	Acknowledging a mental health problem and seeking help, building relationships through participation in care, working towards continuity of care
Omer et al <sup>103</sup>	2015	Continuity of care versus specialist systems	21	1985–2013	Hospitalisation, length of stay, transition of care, and staff and patient views. With regard to patient views, there were more positive reports for continuity of care
Sequeira and Halstead <sup>104</sup>	2002	Restraint and seclusion	23	1975–2001	Client's experience of seclusion, restraint
Strout <sup>105</sup>	2010	Experience of being physically restrained	12	1966–2009	Negative psychological impact, re-traumatisation, perceptions of unethical practices, broken spirit
Sturrock <sup>106</sup>	2010	Experiences of restraint in in-patient areas	5	2000–March 2009	Distressing; should be debriefed; can lead to potentially abusive situations; engendered fear, anxiety and rage; incidents could be prevented
Van der Merwe et al <sup>107</sup>	2009	Views on locked doors	11	Up to 2008	Advantages of locked doors; disadvantages of locked doors by patients, by staff; aggressive incidents and the door status; patients' satisfaction with treatment and the door status; patients' symptoms and the door status.
Van Der Merwe et al <sup>108</sup>	2013	Improving seclusion practice – staff and patient views	39	1960–2006	Patient and staff perception of seclusion, improvement suggestions

## References

- Rethink. *Future Perfect: Mental Health Service Users Set out a Vision for the 21st Century*. Rethink, 2005.
- Royal College of Psychiatrists. *Improving Inpatient Mental Health Services for Black and Minority Ethnic Patients: Recommendations to Inform Accreditation Standards*. Royal College of Psychiatrists, 2010.
- MIND. *Listening to Experience: An independent Inquiry into Acute and Crisis Mental Healthcare*. MIND, 2011.
- Commission for Healthcare Audit and Inspection. *The Pathway to Recovery: A Review of NHS Acute Inpatient Mental Health Services*. Commission for Healthcare Audit and Inspection, 2008.
- Care Quality Commission. *Monitoring the Mental Health Act in 2012/13*. Care Quality Commission, 2014.
- Commission for Healthcare Audit and Inspection. *Count me in 2008: Results of the 2008 National Census of Inpatients in Mental Health and Learning Disability Services in England and Wales*. Commission for Healthcare Audit and Inspection, 2008.
- Staniszewska S, Boardman F, Gunn L, Roberts J, Clay D, Seers K, Brett J, Avital L, Bullock I, O'Flynn N. The Warwick Patient Experiences Framework: patient-based evidence in clinical guidelines. *Int J Qual Health Care* 2014; **26**: 151–7.
- National Institute for Health and Care Excellence (NICE). *Service User Experience in Adult Mental Health: Improving the Experience of Care for People Using Adult NHS Mental Health Services*. NICE, 2011.
- Care Quality Commission. *The State of Care in Mental Health Services 2014 to 2017*. Care Quality Commission, 2017.
- Curtis S, Gesler W, Priebe S, Francis S. New spaces of inpatient care for people with mental illness: a complex 'rebirth' of the clinic? *Health Place* 2009; **15**: 340–8.
- Barbato A, Bajoni A, Rapisarda F, D'Anza V, De Luca LF, Inglese C, et al. Quality assessment of mental health care by people with severe mental disorders: a participatory research project. *Community Ment Health J* 2014; **50**: 402–8.
- Bramesfeld A, Klippel U, Seidel G, Schwartz FW, Dierks M. How do patients expect the mental health service system to act? Testing the WHO responsiveness concept for its appropriateness in mental health care. *Soc Sci Med* 2007; **65**: 880–9.
- Coulter A, Locock L, Ziebland S, Calabrese J. Collecting data on patient experience is not enough: they must be used to improve care. *BMJ* 2014; **348**: g2225.
- National Health Service (NHS) England. *NHS England Review of the Friends and Family Test*. NHS England, 2014.
- Boiko O, Campbell JL, Elmore N, Davey AF, Roland M, Burt J. The role of patient experience surveys in quality assurance and improvement: a focus group study in English general practice. *Health Expect* 2015; **18**: 1982–94.
- Arksey H, O'Malley R. Scoping studies: towards a methodological framework. *Int J Soc Res Methodol* 2005; **8**: 19–32.
- Critical Appraisal Skills Programme (CASP). *CASP Qualitative Checklist*. CASP, 2007. Available at: <http://www.casp-uk.net/>.
- Sandelowski M, Voils CI, Barroso J. Defining and designing mixed research synthesis studies. *Res Sch* 2006; **13**: 29.
- Katsakou C, Marougka S, Garabette J, Rost F, Yeeles K, Priebe S. Why do some voluntary patients feel coerced into hospitalisation? A mixed-methods study. *Psychiatry Res* 2011; **187**: 275–82.
- Katsakou C, Rose D, Amos T, Bowers L, McCabe R, Oliver D, et al. Psychiatric patients' views on why their involuntary hospitalisation was right or wrong: a qualitative study. *Soc Psychiatry Psychiatr Epidemiol* 2012; **47**: 1169–79.
- Bennewith O, Amos T, Lewis G, Katsakou C, Wykes T, Morriss R, et al. Ethnicity and coercion among involuntarily detained psychiatric in-patients. *Br J Psychiatry* 2010; **196**: 75–6.
- Kauppi K, Hätönen H, Adams CE, Välimäki M. Perceptions of treatment adherence among people with mental health problems and health care professionals. *J Adv Nurs* 2015; **71**: 777–88.
- Kontio R, Anttila M, Lantta T, Kauppi K, Joffe G, Valimäki M. Toward a safer working environment on psychiatric wards: service users' delayed perspectives of aggression and violence-related situations and development ideas. *Perspect Psychiatr Care* 2014; **50**: 271–9.
- Kontio R, Joffe G, Putkonen H, Kuosmanen L, Hane K, Holi M, et al. Seclusion and restraint in psychiatry: patients' experiences and practical suggestions on how to improve practices and use alternatives. *Perspect Psychiatr Care* 2012; **48**: 16–24.
- Whittington R, Bowers L, Nolan P, Simpson A, Neil L. Approval ratings of inpatient coercive interventions in a national sample of mental health service users and staff in England. *Psychiatr Serv* 2009; **60**: 792–8.
- Bowers L, Haglund K, Muir-Cochrane E, Nijman H, Simpson A, Van Der Merwe M. Locked doors: a survey of patients, staff and visitors. *J Psychiatr Ment Health Nurs* 2010; **17**: 873–80.
- Stewart D, Burrow H, Duckworth A, Dhillon J, Fife S, Kelly S, et al. Thematic analysis of psychiatric patients' perceptions of nursing staff. *Int J Ment Health Nurs* 2015; **24**: 82–90.



- 28 Wyder M, Bland R, Blythe A, Matarasso B, Crompton D. Therapeutic relationships and involuntary treatment orders: service users' interactions with health-care professionals on the ward. *Int J Ment Health Nurs* 2015; **24**: 181–9.
- 29 Wyder M, Bland R, Herriot A, Crompton D. The experiences of the legal processes of involuntary treatment orders: tension between the legal and medical frameworks. *Int J Law Psychiatry* 2015; **38**: 44–50.
- 30 Dixon-Woods M, Agarwal S, Jones D, Young B, Sutton A. Synthesising qualitative and quantitative evidence: a review of possible methods. *J Health Serv Res Policy* 2005; **10**: 45–53.
- 31 Alexander J. Patients' feelings about ward nursing regimes and involvement in rule construction. *J Psychiatr Ment Health Nurs* 2006; **13**: 543–53.
- 32 Baker JA, Lovell K, Easton K, Harris N. Service users' experiences of 'as needed' psychotropic medications in acute mental healthcare settings. *J Adv Nurs* 2006; **56**: 354–62.
- 33 Bonner G, Lowe T, Rawcliffe D, Wellman N. Trauma for all: a pilot study of the subjective experience of physical restraint for mental health inpatients and staff in the UK. *J Psychiatr Ment Health Nurs* 2002; **9**: 465–73.
- 34 Bowl R. The need for change in UK mental health services: South Asian service users' views. *Ethnicity Health* 2007; **12**: 1–19.
- 35 Chorlton E, Smith I, Jones SA. Understanding how people who use illicit drugs and alcohol experience relationships with psychiatric inpatient staff. *Soc Psychiatry Psychiatr Epidemiol* 2015; **50**: 51–8.
- 36 Cutting P, Henderson C. Women's experiences of hospital admission. *J Psychiatr Ment Health Nurs* 2002; **9**: 705–12.
- 37 Duggins R, Shaw I. Examining the concept of patient satisfaction in patients with a diagnosis of schizophrenia: a qualitative study. *Psychiatric Bull* 2006; **30**: 142–5.
- 38 Fenton K, Larkin M, Boden ZVR, Thompson J, Hickman G, Newton E. The experiential impact of hospitalisation in early psychosis: service-user accounts of inpatient environments. *Health Place* 2014; **30**: 234–41.
- 39 Gilbert H, Rose D, Slade M. The importance of relationships in mental health care: a qualitative study of service users' experiences of psychiatric hospital admission in the UK. *BMC Health Serv Res* 2008; **8**: 92.
- 40 Greenwood N, Hussain F, Burns T, Raphael F. Asian in-patient and carer views of mental health care. Asian views of mental health care. *J Ment Health* 2000; **9**: 397–408.
- 41 Hughes R, Hayward M, Finlay WML. Patients' perceptions of the impact of involuntary inpatient care on self, relationships and recovery. *J Ment Health* 2009; **18**: 152–60.
- 42 Jones A, Crossley D. In the mind of another shame and acute psychiatric inpatient care: an exploratory study. A report on phase one: service users. *J Psychiatr Ment Health Nurs* 2008; **15**: 749–57.
- 43 Labib PLZ, Brownell L. Factors affecting patient satisfaction with the psychiatric ward round: retrospective cross-sectional study. *Psychiatr Bull* 2009; **33**: 295–8.
- 44 Milner G, Jankovic J, Hoosen I, Marrie D. Patients and staff understanding of general adult psychiatry ward rounds. *J Ment Health* 2008; **17**: 492–7.
- 45 Nolan P, Bradley E, Brimblecombe N. Disengaging from acute inpatient psychiatric care: a description of service users' experiences and views. *J Psychiatr Ment Health Nurs* 2011; **18**: 359–67.
- 46 Ridley J, Hunter S. Subjective experiences of compulsory treatment from a qualitative study of early implementation of the Mental Health (Care & Treatment) (Scotland) Act 2003. *Health Soc Care Community* 2013; **21**: 509–18.
- 47 Russo J, Rose D. 'But what if nobody's going to sit down and have a real conversation with you?' Service user/survivor perspectives on human rights. *J Public Ment Health* 2013; **12**: 184–92.
- 48 Stenhouse RC. 'Safe enough in here?': patients' expectations and experiences of feeling safe in an acute psychiatric inpatient ward. *J Clin Nurs* 2013; **22**: 3109–19.
- 49 Kennedy J, Fortune T. Women's experiences of being in an acute psychiatric unit: an occupational perspective. *Br J Occup Ther* 2014; **77**: 296–303.
- 50 Borge L, Hummelvoll JK. Patients' experience of learning and gaining personal knowledge during a stay at a mental hospital. *J Psychiatr Ment Health Nurs* 2008; **15**: 365–73.
- 51 Borge L, Fagermoen MS. Patients' core experiences of hospital treatment: wholeness and self-worth in time and space. *J Ment Health* 2008; **17**: 193–205.
- 52 Cleary M, Horsfall J, Jackson D, O'Hara-Aarons M, Hunt GE. Patients' views and experiences of pro re nata medication in acute mental health settings. *Int J Ment Health Nurs* 2012; **21**: 533–9.
- 53 Cleary M, Hunt GE, Escott P, Walter G. Receiving difficult news. Views of patients in an inpatient setting. *J Psychosoc Nurs Ment Health Serv* 2010; **48**: 40–8.
- 54 Donald F, Duff C, Lee S, Kroschel J, Kulkarni J. Consumer perspectives on the therapeutic value of a psychiatric environment. *J Ment Health* 2015; **24**: 63–7.
- 55 Gunasekara I, Pentland T, Rodgers T, Patterson S. What makes an excellent mental health nurse? A pragmatic inquiry initiated and conducted by people with lived experience of service use. *Int J Ment Health Nurs* 2014; **23**: 101–9.
- 56 Lucas M, Stevenson D. Violence and abuse in psychiatric in-patient institutions: a South African perspective. *Int J Law Psychiatry* 2006; **29**: 195–203.
- 57 Ntsaba GM, Havenga Y. Psychiatric in-patients' experience of being secluded in a specific hospital in Lesotho. *Health SA Gesondheid* 2007; **12**: 3–12.
- 58 Robins CS, Sauvageot JA, Cusack KJ, Suffoletta-Maierle S, Frueh BC. Consumers' perceptions of negative experiences and "sanctuary harm" in psychiatric settings. *Psychiatr Serv* 2005; **56**: 1134–8.
- 59 Roe D, Ronen Y. Hospitalization as experienced by the psychiatric patient: a therapeutic jurisprudence perspective. *Int J Law Psychiatry* 2003; **26**: 317–32.
- 60 Sibitz I, Scheutz A, Lakeman R, Schrank B, Schaffer M, Amering M. Impact of coercive measures on life stories: qualitative study. *Br J Psychiatry* 2011; **199**: 239–44.
- 61 Thibeault CA, Trudeau K, d'Entremont M, Brown T. Understanding the milieu experiences of patients on an acute inpatient psychiatric unit. *Arch Psychiatr Nurs* 2010; **24**: 216–26.
- 62 Thomas SP, Shattell M, Martin T. What's therapeutic about the therapeutic milieu? *Arch Psychiatr Nurs* 2002; **16**: 99–107.
- 63 Johansson H, Eklund M. Patients' opinion on what constitutes good psychiatric care. *Scand J Caring Sci* 2003; **17**: 339–46.
- 64 Brunero S, Lamont S, Fairbrother G. Using and understanding consumer satisfaction to effect an improvement in mental health service delivery. *J Psychiatr Ment Health Nurs* 2009; **16**: 272–8.
- 65 Cleary M, Horsfall J, Hunt GE. Consumer feedback on nursing care and discharge planning. *J Adv Nurs* 2003; **42**: 269–77.
- 66 Eytan A, Bovet L, Gex-Fabry M, Alberque C, Ferrero F. Patients' satisfaction with hospitalization in a mixed psychiatric and somatic care unit. *Eur Psychiatry* 2004; **19**: 499–501.
- 67 Georgieva I, Mulder CL, Wierdsma A. Patients' preference and experiences of forced medication and seclusion. *Psychiatr Q* 2012; **83**: 1–13.
- 68 Greenwood N, Hussain F, Burns T, Raphael F. Asian in-patient and carer views of mental health care. Asian views of mental health care. *J MentHealth* 2000; **9**: 397–408.
- 69 McGuinness D, Dowling M, Trimble T. Experiences of involuntary admission in an approved mental health centre. *J Psychiatr Ment Health Nurs* 2013; **20**: 726–34.
- 70 Olusina AK, Ohaeri JU, Olatawura MO. Patient and staff satisfaction with the quality of in-patient psychiatric care in a Nigerian general hospital. *Soc Psychiatry Psychiatr Epidemiol* 2002; **37**: 283–8.
- 71 Smith D, Roche E, O'Loughlin K, Brennan D, Madigan K, Lyne J, et al. Satisfaction with services following voluntary and involuntary admission. *J Ment Health* 2014; **23**: 38–45.
- 72 Sorgaard KW. Satisfaction and coercion among voluntary, persuaded/persuaded and committed patients in acute psychiatric treatment. *Scand J Caring Sci* 2007; **21**: 214–9.
- 73 Strauss JL, Zervakis JB, Stechuchak KM, Olsen MK, Swanson J, Swartz MS, et al. Adverse impact of coercive treatments on psychiatric inpatients' satisfaction with care. *Community Ment Health J* 2013; **49**: 457–65.
- 74 Anders RL, Olson T, Bader J. Assessment of acutely mentally ill patients' satisfaction of care: there is a difference among ethnic groups. *Issues Ment Health Nurs* 2007; **28**: 297–308.
- 75 Steinert T, Birk M, Flammer E, Bergk J. Subjective distress after seclusion or mechanical restraint: one-year follow-up of a randomized controlled study. *Psychiatr Serv* 2013; **64**: 1012–7.
- 76 Bramesfeld A, Wedegartner F, Elgeti H, Bisson S. How does mental health care perform in respect to service users' expectations? Evaluating inpatient and outpatient care in Germany with the WHO responsiveness concept. *BMC Health Serv Res* 2007; **7**: 99.
- 77 Lilja L, Hellzen O. Former patients' experience of psychiatric care: a qualitative investigation. *Int J Ment Health Nurs* 2008; **17**: 279–86.
- 78 Looi G-ME, Engström Å, Sävenstedt S. A self-destructive care: self-reports of people who experienced coercive measures and their suggestions for alternatives. *Issues Ment Health Nurs* 2015; **36**: 96–103.
- 79 Mayers P, Keet N, Winkler G, Flisher AJ. Mental health service users' perceptions and experiences of sedation, seclusion and restraint. *Int J Soc Psychiatry* 2010; **56**: 60–73.
- 80 Muir-Cochrane E, Oster C, Grotto J, Gerace A, Jones J. The inpatient psychiatric unit as both a safe and unsafe place: implications for absconding. *Int J Ment Health Nurs* 2013; **22**: 304–12.
- 81 Kulkarni J, Gavrilidis E, Lee S, Van Rheenen TE, Grigg J, Hayes E, et al. Establishing female-only areas in psychiatry wards to improve safety and quality of care for women. *Australas* 2014; **22**: 551–6.

- 82 Lindgren B-M, Aminoff C, Graneheim UH. Features of everyday life in psychiatric inpatient care for self-harming: an observational study of six women. *Issues Ment Health Nurs* 2015; **36**: 82–8.
- 83 Shattell M, Melanie Andes M, Thomas S. How patients and nurses experience the acute care psychiatric environment. *Nurs Inq* 2008; **15**: 242–50.
- 84 Giacco D, Fiorillo A, Del Vecchio V, Kallert T, Onchev G, Raboch J, et al. Caregivers' appraisals of patients' involuntary hospital treatment: European multicentre study. *Br J Psychiatry* 2012; **201**: 486–91.
- 85 Svindseth MF, Dahl AA, Hatling T. Patients' experience of humiliation in the admission process to acute psychiatric wards. *Nord J Psychiatry* 2007; **61**: 47–53.
- 86 Thapinta D, Anders RL, Wiwatkunupakan S, Kitsumban V, Vadtanapong S. Assessment of patient satisfaction of mentally ill patients hospitalized in Thailand. *Nurs Health Sci* 2004; **6**: 271–7.
- 87 Chien WT, Chan CW, Lam LW, Kam CW. Psychiatric inpatients' perceptions of positive and negative aspects of physical restraint. *Patient Educ Couns* 2005; **59**: 80–6.
- 88 Ezeobele IE, Malecha AT, Mock A, Mackey-Godine A, Hughes M. Patients' lived seclusion experience in acute psychiatric hospital in the United States: a qualitative study. *J Psychiatr Ment Health Nurs* 2014; **21**: 303–12.
- 89 Holmes D, Kennedy SL, Perron A. The mentally ill and social exclusion: a critical examination of the use of seclusion from the patient's perspective. *Issues Ment Health Nurs* 2004; **25**: 559–78.
- 90 Iversen VC, Sallaup T, Vaaler AE, Helvik A-S, Morken G, Linaker O. Patients' perceptions of their stay in a psychiatric seclusion area. *J Psychiatr Intensive Care* 2011; **7**: 1–10.
- 91 Meehan T, Vermeer C, Windsor C. Patients' perceptions of seclusion: a qualitative investigation. *J Adv Nurs* 2000; **31**: 370–7.
- 92 O'Brien L, Cole R. Mental health nursing practice in acute psychiatric close-observation areas. *Int J Ment Health Nurs* 2004; **13**: 89–99.
- 93 Binnema D. Interrelations of psychiatric patient experiences of boredom and mental health. *Issues Ment Health Nurs* 2004; **25**: 833–42.
- 94 Cutcliffe J, Santos J, Kozel B, Taylor P, Lees D. Raiders of the Lost Art: a review of published evaluations of inpatient mental health care experiences emanating from the United Kingdom, Portugal, Canada, Switzerland, Germany and Australia. *Int J Ment Health Nurs* 2015; **24**: 375–85.
- 95 Duncan E, Best C, Hagen S. Shared decision making interventions for people with mental health conditions. *Cochrane Database Syst Rev* 2010: N.PAG-N. PAG 1p.
- 96 Ford S-B, Bowyer T, Morgan P. The experience of compulsory treatment: the implications for recovery-orientated practice? *Ment Health Soc Inclusion* 2015; **19**: 126–32.
- 97 Gerolamo AM. State of the science: outcomes of acute inpatient psychiatric care. *Arch Psychiatr Nurs* 2004; **18**: 203–14.
- 98 Hopkins JE, Loeb SJ, Fick DM. Beyond satisfaction, what service users expect of inpatient mental health care: a literature review. *J Psychiatr Ment Health Nurs* 2009; **16**: 927–37.
- 99 Katsakou C, Priebe S. Patient's experiences of involuntary hospital admission and treatment: a review of qualitative studies. *Epidemiol Psychiatr Soc* 2007; **16**: 172–8.
- 100 McHale J, Felton A. Self-harm: what's the problem? A literature review of the factors affecting attitudes towards self-harm. *J Psychiatr Ment Health Nurs* 2010; **17**: 732–40.
- 101 Maatta S. Exploring male and female patients' experiences of psychiatric hospital care: A critical analysis of the literature. *Issues Ment Health Nurs* 2009; **30**: 174–80.
- 102 Newman D, O'Reilly P, Lee SH, Kennedy C. Mental health service users' experiences of mental health care: an integrative literature review. *J Psychiatr Ment Health Nurs* 2015; **22**: 171–82.
- 103 Omer S, Priebe S. Continuity across inpatient and outpatient mental health care or specialisation of teams? A systematic review. *Eur Psychiatry* 2015; **30**: 258–70.
- 104 Sequeira H, Halstead S. Restraint and seclusion: service user views. *J Adult Protection* 2002; **4**: 15–24.
- 105 Strout TD. Perspectives on the experience of being physically restrained: an integrative review of the qualitative literature. *Int J Ment Health Nurs* 2010; **19**: 416–27.
- 106 Sturrock A. Restraint in inpatient areas: the experiences of service users. *Ment Health Pract* 2010; **14**: 22–6.
- 107 Van der Merwe M, Bowers L, Jones J, Simpson A, Haglund K. Locked doors in acute inpatient psychiatry: a literature review. *J Psychiatr Ment Health Nurs* 2009; **16**: 293–9.
- 108 Van Der Merwe M, Muir-Cochrane E, Jones J, Ziggili M, Bowers L. Improving seclusion practice: implications of a review of staff and patient views. *J Psychiatr Ment Health Nurs* 2013; **20**: 203–15.



## 100 words

### 100 words...on psychobiotics

Timothy G. Dinan

That gut microbes, collectively called the microbiota, influence brain development and functioning is viewed as a new paradigm in neuroscience with implications for psychiatry. These gut microbes communicate with the brain via a number of routes including the vagus nerve and the production of molecules such as short-chain fatty acids. In major depressive disorder the gut microbiota shows a significant decrease in microbial diversity which is associated with a peripheral inflammatory phenotype. Psychobiotics are bacteria which, when ingested in appropriate amounts, have positive mental health benefits. Preliminary studies with bacteria such as *Bifidobacterium longum* indicate anxiolytic activity in healthy volunteers.

Copyright © The Royal College of Psychiatrists 2019

The British Journal of Psychiatry (2019)  
214, 338. doi: 10.1192/bjp.2019.81