

Seedbank Depletion and Emergence Patterns of Giant Ragweed (*Ambrosia trifida*) in Minnesota Cropping Systems

Jared J. Goplen, Craig C. Sheaffer, Roger L. Becker, Jeffrey A. Coulter, Fritz R. Breitenbach, Lisa M. Behnken, Gregg A. Johnson, and Jeffrey L. Gunsolus*

In the midwestern United States, biotypes of giant ragweed resistant to multiple herbicide biochemical sites of action have been identified. Weeds with resistance to multiple herbicides reduce the utility of existing herbicides and necessitate the development of alternative weed control strategies. In two experiments in southeastern Minnesota, we determined the effect of six 3 yr crop-rotation systems containing corn, soybean, wheat, and alfalfa on giant ragweed seedbank depletion and emergence patterns. The six crop-rotation systems included continuous corn, soybean–corn–corn, corn–soybean–corn, soybean–wheat–corn, soybean–alfalfa–corn, and alfalfa–alfalfa–corn. The crop-rotation system had no effect on the amount of seedbank depletion when a zero-weed threshold was maintained, with an average of 96% of the giant ragweed seedbank being depleted within 2 yr. Seedbank depletion occurred primarily through seedling emergence in all crop-rotation systems. However, seedling emergence tended to account for more of the seedbank depletion in rotations containing only corn or soybean compared with rotations with wheat or alfalfa. Giant ragweed emerged early across all treatments, with on average 90% emergence occurring by June 4. Duration of emergence was slightly longer in established alfalfa compared with other cropping systems. These results indicate that corn and soybean rotations are more conducive to giant ragweed emergence than rotations including wheat and alfalfa, and that adopting a zero-weed threshold is a viable approach to depleting the weed seedbank in all crop-rotation systems.

Nomenclature: Giant ragweed, *Ambrosia trifida* L. AMBTR, alfalfa, *Medicago sativa* L., corn, *Zea mays* L., soybean, *Glycine max* (L.) Merr., wheat, *Triticum aestivum* L.

Key words: Crop rotation, herbicide resistance, weed emergence, weed seedbank depletion.

Giant ragweed is one of the most competitive agricultural weeds plaguing crops in the midwestern United States (Webster et al. 1994). A single giant ragweed plant m^{-2} has the potential to reduce soybean yields by 45 to 77%, and 1.4 giant ragweed plants m^{-2} can reduce corn yields by up to 90% (Harrison et al. 2001; Webster et al. 1994). As an annual, giant ragweed relies on the seedbank to persist in agricultural fields, and seed production in soybean and field margins is nearly 2,000 seeds $plant^{-1}$, indicating high potential for seedbank replenishment (Fenner 1995; Goplen et al. 2016).

Giant ragweed control in corn and soybean rotations has become complicated due to the development of

resistance to multiple herbicide mechanisms of action, including both acetolactate synthase (ALS) inhibitors and glyphosate (Heap 2015). The availability of multiple glyphosate-resistant crops beginning in 1996 allowed producers to use glyphosate as an effective postemergence, broad-spectrum herbicide with low cost, which led to glyphosate being used as a stand-alone herbicide on millions of hectares of cropland (Duke and Powles 2008). The widespread and repeated use of glyphosate, paired with application to large weeds, caused tremendous selection pressure on weed populations and resulted in glyphosate-resistant weeds. When paired with ALS-inhibitor resistance, which had previously developed in giant ragweed, resistance to multiple herbicide sites of action was stacked within the same plant. As herbicide-resistant giant ragweed becomes more prevalent, there is an increased need for integrative forms of weed control (Shaner and Beckie 2014).

Crop rotations have long been recognized as an effective means to control a diversity of weeds (Leighty 1938; Liebman and Dyck 1993). Diverse crop rotations provide the opportunity to use

DOI: 10.1614/WS-D-16-00084.1

First, second, third, fourth, seventh, and eighth authors: Graduate Student, Professor, Professor, Associate Professor, Associate Professor, and Professor, Department of Agronomy and Plant Genetics, University of Minnesota, 1991 Upper Buford Circle, Saint Paul, MN 55108; fifth and sixth authors: Extension Educator and Integrated Pest Management Specialist, University of Minnesota, 863 30th Ave. SE, Rochester, MN 55904. Corresponding author's E-mail: gople007@umn.edu

different weed control strategies, including a variety of mechanical and chemical methods. Planting crops that permit the use of alternative herbicide sites of action makes control of resistant weeds more manageable and reduces the potential of weeds developing additional resistance (Norsworthy et al. 2012). More diverse crop rotations also provide a more suitable habitat for seed predators that reduce weed seedbanks (Davis and Liebman 2003; Westerman et al. 2005). Overall, crop sequences that vary in patterns of resource competition, soil disturbance, and mechanical damage create unstable environments hostile to any particular weed species, ultimately decreasing weed populations (Liebman and Dyck 1993; Schreiber 1992). Using crop rotations that diversify herbicide sites of action, in addition to promoting weed seedbank depletion via seed decay and predation, offers potential to effectively manage herbicide-resistant giant ragweed over the long term (Chee-Sanford et al. 2006).

Seed predation by rodents and invertebrates has been shown to remove nearly 90% of giant ragweed seeds within one year in no-tillage corn (Harrison et al. 2003). Seed predation increases in wheat and alfalfa compared with annual row crops due to greater crop canopy (Hartzler et al. 2007; Westerman et al. 2005). In addition, alfalfa is harvested several times throughout the growing season via mowing, eliminating giant ragweed seed production and replenishment of the weed seedbank. Wheat increases early-season competition with emerging giant ragweed by being planted earlier in the growing season than corn or soybean and in narrower rows. Additionally, wheat allows the use of herbicides with alternative mechanisms of action that are more effective against herbicide-resistant populations of giant ragweed. In the event of weed escapes, wheat is harvested prior to giant ragweed seed production (Goplen et al. 2016), preventing replenishment of the seedbank and allowing time for multiple mechanical and chemical weed control options following wheat harvest.

Although crop rotation is known to be an effective form of weed control, the effects of crop-rotation systems on giant ragweed seedbank dynamics have not been thoroughly evaluated. The objectives of this research were to determine how cropping systems common to the midwestern United States affect giant ragweed's (1) timing of emergence, (2) total emergence, and (3) seedbank depletion after two growing seasons. This research will help define which crop-rotation systems promote soil conditions most conducive to minimizing giant ragweed emergence and

maximizing seedbank depletion, allowing crop producers to determine the most effective ways to proactively manage herbicide-resistant giant ragweed infestations.

Materials and Methods

Site Details. Two replicated experiments were initiated in 2012 and 2013 at different sites near Rochester, MN (43.91°N, 92.56°W). Both sites were on a Port Byron silt loam (fine-silty, mixed, superactive, mesic Typic Hapludolls) with a pH of 7.0 and 4.0% organic matter with a history of a 2-yr corn–soybean rotation, with soybean immediately preceding both experiments. The research sites had known populations of giant ragweed resistant to glyphosate and ALS-inhibitor herbicides.

Crop Management. Each experiment had six crop rotations arranged in a randomized complete block design with four replications. Crop-rotation treatments consisted of: continuous corn (CCC); soybean–corn–corn (SCC); corn–soybean–corn (CSC); soybean–wheat–corn (SWC); soybean–alfalfa–corn (SAC); and alfalfa–alfalfa–corn (AAC). Plots were 10 by 15 m. Corn and alfalfa cultivars had resistance to glyphosate and corn and soybean cultivars were glufosinate-resistant. DEKALB[®] DKC53-78RIB corn was planted at 86,500 seeds ha⁻¹ in 76 cm rows. Soybean plots were planted with Stine[®] 19LD08 at 345,900 seeds ha⁻¹ in 76 cm rows. Inoculated alfalfa (DEKALB[®] DKA41-18RR) was direct seeded with a no-till drill at 16.8 kg ha⁻¹ in 19 cm rows. MN RB07 wheat was direct seeded with a no-till drill at 135 kg ha⁻¹ in 19 cm rows. Fertilizer was applied according to University of Minnesota guidelines (Kaiser et al. 2011). Phosphorus, potassium, and sulfur were broadcast using mono-ammonium phosphate, potash, and ammonium sulfate, respectively, across the entire study area in the fall of each year to maintain adequate levels of these nutrients for all crops grown. At planting, ammonium nitrate was applied at 191, 135, 0, and 129 kg N ha⁻¹ for corn following corn or wheat, corn following 1 yr of alfalfa, corn following 2 yr of alfalfa, and wheat following soybean, respectively.

Corn plots were chisel plowed to a depth of 20 cm in the fall after corn harvest and stover chopping, and were field cultivated twice in the spring just prior to planting. Soybean plots were field cultivated twice in the spring just prior to planting. Soybean stubble following harvest was chisel plowed to a depth of 20 cm in the fall when corn was to be planted the following year and was left fallow when wheat or alfalfa was to be seeded the following year.

Wheat was no-tilled into standing soybean stubble, and wheat stubble was chisel plowed to a depth of 20 cm in the fall after harvest at the same time as chisel plowing occurred in the other crop-rotation systems. Alfalfa plots that were seeded in the first year of the rotation received a single pass with a field cultivator prior to planting, while alfalfa plots seeded in the second year of the rotation were no-till seeded into soybean stubble.

Fields were scouted for insect pests and diseases, and none reached levels warranting treatment according to University of Minnesota guidelines (University of Minnesota Extension 2015). Wheat plots were sprayed prophylactically with tebuconazole (alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol) (Bayer CropScience® Folicur 3.6F) at 126 g ai ha⁻¹ at flower initiation (Zadoks 61) to prevent the development of *Fusarium* head blight (*Fusarium graminearum* L.) (Simmons et al. 1995).

A zero-weed threshold was maintained throughout the study to determine giant ragweed seedbank depletion. Due to the presence of glyphosate-resistant and ALS inhibitor-resistant giant ragweed populations, herbicides specifically targeting resistant giant ragweed were used. When herbicides with residual activity on giant ragweed were used, quadrats where emergence was monitored were covered at time of application to prevent herbicide coverage. Quadrats were not covered for POST applications of herbicides without residual activity, as seedlings were counted and removed prior to herbicide application. Corn and soybean plots received a single PRE application of S-metolachlor at 2.14 kg ai ha⁻¹ on the date of planting each year. Corn and soybean plots received two POST applications of glufosinate-ammonium at 450 g ai ha⁻¹ at approximately 3 and 6 wk post-planting. Alfalfa plots received a single application of 2,4-DB, dimethylamine salt at 1.12 kg ae ha⁻¹ 2 wk following planting in the seeding year, while second-year alfalfa received no herbicide application. Wheat plots received a tank-mixed POST application of a prepackaged mixture of clopyralid monoethanolamine salt and fluroxypyr, 1-methylheptyl ester at 105 g ae ha⁻¹ each (Widematch) and MCPA isooctyl (2-ethylhexyl) ester at 389 g ae ha⁻¹ approximately 2 wk post-planting, when wheat was beyond the 3-leaf stage (Zadoks 13) (Simmons et al. 1995). Weeds escaping herbicide control were hand weeded to ensure there were no inputs into the seedbank.

Seedbank Monitoring. Giant ragweed seedbank densities were determined after emergence terminated in mid-July in the initial and final year of each crop-

rotation system. Seedbank samples in the first year at the first experimental location were taken from three fixed quadrat locations in each plot by sampling a single 25 by 40 cm area to a depth of 15 cm. Due to the amount of time necessary to extract each sample, an alternative sampling method was used to determine seedbank densities in the final year at the first experimental location and in both the initial and final years at the second experimental location. The alternative sampling procedure, adapted from Forcella (1992), used a 10-cm-diameter golf-hole cutter to obtain a composite of 10 samples to a 15 cm depth collected from the same three quadrat locations in each plot. Weed seeds were separated from combined samples using a modified version of a physical extraction procedure adapted from Ball and Miller (1989), Cardina and Sparrow (1996), and Standifer (1980), in which compiled samples were wet sieved to separate seeds. Samples were soaked with water and mixed several times over 20 min using a paint-stirring attachment on an electric drill. Once the soil was in suspension, samples were poured through a 0.16 cm sieve to extract seeds. Remaining soil was soaked again and mixed until the entire sample passed through the sieve. A low-pressure shower of water was sprayed on the sample to speed the passing of soil through the sieve. Once organic material >0.16 cm was separated from soil, the seedbank extract was placed in a 60 C forced-air oven for 2 d to dry the sample before seeds and seed fragments were handpicked from the samples. Seeds were then determined to be potentially viable or nonviable by dissecting the seeds to determine the presence of an intact embryo and counted, similar to methods defined previously (Ball and Miller 1989; Cardina and Sparrow 1996; Standifer 1980).

Emergence Monitoring. Giant ragweed emergence counts were made on a weekly basis starting at the onset of emergence and continued for at least 10 wk or until emergence ceased each year. Giant ragweed emergence was monitored in six permanent 30 by 76 cm quadrats within each plot. Three quadrats were placed between rows, and three quadrats were placed over the crop row. Each week, seedlings were counted and removed from the quadrat by clipping seedlings at the soil surface without disturbing the soil.

Environmental Effects. Daily precipitation, soil temperature, and growing degree days (GDDs) were monitored to determine their effects on giant ragweed emergence. Daily precipitation and minimum and maximum air temperatures were obtained from the National Weather Service station nearest the study

locations. Soil temperature was monitored at a 5 cm depth using temperature sensors (Hobo Water Temp Pro v. 2), logging temperature at hourly intervals (Figure 1). GDDs were calculated using Equation 1,

$$\text{GDD} = \sum_{S_2}^{S_1} \frac{(T_{\max} + T_{\min})}{2} - b_0 \quad [1]$$

where T_{\max} is the maximum daily air temperature, T_{\min} is the minimum daily air temperature, b_0 is the base temperature (2 C), and S_1 and S_2 are April 1 and July 31, respectively (Table 1) (Abul-Fatih and Bazzaz 1979).

Statistical Analysis. Seedbank depletion, total giant ragweed emergence each year, and the proportion of the seedbank depleted due to emergence were each




Figure 1. Mean daily soil temperature throughout the giant ragweed emergence period at 5 cm depth in the second year of each crop-rotation system in 2013 and 2014.

^aC, S, W, and A represent the sequence of corn, soybean, wheat, and alfalfa in each 3 yr crop-rotation system.

analyzed using the MIXED procedure of SAS, with crop-rotation system considered a fixed effect and experimental location, block (nested within location), interactions, and subsampling considered random effects (SAS Institute 2012). Data from both experimental locations were combined following analysis of interactions. Mean comparisons were made for each set of analyses using Fisher's protected LSD test ($\alpha = 0.05$) and transformed means (described below) were back transformed for presentation.

Seedbank depletion was calculated as a percentage of the initial giant ragweed seedbank density. Since seedbank depletion between the first and third years of each crop-rotation system was a percentage and exhibited a skewed distribution, data were arcsine transformed prior to analysis. The total giant ragweed emergence in each year exhibited a nonnormal distribution and was transformed with the natural log transformation prior to analysis. Additionally, the initial seedbank density was included as a random effect covariate, to control for spatial variation in the analysis of total giant ragweed emergence each year. Since the initial seedbank samples were taken after giant ragweed emergence, the proportion of the seedbank depleted by emergence was calculated as the total giant ragweed emergence from the second and third year of the crop-rotation systems, therefore representing the emergence that occurred between the initial and final seedbank samples. The proportion of seedbank depletion due to emergence exhibited a skewed distribution and was transformed with the natural log transformation prior to analysis.

To evaluate emergence timing of giant ragweed, weekly emergence counts were converted to a cumulative percentage of the total seedlings that emerged

Table 1. Mean air temperature, total growing degree days (GDDs), and total precipitation by month and across the growing season (April to October) during the study period from the nearest National Weather Service station.^a

Month	Air temperature (C)				GDD ^b				Total precipitation (mm)			
	2012	2013	2014	2015	2012	2013	2014	2015	2012	2013	2014	2015
April	9.5	4.6	5.7	8.7	225	100	126	203	68	173	144	125
May	17.4	13.2	14.4	14.3	476	350	384	380	123	311	45	133
June	21.7	19.4	20.4	19.6	590	523	553	529	83	174	185	113
July	25.0	21.9	19.8	20.8	714	616	550	584	83	54	57	126
August	20.7	21.0	21.4	19.2	578	590	601	533	50	46	136	80
September	16.6	17.8	15.4	19.0	437	474	403	510	33	31	90	73
October	9.3	8.7	8.6	10.5	225	214	205	246	47	76	60	24
Mean	17.2	15.2	15.1	16.0	—	—	—	—	—	—	—	—
Cumulative	—	—	—	—	3,246	2,867	2,821	2,984	485	865	717	675

^a Weather data obtained from site identification: KRST (43.9041°N, -92.4916°W).

^b GDDs calculated using Equation 1 (2 C base temperature).

each year. Cumulative percent emergence of giant ragweed in each crop-rotation system was modeled using a logistic regression equation as a function of day of year (doy) (Equation 2). The emergence data in both experimental locations were similar and were combined for analysis. The emergence rate (ER) and inflection point (IP) of Equation 2

$$\% \text{ emerged} = \frac{100}{1 + e^{-ER(\text{doy} - \text{IP})}} \quad [2]$$

were estimated for the second year of each crop-rotation system and were used to predict the dates when 50 and 90% emergence occurred in each crop-rotation system. Only emergence in the second year of each crop-rotation system was modeled, because all four crops were present in the second year, illustrating the greatest cropping diversity effects on giant ragweed emergence.

Results and Discussion

Total Emergence. Across both experimental locations, 125, 37, and 6 seedlings $\text{m}^{-2} \text{yr}^{-1}$ emerged in years 1, 2, and 3 of the crop-rotation systems, respectively, representing a major threat for crop yield and control costs. Initial seedbank densities over the study locations ranged from 0 to 317 seeds m^{-2} . Due to spatial variation of the initial weed seedbank at both locations, estimates of the starting seedbank density were included as a covariate in analysis of total emergence in each year of each cropping system ($R^2 = 0.31$, $P < 0.001$). In the first year, there were no differences in giant ragweed emergence among crop-rotation systems ($P = 0.73$; Table 2), which was expected, since all crops were planted into a site with the same management history. Additionally, crop planting was delayed in the first year of the cropping systems at the second experimental location in 2013 due to large amounts of precipitation (Table 1), resulting in a large percentage of giant ragweed emergence prior to crop planting, thus minimizing any cropping effects.

There were differences in total giant ragweed emergence in the second year of the crop-rotation systems, in that second-year alfalfa had the least emergence of all crop-rotation systems. Although not significantly different from the continuous corn treatment, corn following soybean had the greatest amount of giant ragweed emergence in the second year of the crop-rotation systems (Table 2). The other crops planted following soybean, including wheat and alfalfa in the SWC and SAC treatments, had less emergence of giant ragweed than SCC in the second year of the

Table 2. Total giant ragweed emergence in each year, percentage of seedbank depletion, and the percentage of depletion accounted for by emergence in each crop-rotation system, across both experimental locations in 2012 to 2015.^a

Crop-rotation system ^d	Emergence ^b			Seedbank ^c	
	Year 1	Year 2	Year 3	Depletion	Emergence
	— Seedlings m^{-2} —			— % —	
CCC	3.4 ns	5.6 ab	1.1 ab	97.7 ns	100 ab
SCC	7.0	10.5 a	0.1 c	95.7	100 a
CSC	4.7	4.8 b	1.5 a	90.6	96 bc
SWC	6.5	4.1 b	0.5 abc	94.7	61 bc
SAC	7.1	4.6 b	0.5 bc	98.4	74 bc
AAC	4.4	0.8 c	0.3 bc	99.0	41 c

^a Means with different letters indicate a significant difference at the 0.05 probability level using Fisher's protected LSD.

^b Total seedling emergence in each year is the corrected mean from the seedbank density covariate.

^c Seedbank depletion represents the percentage of seeds depleted between the first and third year of the crop-rotation systems, while emergence represents the percentage of the seedbank depletion that was accounted for by emergence during the same time period.

^d C, S, W, and A represent the sequence of corn, soybean, wheat, and alfalfa in each 3 yr crop-rotation system.

rotations, despite similar overwinter and early-spring conditions. However, the pre-planting tillage and planting date differed among the SWC, SAC, and SCC systems in the second year of the rotations, with alfalfa and wheat planted an average of 22 d prior to corn and soybean. There were differences in total emergence in the third year of the crop-rotation systems, despite having consistent tillage type and timing the preceding fall and spring prior to planting (Table 2). Overall, differences in emergence could have been due to variations in soil temperature (Figure 1) and crop residues, since even slight changes in planting date, cultivation timing, and residue management can influence vertical seed movement, seed dormancy dynamics, and overall emergence of giant ragweed (Buhler 1995; Buhler et al. 1997; Cousens and Moss 1990; Dyer 1995; Staricka et al. 1990).

The second year of the AAC system had the least giant ragweed emergence among crop-rotation systems, which was expected, since giant ragweed is least adapted to the perennial environment of alfalfa. In the fall of the first year of the AAC system, there was significant alfalfa canopy coverage, likely buffering the soil environment from extreme temperature changes throughout the winter. In the spring of the second year, this canopy coverage likely contributed to reduced fluctuations in soil temperature and cooler soil temperature in the established alfalfa than in the exposed soil of the other treatments, which

was most extreme from May 20 to June 12 in 2014 (Figure 1). These differences in soil temperature were minimized after June 12 in 2014, as alfalfa was harvested on this date. Additionally, less tillage occurred in the AAC system. Previous work has shown that weed emergence depth is shallower with no-tillage than chisel and moldboard plowing (Buhler and Mester 1991). Seeds of large-seeded weeds like giant ragweed tend to remain near the soil surface with less intensive tillage, which has been shown to inhibit establishment of these species (Lueschen and Anderson 1980). Therefore, lower soil temperatures along with less intensive tillage likely provided a soil environment less conducive to giant ragweed emergence. This environment may also have been more conducive to seedbank degradation through seedling mortality, seed degradation, and seed predation, resulting in less emergence with similar levels of seedbank depletion (Table 2).

Emergence Timing. Giant ragweed exhibited similar emergence patterns in each year of the crop-rotation system treatments, following a logistic growth curve relative to date (Figure 2; Table 3). Only emergence from the second year of the crop-rotation system is presented here, because corn, soybean, wheat, and alfalfa were all planted in the second year and represent the greatest cropping diversity in a single year. Giant ragweed began emerging slowly in the early weeks of each growing season before having a period of rapid emergence throughout May. Emergence tapered

off and nearly terminated mid-June. Giant ragweed emergence initiated at a similar date in 2013 and 2014, but emergence occurred slightly more rapidly in 2013, which was likely caused by greater precipitation from April to June in 2013 (Table 1). Despite slight yearly differences, across all treatments, 50 and 90% of giant ragweed emergence in all years occurred on May 21 and June 4, respectively (Table 3), indicative of the early emergence pattern of giant ragweed in much of the midwestern United States (Werle et al. 2014). Giant ragweed in this study did not exhibit a biphasic emergence pattern, as has been identified in Ohio (Schutte et al. 2012). All crop-rotation systems exhibited a similar pattern of cumulative emergence, except for the second year alfalfa in the AAC system. The second year of the AAC system achieved 50% emergence at the same time as the other crop-rotation systems, indicated by the similarity in inflection points of the crop-rotation systems (Table 3). However, the AAC system had a longer duration of emergence, reflected by the slower emergence rate, and did not reach 90% emergence until June 18, several weeks after the other crop-rotation systems (Table 3; Figure 2). Giant ragweed emergence has been shown to be associated with the accumulation of thermal time (Archer et al. 2006; Davis et al. 2013; Schutte et al. 2008; Werle et al. 2014). The prolonged pattern of emergence of giant ragweed in the second year of the AAC system may be related to the greater early-season crop canopy of alfalfa, which reduced the amount of solar radiation reaching the soil surface, thereby decreasing soil temperature and slowing the accumulation of thermal time at the soil level (Figure 1).




Figure 2. Cumulative percentage of giant ragweed emergence in the second year of each crop-rotation system at both experimental locations based on the best-fit logistic regression equations (Table 3).

^aC, S, W, and A represent the sequence of corn, soybean, wheat, and alfalfa in each 3 yr crop-rotation system.

Seedbank Depletion. There were no differences in giant ragweed seedbank depletion among crop-rotation systems when a zero-weed threshold was maintained ($P = 0.57$). On average, 96% of the giant ragweed seedbank was depleted in 2 yr in any crop-rotation system (Table 2), indicating that the giant ragweed seedbank is short-lived regardless of the cropping system. Harrison et al. (2007) found that depletion of the giant ragweed seedbank was dependent on burial depth and that seeds closer to the soil surface were degraded more quickly than seeds >10 cm deep. A small percentage of giant ragweed seed remaining in the seedbank has been shown to persist for up to 15 yr, which exemplifies the importance of long-term weed management (Hartnett et al. 1987; Loux and Berry 1991).

There are multiple ways weed seeds can be depleted from the seedbank. Weed seeds can germinate and

Table 3. Parameter estimates and predicted dates of 50 and 90% cumulative giant ragweed emergence in the second year of each crop-rotation system, across both experimental locations.

Crop-rotation system ^a	Percentage emerged ^b		Parameter estimates ^c			
	50%	90%	ER	SE _{ER}	IP	SE _{IP}
	Date					
CCC	May 22 ns	June 3 a	0.19	0.02	142	0.7
SCC	May 22	June 2 a	0.20	0.02	142	0.7
CSC	May 21	May 31 a	0.22	0.03	141	0.7
SWC	May 22	June 5 a	0.15	0.02	142	0.8
SAC	May 21	June 5 a	0.15	0.02	141	0.8
AAC	May 21	June 18 b	0.08	0.01	140	1.3

^a C, S, W, and A represent the sequence of corn, soybean, wheat, and alfalfa in each 3 year crop-rotation system.

^b Dates for percentage emerged are calculated from the logistic regression parameter estimates. Dates with different letters indicate significant difference.

^c Parameter estimates are from the best-fit logistic regressions (Equation 2) from each crop-rotation system based on cumulative giant ragweed emergence. Abbreviations: ER, emergence rate; IP, inflection point; SE_{ER}, standard error for ER parameter estimate; SE_{IP}, standard error for IP parameter estimate.

emerge or die, fungi and other soil microorganisms can decay the seeds, and predators such as birds and rodents can consume seeds (Buhler et al. 1997; Chee-Sanford et al. 2006; Kremer 1993). Each of these mechanisms of seedbank degradation has potential to cause significant seedbank losses. Harrison et al. (2003) found that up to 90% of giant ragweed seeds deposited on the soil surface of a no-tillage cornfield can be eliminated by predation in a single year. Additionally, the rate of seed predation increases as the crop canopy develops within a field, with wheat and alfalfa typically having greater seed predation than corn (Hartzler et al. 2007; Westerman et al. 2005).

A large portion of giant ragweed seedbank depletion was due to emergence in all crop-rotation systems, ranging from 41 to 100% among crop-rotation systems (Table 2). In the CCC, SCC, and CSC systems, nearly 100% of seedbank depletion was due to emergence, although CCC and CSC did not have significantly greater depletion due to emergence than SWC and SAC, which accounted for 61 and 74%, respectively. Less emergence in the second year of the AAC treatment likely resulted in the AAC treatment having less seedbank depletion due to emergence than the CCC and SCC crop-rotation systems (Table 2). These results indicate that there was slightly more depletion of the giant ragweed seedbank due to factors other than emergence in the AAC system, which is supported by Brust and House (1988), who found that depletion of weed seedbanks due to seed predators and soil microorganisms is greater in more diverse cropping systems with more habitat for seed predators.

Conclusions. These results align with previous research indicating that giant ragweed is a relatively early-emerging weed with a short duration of emergence compared with other common weeds in the midwestern United States (Werle et al. 2014). This early-season emergence pattern indicates that there is potential to enhance giant ragweed control through the timing of field operations. Although delayed planting may reduce crop yield potential, it allows a greater percentage of seedlings to emerge and be destroyed prior to planting when tillage and/or herbicides can be used to control early-emerging weeds just prior to planting (Gill 1996; Walsh and Powles 2007). Crops also have a faster rate of growth when planted later in the growing season, providing an additional competitive advantage over weeds (Tsimba et al. 2013). It is also expected that delayed planting will improve control of giant ragweed populations that have a biphasic emergence pattern like those found in Ohio, since these populations still have an early flush of emergence that can be affected by planting date (Schutte et al. 2012). Additionally, delayed planting provides the opportunity for PRE herbicide residual activity to extend weed control later into the growing season.

This study indicates that the giant ragweed seedbank is short-lived and that crop-rotation systems do not differ in the amount of seedbank depletion when a zero-weed threshold is implemented. More specifically, these results indicate that weed seed inputs in the cropping systems studied only need to be prevented for 2 yr to reduce the giant ragweed seedbank by 96%. Implementing a zero-weed threshold may be easier if total annual

emergence is reduced. Corn following soybean had the greatest total emergence of giant ragweed. Conversely, emergence in the AAC system was less than that of the other cropping systems, indicating that the inclusion of alfalfa in the cropping system has the greatest potential to improve giant ragweed control. Although the emergence period was longer in the AAC system, the harvesting schedule of established alfalfa prevents seedbank inputs without reliance on herbicides. Overall, these results indicate that there is potential to manage fields infested with giant ragweed by eliminating seedbank inputs and depleting the weed seedbank to ultimately improve herbicide-resistant giant ragweed control.

Acknowledgments

This research was funded by the Monsanto Graduate Fellowship, the Rapid Agricultural Response Fund of the Minnesota Agricultural Experiment Station, and the Torske Klubben Graduate Fellowship. The authors would also like to express appreciation to staff and students for their assistance, in particular Brad Kincaid and Doug Miller.

Literature Cited

- Abul-Fatih HA, Bazzaz FA (1979) The biology of *Ambrosia trifida* L. II. germination, emergence, growth and survival. *New Phytol* 83:817–827
- Archer DW, Forcella F, Korth A, Kuhn A, Eklund J, Spokas K (2006). *WeedCast*. <http://www.ars.usda.gov/services/software/download.htm?softwareid=112>. Accessed April 23, 2013
- Ball DA, Miller SD (1989) A comparison of techniques for estimation of arable soil seedbanks and their relationship to weed flora. *Weed Res* 29:365–373
- Brust GE, House GJ (1988) Weed seed destruction by arthropods and rodents in low-input soybean agroecosystems. *Am J Alt Agric* 3:19–25
- Buhler DD (1995) Influence of tillage systems on weed population dynamics and management in corn and soybean in the central USA. *Crop Sci* 35:1247–1258
- Buhler DD, Mester TC (1991) Effect of tillage systems on the emergence depth of giant (*Setaria faberi*) and green foxtail (*Setaria viridis*). *Weed Sci* 39:200–203
- Buhler DD, Hartzler RG, Forcella F (1997) Implications of weed seedbank dynamics to weed management. *Weed Sci* 45:329–336
- Cardina J, Sparrow DH (1996) A comparison of methods to predict weed seedling populations from the soil seedbank. *Weed Sci* 44:46–51
- Chee-Sanford JC, Davis AS, Sims GK (2006) Do microorganisms influence seed-bank dynamics? *Weed Sci* 54:575–587
- Cousens R, Moss SR (1990) A model of the effects of cultivation on the vertical distribution of weed seeds within the soil. *Weed Res* 30:61–70
- Davis AS, Liebman M (2003) Cropping system effects on giant foxtail (*Setaria faberi*) demography. 1. Green manure and tillage timing. *Weed Sci* 54:919–929
- Davis AS, Clay S, Cardina J, Dille A, Forcella F, Lindquist J, Sprague C (2013) Seed burial physical environment explains departures from regional hydrothermal model of giant ragweed (*Ambrosia trifida*) seedling emergence in U.S. Midwest. *Weed Sci* 61:415–421
- Duke SO, Powles SB (2008) Glyphosate: a once-in-a-century herbicide. *Pest Manag Sci* 64:319–325
- Dyer WE (1995) Exploiting weed seed dormancy and germination requirements through agronomic practices. *Weed Sci* 43:498–503
- Fenner M (1995) Ecology of seed banks. Pages 507–528 in Kigel J & Galili G, eds. *Seed Development and Germination*. New York: Marcel Dekker
- Forcella F (1992) Prediction of weed seedling densities from buried seed reserves. *Weed Res* 32:29–38
- Goplen J, Sheaffer CC, Becker RL, Coulter JA, Breitenbach FR, Behnken LM, Johnson GA, Gunsolus JL (2016) Giant ragweed (*Ambrosia trifida*) seed production and retention in soybean and field margins. *Weed Technol* 30:246–253
- Gill GS (1996) Ecology of annual ryegrass. *Plant Prot Q* 11:195–198
- Harrison SK, Regnier EE, Schmoll JT (2003) Postdispersal predation of giant ragweed (*Ambrosia trifida*) seed in no-tillage corn. *Weed Sci* 51:955–964
- Harrison SK, Regnier EE, Schmoll JT, Harrison JM (2007) Seed size and burial effects on giant ragweed (*Ambrosia trifida*) emergence and seed demise. *Weed Sci* 55:16–22
- Harrison SK, Regnier EE, Schmoll JT, Webb JE (2001) Competition and fecundity of giant ragweed in corn. *Weed Sci* 49:224–229
- Hartnett DC, Hartnett BB, Bazzaz FA (1987) Persistence of *Ambrosia trifida* populations in old fields and responses to successional changes. *Am J Bot* 74:1239–1248
- Hartzler R, Liebman M, Westerman P (2007). Weed seed predation in agricultural fields. Iowa State University Extension. <http://www.weeds.iastate.edu/mgmt/2006/seedpredation.pdf>. Accessed December 10, 2015
- Heap I (2015). The International Survey of Herbicide Resistant Weeds. <http://www.weedscience.org>. Accessed December 10, 2015
- Kaiser DE, Lamb JA, Eliason R (2011). *Fertilizer Guidelines for Agronomic Crops in Minnesota*. St. Paul, MN: University of Minnesota Extension BU-06240-5. <http://www.extension.umn.edu/agriculture/nutrient-management/nutrient-lime-guidelines/fertilizer-recommendations-for-agronomic-crops-in-minnesota/docs/BU-6240S-PUB.pdf>. Accessed December 10, 2015
- Kremer RJ (1993) Management of weed seed banks with microorganisms. *Ecol Appl* 3:42–52
- Leighty CE (1938) Crop rotation. Pages 406–430 in, *Soils and Men: Yearbook of Agriculture 1938*. Washington, DC: U.S. Government Printing Office
- Liebman M, Dyck E (1993) Crop rotation and intercropping strategies for weed management. *Ecol Appl* 3:92–122
- Loux MM, Berry MA (1991) Use of a grower survey for estimating weed problems. *Weed Technol* 5:460–466
- Lueschen WE, Andersen RN (1980) Longevity of velvetleaf (*Abutilon theophrasti*) seeds in soil under agricultural practices. *Weed Sci* 28:341–346
- Norsworthy JK, Ward SM, Shaw DR, Llewellyn RS, Nichols RL, Webster TM, Bradley KW, Frisvold G, Powles SB, Burgos NR, Witt WW, Barrett M (2012) Reducing risks of herbicide resistance: best management practices and recommendations. *Weed Sci* 60(sp1): 31–62

- SAS Institute. (2012). The SAS System for Windows. v. 9.4. Cary, NC: SAS Institute
- Schreiber MM (1992) Influence of tillage, crop rotation, and weed management on giant foxtail (*Setaria faberi*) population dynamics and corn yield. *Weed Sci* 40:645–653
- Schutte BJ, Regnier EE, Harrison SK (2012) Seed dormancy and adaptive seedling emergence timing in giant ragweed (*Ambrosia trifida*). *Weed Sci* 60:19–26
- Schutte BJ, Regnier EE, Harrison SK, Schmoll JT, Spokas K, Forcella F (2008) A hydrothermal seedling emergence model for giant ragweed (*Ambrosia trifida*). *Weed Sci* 56:555–560
- Shaner DL, Beckie HJ (2014) The future of weed control and technology. *Pest Manag Sci* 70:1329–1339
- Simmons S, Oelke E, Anderson P (1995). Growth and Development Guide for Spring Wheat. St. Paul, MN: University of Minnesota Extension. <http://www.extension.umn.edu/agriculture/small-grains/growth-and-development/spring-wheat>. Accessed December 10, 2015
- Standifer LC (1980) A technique for estimating weed seed populations in cultivated soil. *Weed Sci* 28:137–138
- Staricka JA, Burford PM, Allmaras RR, Nelson WW (1990) Tracing the vertical distribution of simulated shattered seeds as related to tillage. *Agron J* 82:1131–1134
- Tsimba R, Edmeades GO, Millner JP, Kemp PD (2013) The effect of planting date on maize: phenology, thermal time durations and growth rates in a cool temperate climate. *Field Crops Res* 150:145–155
- University of Minnesota Extension. (2015). Crop Production. <http://www.extension.umn.edu/agriculture/crops>. Accessed December 10, 2015
- Walsh MJ, Powles SB (2007) Management strategies for herbicide-resistant weed populations in Australian dryland crop production systems. *Weed Technol* 21:332–338
- Webster TM, Loux MM, Regnier EE, Harrison SK (1994) Giant ragweed (*Ambrosia trifida*) canopy architecture and interference studies in soybean (*Glycine max*). *Weed Technol* 8:559–564
- Werle R, Sandell LD, Buhler DD, Hartzler RG, Lindquist JL (2014) Predicting emergence of 23 summer annual weed species. *Weed Sci* 62:267–279
- Westerman PR, Liebman M, Menalled FD, Heggenstaller AH, Hartzler RG, Dixon PM (2005) Are many little hammers effective? Velvetleaf (*Abutilon theophrasti*) population dynamics in two- and four-year crop rotation systems. *Weed Sci* 53:382–392

Received May 23, 2016, and approved July 26, 2016.

Associate Editor for this paper: Martin M Williams II, USDA-ARS