

Psychometric validation of a knowledge questionnaire on micronutrients among adolescents and its relationship to micronutrient status of 15–19-year-old adolescent boys, Hyderabad, India

Little Flower Augustine¹, Shahnaz Vazir², Sylvia Fernandez Rao², M Vishnuvardhana Rao³, A Laxmaiah⁴, P Ravinder¹, V Vikas Rao¹ and K Madhavan Nair^{1,*}

¹Division of Micronutrient Research, National Institute of Nutrition, Indian Council of Medical Research, Jamai-Osmania, Hyderabad 500007, Andhra Pradesh, India: ²Division of Behavioral Sciences, National Institute of Nutrition, Indian Council of Medical Research, Hyderabad, India: ³Division of Biostatistics, National Institute of Nutrition, Indian Council of Medical Research, Hyderabad, India: ⁴Division of Community Studies, National Institute of Nutrition, Indian Council of Medical Research, Hyderabad, India

Submitted 23 June 2011: Accepted 19 December 2011: First published online 10 February 2012

Abstract

Objective: To develop a psychometrically valid questionnaire for testing knowledge on micronutrients and to assess the relationship between knowledge and biomarkers of micronutrient status among adolescents.

Design: Cross-sectional, institution-based, validity and reliability study.

Setting: Seven higher secondary schools were covered in the limits of Greater Hyderabad Municipal Corporation, Hyderabad, India.

Subjects: Students aged 15–19 years, *n* 92 for the pre-test, *n* 108 for test–retest and *n* 109 for studying the relationship between knowledge and biomarkers of Fe, retinol, ascorbic acid, α -tocopherol, folic acid and vitamin B₁₂ status.

Results: From an item pool of 106, thirty-one items were selected based on content validity. Statistical tools to obtain a valid and reliable questionnaire among adolescent boys and girls resulted in eighteen items with a difficulty index of 0.11–0.86, discrimination index of 0.20–0.72 and validity index (point bi-serial correlation) of 0.10–0.62. Reliability as measured by Cronbach's α was 0.71 and the intra-class correlation coefficient was 0.80. A Bland–Altman plot showed good agreement between test and retest scores. The mean response score to the eighteen-item questionnaire was 5.2 (SD 2.68). The mean values of serum retinol were significantly different ($P = 0.022$) between groups below (24.8 (SD 6.64) $\mu\text{g}/\text{dl}$) and above (28.0 (SD 7.67) $\mu\text{g}/\text{dl}$) the 50th percentile of knowledge score. The relationship persisted after controlling for economic status as a covariate using analysis of covariance ($P = 0.018$). Other micronutrients did not show any significant relationship.

Conclusions: A valid and reliable eighteen-item knowledge questionnaire was constructed and found to have a significant positive relationship with plasma retinol status alone.

Keywords
Knowledge questionnaire
Adolescents
Micronutrients
Retinol
Psychometric indices
India

Micronutrient deficiencies are rampant in India among all age groups. The intervention strategies include supplementation, fortification and dietary diversification, the latter being the most sustainable approach. This strategy can be promoted through intense education and mass awareness programmes. Lack of culture-appropriate nutrition education has been identified as one of the weakest links of nutrition interventions in India targeting micronutrient malnutrition⁽¹⁾. One of the obstacles for this could be the lack of information on current nutrition knowledge among different population groups, which in

turn can be attributed to the lack of validated assessment tools for testing knowledge among target groups. The need for valid and reliable tools in nutrition education research for assessment of knowledge is now well recognized on account of the realization that the poor association between nutritional knowledge and dietary intake may be due to the poor assessment of knowledge^(2–12).

Development of a valid questionnaire based on pre-defined criteria was suggested by Kline⁽²⁾. Applying these criteria, a general nutrition knowledge questionnaire was

*Corresponding author: Email nairthayil@gmail.com

developed for the US adult population which has been translated, pre-tested and used widely⁽³⁻⁵⁾. Attempts have been made elsewhere to develop reliable questionnaires for nutrition education research, focusing on general nutrition knowledge of adults^(6,7). Tools have also been developed for use among elementary-school teachers and children aimed at measuring the dietary intake or general nutrition knowledge⁽⁸⁻¹⁰⁾. Additionally, there have been needs-based attempts in the past to develop questionnaires addressing specific areas of nutrition knowledge like fat, fibre and cholesterol^(10,11). A questionnaire was also developed for assessment of physical activity, diet, weight-related attitudes and cultural identity among American Indian children⁽¹²⁾.

Furthermore, studies on the relationship between nutrition knowledge and nutritional status are often restricted to either intake data or body weight⁽¹³⁻¹⁵⁾. Studies attempting to relate nutrition knowledge to the biomarkers of nutrients are difficult to find. Such information can however be helpful for strengthening validity in the development of nutrition knowledge questionnaires. A valid and reliable questionnaire can help nutrition scientists to monitor the impact of interventions aimed at improving the dietary intake of micronutrients.

There are no validated tools in the literature to assess knowledge on micronutrients, deficiency of which is an emerging area of concern in developing countries like India. Since adolescence is an age group where maximum behaviour change is possible⁽¹⁶⁾, the aim of the present study was to develop a psychometrically valid and reliable questionnaire for testing knowledge on micronutrients. The study also attempted to apply the validated questionnaire among a group of adolescents whose micronutrient status was already assessed, to understand the relationship between knowledge and biomarkers of micronutrient status.

Methods

All procedures and protocols were approved by the Institutional Ethics Committee. Permission was obtained from the principals of the participating schools and the Directorate of Intermediate Education, Government of Andhra Pradesh – a southern state of India. The study was carried out in 2009–2010 in Government-run schools which cater to students belonging to low- and middle-income families. A written informed consent from parents, and assent from students, was obtained.

Development of questionnaire

The questionnaire was constructed using the method suggested by Kline⁽²⁾ and Garret and Woodworth⁽¹⁷⁾. The process of questionnaire construction consisted of forming an item pool, carrying out a content analysis, and analysing the item difficulty, discriminative power, validity and reliability.

Item pool

The item pool addressed specific micronutrients on the basis of: (i) those that were known to be deficient among the population; (ii) those that had a reliable biomarker; and (iii) intake data being available. Among the fat-soluble vitamins, vitamins A and E were selected. Vitamin C, vitamin B₁₂ and folic acid were addressed among the water-soluble vitamins. Fe was the only mineral considered on account of its low intake and widespread deficiency. The questions were designed within three major domains: (i) functions; (ii) signs of deficiencies; and (iii) sources of the micronutrients. A few questions on general health, nutrition and national programmes were also included (Table 1). An item pool of 106 questions was constructed from the literature, available validated questionnaires and by consulting experts in the field of nutrition. The item pool was subjected to content analysis.

Table 1 Knowledge segments covered under the domains of function, deficiency and food sources in the item pool

No.	Segment	No. of items
1.	Micronutrient deficiency Causal relationship: anaemia, night blindness and scurvy Emerging evidence: non-communicable diseases and role in common infections like cold	27
2.	Awareness on nutrients and nutrition Balanced diet and ability to identify food sources of carbohydrates, protein, fat, vitamins and minerals in the diet	23
3.	Food sources of micronutrients Fe, vitamin A, vitamin C, folic acid, B vitamins, vitamin E and vitamin B ₁₂	21
4.	Micronutrient functions Maintaining general health and vigour, synthesis of Hb, role in cognition, brain development, work capacity, attention concentration and depression	11
5.	Nutrient availability and storage in the body Emphasis on absorption promoters and inhibitors, storage of fat-soluble and water-soluble vitamins	13
6.	Processing losses Washing prior to cooking and cooking per se	6
7.	Awareness on hygiene Hand washing and worm infestation	2
8.	Awareness on national programmes Addressing micronutrient deficiencies – vitamin A, Fe, folic acid and iodine	3
	Total number of items	106

Content analysis

Content analysis was done by the judgement method. The questions were direct and no negative phrasing was used. Two independent reviews were carried out by a panel of five experts from the fields of nutrition, psychology, medicine and basic sciences to select the questions that were appropriate, accurate and interpretable. The acceptance, rejection or modification of an item was based on majority opinion.

Pre-test

The questionnaire was translated to the regional language, Telugu, using the forward-backward translation method and pre-tested in a group of adolescents aged 15–19 years studying in 12th grade. The questionnaire was self-administered. The group consisted of equal number of students from science and commerce streams (n 92) from both genders (thirty-seven girls and fifty-five boys).

Scoring

The thirty-one-item questionnaire consisted of twenty-eight closed-end questions and three open-ended questions. The closed-end questions consisted of sixteen 'yes/no' questions and twelve multiple-choice questions. There was only one correct answer to each question and an option of 'not sure' was used to reduce response bias. The responses were scored by awarding a mark of 1 to the correct response and a mark of 0 to an incorrect or blank response.

Item difficulty index

The item difficulty index (p) was determined by the number of students in the group who performed correctly on the item and expressed as a ratio of correct responses to the total number of questions. An item with a value of p below 0.20 was considered difficult, while p above 0.80 indicated an easy item⁽¹⁷⁾.

Item discrimination index

The steps involved in determining the item discrimination index (D) consisted of identifying the 27% of students who had the highest overall scores and the 27% with the lowest scores, representing the tail from the mean of the standard normal distribution of response error⁽¹⁷⁾. The p levels for the upper and lower groups were determined as described above, and then the p levels of the lower group were subtracted from those of the upper group; the resulting values were considered as D . The higher the D , the more discriminating was the item. A value of $D = 0.20$ – 0.29 was considered 'acceptable', $D = 0.30$ – 0.39 as 'good' and $D > 0.40$ was considered 'excellent' as per available literature⁽¹⁸⁾.

Item validity index

The validity index of each item (calculated as point bi-serial r) was determined by the extent to which the given

item could discriminate among students who differed sharply in the function measured by the test as a whole and was determined by the inter-correlation between each item in a scale⁽⁷⁾. Point bi-serial r is mathematically equivalent to Pearson product moment correlation and is used when one of the variables is dichotomous. A point bi-serial r of 0.2 was considered valid⁽²⁾ except in circumstances where an item was considered essential in terms of content validity. In such cases, an r value of ≥ 0.1 was considered a minimum requirement as described in the literature⁽¹⁷⁾.

Internal consistency

The rational equivalence method using Cronbach's α was employed to assess the reliability of the test. Cronbach's α estimates the reliability of test scores with respect to (i) how well the individual items of the scores fit together and (ii) whether they assess the same construct. It assesses the inter-correlations of the items in the test and the correlations of the items with the test as a whole. A value of α above 0.70 was considered acceptable on the internal consistency as per available literature^(2,19) and was done using the SPSS statistical software package version 16.0 (SPSS Inc., Chicago, IL, USA).

Test-retest reliability

Considering a minimum of five observations for each item and a drop-out rate of 20%, in total 108 students were considered sufficient for the study on test-retest reliability of the eighteen-item questionnaire that was developed. The questionnaire was self-administered. The time taken to complete the questionnaire was noted down using a wrist watch. The students were kept unaware of the retest that was done after a gap of 7 d⁽⁹⁾. A total of 100 students completed both the contacts of the test-retest. A Bland-Altman plot was constructed and the intra-class correlation coefficient (ICC) was calculated for assessing the agreement between the scores^(20,21). An ICC above 0.75 was considered acceptable⁽²¹⁾.

Relationship between biomarkers and knowledge on micronutrients

The knowledge questionnaire was applied to a group of students participating in a 2-year study on 'Stress, allostatic load and micronutrient status among students: impact of dietary advice'. For this, five single-gender schools catering to male students from three different geographical locations of Greater Hyderabad Municipal Corporation, Andhra Pradesh, India were selected. Students with congenital anomalies, physical handicaps or any other chronic health problems requiring medical attention and those taking micronutrient tablets for the past year were excluded from the study. A total of 380 students were recruited for the study using probability proportional to size sampling, from whom a blood sample was collected for analysis of micronutrient status in a randomly selected sub-sample of 150. One hundred

and nine of these participants were available for testing the validated knowledge questionnaire, which was done after a gap of 6 months.

Micronutrient status

The baseline blood samples were collected after an overnight fast and analysed for Hb⁽²²⁾ and plasma ferritin⁽²³⁾, folic acid and vitamin B₁₂ (RIA kit by Siemens Inc., LA, USA). Simultaneous determination of plasma retinol and α -tocopherol was carried out by HPLC (Thermo Finnigan, Herts, UK)⁽²⁴⁾. An ascorbic acid assay was done on the same day using the α,α -bipyridyl micromethod⁽²⁵⁾. The ferritin ELISA and HPLC methods for retinol and α -tocopherol were in-house methods and are continuously validated by the VITAL-EQA (Vitamin A Laboratory – External Quality Assurance) programme of the Centers for Disease Control and Prevention (Atlanta, GA, USA). The percentage inadequacy of micronutrient status was computed using cut-off values derived from the literature⁽²⁶⁾.

Economic status

Economic status was derived from the information collected on twelve household assets by computing the standard of living index⁽²⁷⁾.

Statistical analysis

All statistical analyses were done using SPSS version 16.0. Descriptive statistics were used to summarize the results. Item analysis (item difficulty index p , item discrimination index D , point bi-serial r and Cronbach's α) was done for psychometric validation of the thirty-one-item questionnaire. The Bland–Altman plot was constructed using the MedCalc statistical program version 11.5.1.0 (MedCalc Software, Mariakerke, Belgium). The relationship between knowledge and biomarkers of micronutrient status was tested by the two-tailed independent-samples t test between groups above and below the 50th percentile of knowledge score. For micronutrients which were not normally distributed, the non-parametric Mann–Whitney U test was used. Analysis of covariance was performed taking the standard of living index as a covariate, knowledge as the independent variable and micronutrient status as the dependent variable.

Results

The content analysis of the 106 items resulted in the selection of six multiple-choice questions, twenty-one yes/no questions and four one-word-answer type questions. There was no significant gender difference in the scores of respondents during the pre-test ($P=0.995$) and therefore only pooled analysis was done.

Psychometric indices

The item difficulty index p ranged from 0.11 to 0.86. There were three relatively easy questions ($p=0.84$ – 0.86) and

three difficult questions ($p=0.11$ – 0.19) and all the rest were in the desirable range of difficulty (Table 2).

The item discriminative index D ranged from -0.32 to 0.72 . Eight items with a discriminative index below 0.20 were discarded as poorly discriminating questions (Table 2).

Twelve questions had item validity index r above 0.2 . Due to good content validity, six items in the range of r from 0.1 to 0.2 were retained, resulting in an eighteen-item questionnaire (Table 3).

A Cronbach's α value of 0.71 was obtained for the eighteen-item scale. The domains were combined in the shortened version. The Bland–Altman plot showed a mean of 0.8 with few outliers (Fig. 1). The ICC of 0.80 was significant at $P<0.01$. The final questionnaire consisted of twelve items on functions/deficiencies, five items on dietary source and one question on hand washing (hygiene). The questionnaire took about 15 min to complete.

Relationship between the valid eighteen-item questionnaire and micronutrient status

The mean age of the students was 16.6 (SD 1.24) years; their mean height was 163.7 (SD 6.95) cm, mean weight was 47.5 (SD 7.37) kg and mean BMI was 17.7 (SD 2.23) kg/m².

The knowledge scores of the students ranged from 0 to 14 , with a mean of 5.2 (SD 2.68). Fifty-nine per cent of the students scored below the 50th percentile of knowledge score, i.e. a score of less than 5.0 .

Micronutrient status

The mean values of micronutrients were 13.9 (SD 1.63) g/dl for Hb, 44.6 (SD 80.1) ng/ml for ferritin, 0.42 (SD 0.29) mg/dl for ascorbic acid, 26.3 (SD 7.4) μ g/dl for plasma retinol, 328.9 (SD 164.42) μ g/dl for α -tocopherol, 5.5 (SD 1.9) ng/ml for folic acid and 246.0 (SD 110.00) pg/ml for vitamin B₁₂.

Knowledge and micronutrient status among students

Plasma retinol status was significantly different between the groups below and above the 50th percentile of knowledge score ($P=0.022$). Other biomarkers of micronutrient status did not show any significant relationship with knowledge score (Table 4). The analysis of covariance performed with micronutrient status as the dependent variable and standard of living index as a covariate also confirmed a significant difference in retinol status based on knowledge score ($P=0.018$).

Discussion

Several studies assessing knowledge have been found to report on a questionnaire being 'pre-tested' without providing the statistical attributes that establish its reliability and validity^(28–31). A structured approach of scaling is of prime importance for the construction of a knowledge questionnaire in order to establish its validity. To our understanding, there are no validated questionnaires

Table 2 Difficulty index and discrimination index values of questions in the knowledge questionnaire

No.	Item	<i>p</i>	<i>D</i>	No.	Item	<i>p</i>	<i>D</i>
1.	Example of a vitamin	0.61	0.24	16.	Animal foods as the major food source of vitamin B ₁₂	0.30	0.32
2.	Iron as an essential mineral for the body	0.86	0.32	17.	Vitamin C as the absorption promoter of iron†	0.44	0.24
3.	Low intake of iron as a reason for anaemia	0.57	0.60	18.	Groundnuts as a source of vitamin E	0.51	0.04*
4.	Less activity exhibited by anaemic	0.39	0.04*	19.	Fruits & vegetables as a source of B vitamins	0.74	-0.32*
5.	Iron as an integral part of haemoglobin	0.41	0.72	20.	Washing hands for prevention of infections (iron)	0.47	0.56
6.	Familiarity on folic acid	0.84	0.20	21.	Beta-carotene as pro-vitamin A†	0.39	0.24
7.	Role of folic acid in red blood cell formation	0.39	0.24	22.	Colour code for beta-carotene	0.20	0.08*
8.	Vitamin A deficiency as the cause for night blindness	0.45	0.28	23.	Lime as a good source of vitamin C	0.68	0.08*
9.	Familiarity with 'fat-soluble vitamins'	0.75	0.32	24.	Cooking losses of B vitamins†	0.43	0.20
10.	Example of a fat-soluble vitamin	0.22	0.60	25.	Fresh vegetables & fruits as source of B vitamins	0.79	0.04*
11.	Familiarity with 'water-soluble vitamins'	0.37	0.36	26.	Cooking loss of vitamin C	0.30	0.44
12.	Example of a water-soluble vitamin	0.19	0.52	27.	Banana as an easily available source of B vitamins	0.49	0.52
13.	Phrynoderma as a deficiency of essential fatty acid and vitamin E	0.27	0.08*	28.	Food as major source of micronutrients for the body†	0.58	0.20
14.	Vitamin C deficiency as a cause for spongy bleeding gums	0.37	0.44	29.	Fruits and vegetables as source of vitamins and minerals	0.86	0.16*
15.	Vitamin C deficiency pre-disposing to infection†	0.28	0.24	30.	Cereals & pulses as starch-rich foods	0.13	0.20
				31.	Pulses as protein-rich foods	0.11	0.24

The item difficulty index *p* is the ratio of the number of correct responses to the total number of responses (*n* 92). The higher the index value, lower the difficulty. *D* is the item discrimination index.

*Indicates the items discarded due to low *D*.

†Indicates the items discarded due to low item validity index (*r*) value of 0.0335 to 0.1319.

Table 3 Summary of psychometric properties of the eighteen selected items

Index	Category	Range	<i>n</i>
Item difficulty index, <i>p</i> ⁽¹⁷⁾	Difficult	<0.20	3
	Ideal	0.20-0.79	13
	Easy	>0.80	2
Item discrimination index, <i>D</i> ⁽¹⁸⁾	Acceptable	0.20-0.29	7
	Good	0.30-0.39	3
	Excellent	>0.40	8
Item validity index, <i>r</i> ^{*(17)}	Low	0.1-0.2	6
	Good	0.2-0.7	12

n, number of items.

*Values less than 0.1 not considered.




Fig. 1 Bland-Altman plot showing the agreement between test and retest scores on the knowledge questionnaire on nutrients among 15-19-year-old students, Hyderabad, India (*n* 100). The differences between the retest and test scores were plotted against the means and the limits of agreement were calculated. — represents the mean difference between the retest and the test (bias); - - - represent the 95% confidence interval of bias; - - - - represent the upper and lower 95% limits of agreement

addressing micronutrients, deficiency of which is of major concern in developing countries like India. Therefore, our study is the first of its kind in constructing and psychometrically validating a knowledge questionnaire on micronutrients. Further, we also tested the relationship of knowledge assessed by this questionnaire with biomarkers of micronutrient status among adolescents.

The item pool used for the questionnaire development addressed the micronutrients which had an established evidence of deficiency or where a national database was available with reference to the intake or widespread deficiency⁽³²⁾. Even though Zn and vitamin D deficiencies are emerging problems in India, there was no national database on intake or deficiency nor was there any RDA for these nutrients in India at the time of designing the study. In addition, there are no reliable markers for mapping Zn deficiency⁽³³⁾. For these reasons we did not

Table 4 Indicators of micronutrient status according to knowledge score below or above the 50th percentile: 15–19-year-old students, Hyderabad, India

Biomarker	Below 50th percentile		Above 50th percentile		P value
	Mean	SD	Mean	SD	
Hb g/dl	13.7	2.36	14.1	2.64	0.346
Ascorbic acid (mg/dl)	0.42	0.282	0.42	0.308	0.758
Retinol (μ g/dl)	24.8	6.69	28.0	7.67	0.022*
α -Tocopherol (μ g/dl)	326	200.0	331	99.5	0.369
Folic acid (ng/ml)	5.4	1.84	5.6	2.09	0.418
Vitamin B ₁₂ (pg/ml)	244	108.1	248	114.9	0.946
Ferritin (ng/ml) [†]	46.9	101.47	40.4	35.06	0.758

The knowledge score 'below 50th percentile' indicates a score of ≤ 5.0 (n 64) and 'above 50th percentile' indicates a score of > 5.0 (n 45). Comparisons are made between the means along the rows.

*Significantly different means.

[†]The high variation observed is due to one extreme value of 821 ng/ml. The data were analysed after logarithmic transformation.

include Zn and vitamin D in the item pool. Iodine was not included on account of the universal salt iodization programme⁽³⁴⁾ currently in operation in the country.

The selection of questions from the item pool was perhaps the most difficult step involved in our study. The expertise of the panel with regard to the domain of knowledge tested and the anticipated difficulty level of the item were considered while selecting questions. The nutrition component in the school curriculum, and nutrition messages propagated by the Government of India through the popular electronic media (television), were the basis for this. Accordingly, a number of questions on dietary sources that were judged to be inappropriate were deleted from the item pool. Questions related to nutrient absorption were also deleted as they were judged to be too technical for the group being studied. Care was also taken to select items which were simple, easily understandable and suitable for 15–19-year-old adolescent students of low and middle income level of economic status. Thus the selected thirty-one questions mostly attempted to test basic knowledge on micronutrients.

To obtain a valid questionnaire applying appropriate statistical methods, the thirty-one items were pre-tested in a group of adolescents. The statistical tools applied, such as item difficulty index, item discrimination index, item validity index and reliability test, provided an eighteen-item questionnaire. The spread of data, difficulty level and discriminative power of this eighteen-item questionnaire were within acceptable ranges. Nine items belonging to the domain of dietary sources did not show significant item to total correlation and were discarded. Some of the items had low inter-correlations but were retained due to content validity. The reliability of the questionnaire of $\alpha = 0.71$ was also within the acceptable range and is in agreement with the widely accepted general nutrition knowledge test⁽³⁾ and a knowledge questionnaire developed for US children which had an α of 0.7⁽¹²⁾. Thus we constructed a short self-administered questionnaire that required 15 min to complete with no gender-specific items and so can be used for adolescents.

This had an added advantage of easy assessment of knowledge useful in pre–post knowledge assessment studies on micronutrients.

In an attempt to understand whether true knowledge will be reflected in biomarkers of micronutrient status, we applied this questionnaire in a group of adolescent boys. The prevalence of anaemia among this study group was 21%, which is similar to the anaemia prevalence among males (15–49 years) in the State of Andhra Pradesh as well as the national average⁽³⁴⁾. Further, the prevalence of subclinical vitamin A deficiency of 17.5% and ascorbic acid deficiency of 75.0% reported in the present study were similar to those in a study among apparently healthy adults from western India which reported 21% and 77% respectively⁽³⁵⁾. The other micronutrient deficiencies observed in the study population were similar to those reported in the literature in a similar group from a similar location⁽³⁶⁾, implying the representativeness of the study sample.

Among the micronutrients tested, plasma retinol alone showed a positive relationship with knowledge score, suggesting that knowledge on micronutrients can influence the status of at least some nutrients. This relationship persisted after controlling for economic status, a known confounder for micronutrient status on account of affordability and thereby intake⁽³⁷⁾.

Evidence relating knowledge, dietary intake and biomarkers is meagre except for one study among nineteen volunteers that evaluated the impact of dietary counselling among cancer survivors, with an experimental group receiving dietary counselling to increase fruit and vegetable consumption. The plasma levels tested were for carotenoids, vitamin C, α -tocopherol and cholesterol. Plasma carotenoid levels and ascorbic acid showed a change with acceptance and compliance to diet therapy⁽³⁸⁾. Another study from western India, where population characteristics were similar to our study group, examined the relationship between dietary habits, plasma retinol and plasma ascorbic acid among 214 men and 108 women. Results showed a dose–response relationship of intake with

serum retinol alone and not with vitamin C⁽³⁵⁾. The findings were explained on the basis that vitamin A is fat soluble, can be stored in the body and therefore better reflects the dietary habits of the individual. The present study findings can also be explained along similar lines, with knowledge influencing dietary habits which are reflected in biomarker levels. Data on fat intake were not collected. However, the study in western India was done in a population where socio-economic status was similar to our study participants, and it is believed that fat intake would also be similar in both groups. In the absence of any other reported literature, it would not be wrong to consider retinol status as a good biomarker sensitive to knowledge on micronutrients and which may change with changes in knowledge following dietary advice.

The implications of our study findings are varied. First, they emphasize the need to use validated questionnaires in nutrition research involving qualitative aspects like knowledge. Second, they show that knowledge on micronutrients, if assessed correctly, may relate to the blood levels of specific micronutrients of the target group as reported herein for plasma retinol. Increasing the knowledge of the population increases the sustainability of interventions as well as voluntary compliance by the target population.

Although parental factors can influence food choices, the participants of the present study were close to adulthood and therefore culturally possessed a certain amount of autonomy in their food choices. The majority of the participants had a higher educational qualification than their parents and contributed in performing family food chores. This background information on the study participants gives us sufficient confidence in using them as 'change agents'.

There are limitations that need to be considered when interpreting the study results. The generalizability of the test-retest data may be limited since the study was done in girls alone, even though the pre-test showed no gender difference in scoring. Dietary intake data were not collected, which could have provided more conclusive evidence for the relationship we found. A positive relationship of knowledge with micronutrients like folic acid (where body stores are not available) cannot be ruled out, on account of the time lag of 6 months between the assessments of micronutrient status and knowledge.

Acknowledgements

The authors wish to acknowledge the funding received from the Indian Council of Medical Research for carrying out the study. The Fellowship received by L.F.A. from the University Grants Commission, Government of India, is also acknowledged. The authors declare that there is no conflict of interest. All authors have participated in the research reported and all have agreed to be an author of

this paper. K.M.N. conceptualized and designed the study, supervised the study protocol and edited the manuscript. S.V. conceptualized the psychometric validation of the test scale, was involved in development of the questionnaire at various stages and edited the manuscript. L.F.A. was responsible for designing the questionnaire, conduct of studies, analysis of ferritin and has written the manuscript. A.L. did clinical examination for the project and applied inclusion/exclusion criteria. M.V.R. was responsible for data analysis. S.F.R. was responsible for application of the validated scale in the study on 'Stress, allostatic load and micronutrient status among students: impact of dietary advice' and was involved in interpreting the test scales. P.R. analysed biomarkers of micronutrient status and V.V.R. analysed plasma retinol and α -tocopherol. All authors have read and contributed to the final manuscript. The authors thank the Nutrition Society of India for the opportunity to present part of the results of the study at its 42nd Annual Conference held in Mumbai, India, 19–20 November 2010. They also thank Dr B. Sesikeran MD, Director, National Institute of Nutrition, for his support and encouragement.

References

1. Vijayaraghavan K (2002) Control of micronutrient deficiencies in India; obstacles and strategies. *Nutr Rev* **60**, 5 Pt 2, S73–S76.
2. Kline P (1993) *The Handbook of Psychological Testing*. London: Routledge.
3. Parmenter K & Wardle J (1999) Development of a general nutrition knowledge questionnaire for adults. *Eur J Clin Nutr* **53**, 298–308.
4. Hendrie GA, Cox DN & Coveney J (2008) Validation of the general nutrition knowledge questionnaire in an Australian community sample. *Nutr Diet* **65**, 72–77.
5. Shaaban SY, Nassar MF, Abd Elhamid DM *et al.* (2009) Nutritional knowledge and attitude of adolescent school girls living in Cairo. *Res J Med Med Sci* **4**, 421–427.
6. Turconi G, Celsa M, Rezzani C *et al.* (2003) Reliability of a dietary questionnaire on food habits, eating behaviour and nutritional knowledge of adolescents. *Eur J Clin Nutr* **57**, 753–763.
7. Anderson AS, Bell A, Adamson A *et al.* (2001) A questionnaire assessment of nutrition knowledge – validity and reliability issues. *Public Health Nutr* **5**, 497–503.
8. Lambert LG, Monroe A & Wolff L (2010) Mississippi elementary school teachers' perspectives on providing nutrition competencies under the framework of their school wellness policy. *J Nutr Educ Behav* **42**, 271–276.
9. Zarnowiecki D, Dollman J & Sinn N (2011) A tool for assessing healthy food knowledge in 5–6-year-old Australian children. *Public Health Nutr* **14**, 1177–1183.
10. Resnicow K, Hearn M, Delano RK *et al.* (1997) Development of a nutrition knowledge scale for elementary school students: toward a national surveillance system. *J Nutr Educ* **28**, 156–164.
11. Steenhuis IHM, Brug J, Van Assema P *et al.* (1996) The validation of a test to measure knowledge about the fat content of food products. *Nutr Health* **10**, 331–339.
12. Stevens J, Cornell CE, Story M *et al.* (1999) Development of a questionnaire to assess knowledge, attitudes, and behaviors in American Indian children. *Am J Clin Nutr* **69**, 4 Suppl., 773S–781S.

13. O'Brien G & Davies M (2007) Nutrition knowledge and body mass index. *Health Educ Res* **22**, 571–575.
14. Nuss H, Freeland-Graves J, Clarke K *et al.* (2007) Greater nutrition knowledge is associated with lower 1-year postpartum weight retention in low-income women. *J Am Diet Assoc* **107**, 1801–1806.
15. Worsley A (2002) Nutrition knowledge and food consumption: can nutrition knowledge change food behaviour? *Asia Pacific J Clin Nutr* **11**, Suppl. 3, S579–S585.
16. Planning Commission (2002–2007) *Report of the Sub-group on Adolescents for the Tenth Five Year Plan*. New Delhi: Planning Commission, Government of India; available at http://planningcommission.nic.in/aboutus/committee/wrkgrp/wg_adolcntr.pdf
17. Garret HE & Woodworth RS (1969) The reliability and validity of test scores. In *Statistics In Psychology and Education*, pp. 337–370. New York: David McKay Company Inc.
18. Ebel RL (1972) *Essentials of Educational Measurement*, 1st ed. Englewood Cliffs, NJ: Prentice Hall.
19. Wilson AM, Magarey AM & Masterson N (2008) Reliability and relative validity of a child nutrition questionnaire to simultaneously assess dietary patterns associated with positive energy balance and food behaviours, attitudes, knowledge and environments associated with healthy eating. *Int J Behav Nutr Phys Act* **5**, 5.
20. Myles PS & Cui J (2007) Using the Bland–Altman method to measure agreement with repeated measures. *Br J Anaesth* **99**, 309–311.
21. Rosner B (2006) Multisample inference. In *Fundamentals of Biostatistics*, 6th ed., pp. 568–571. Belmont, CA: Duxbury Thomson-Brooks/Cole.
22. International Nutritional Anaemia Consultative Group (1985) *Measurement of Iron Status. A Report of the International Nutritional Anaemia Consultative Group (INACG)*, pp. 4–8. Washington, DC: The Nutrition Foundation Inc.
23. Pawashe AB, Raman L, Nair M *et al.* (1987) Validity of using capillary blood for the measurement of plasma ferritin. *Clin Chim Acta* **163**, 119–120.
24. Bieri JG, Tolliver TJ & Catignani GL (1979) Simultaneous determination of α -tocopherol and retinol in plasma or red cells by high pressure liquid chromatography. *Am J Clin Nutr* **32**, 2143–2149.
25. Zannoni V, Lynch M & Goldstein S (1974) A rapid micromethod for the determination of ascorbic acid in plasma and tissues. *Biochem Med* **11**, 41–48.
26. Young DS (1987) Implementation of SI units for clinical laboratory data, style specifications and conversion tables. *Ann Intern Med* **106**, 114–129.
27. International Institute for Population Sciences & ORC Macro (2000) *National Family and Health Survey (NFHS-2)*. Mumbai: IIPS.
28. Lakshman RR, Sharp SJ, Ong KK *et al.* (2010) A novel school-based intervention to improve nutrition knowledge in children: cluster randomised controlled trial. *BMC Public Health* **10**, 123.
29. Lanerolle P & Atukorala S (2006) Nutrition education improves serum retinol concentration among adolescent school girls. *Asia Pac J Clin Nutr* **15**, 43–49.
30. Joshi N & Vijayalaxmi KG (2009) Nutritional education tool to improve overall dietary attitude and knowledge among young women. *J Hum Ecol* **25**, 187–191.
31. Nazni P & Vimala S (2010) Nutrition knowledge, attitude and practice of college sportsmen. *Asian J Sports Med* **1**, 93–100.
32. National Nutrition Monitoring Bureau (2003) *Prevalence of Micronutrient Deficiencies. NNMB Technical Report Series* no. 21. Hyderabad: National Institute of Nutrition; available at <http://www.nnmbindia.org/NNMB%20MND%20REPORT%202004-Web.pdf>
33. International Zinc Nutrition Consultative Group, Brown KH, Rivera JA *et al.* (2004) International Zinc Nutrition Consultative Group (IZiNCG) technical document #1. Assessment of the risk of zinc deficiency in populations. *Food Nutr Bull* **25**, 1 Suppl. 2, S130–S161.
34. International Institute for Population Sciences & ORC Macro (2007) *National Family and Health Survey (NFHS-3)*. Mumbai: IIPS.
35. Chiplonkar SA, Agte VV, Mengale SS *et al.* (2002) Are lifestyle factors good predictors of retinol and vitamin C deficiency in apparently healthy adults? *Eur J Clin Nutr* **56**, 96–104.
36. Sivakumar B, Nair KM, Sreeramulu D *et al.* (2006) Effect of micronutrient supplement on health and nutritional status of school children: biochemical status. *Nutrition* **22**, 1 Suppl., S15–S25.
37. Monsivais P, Aggarwal A & Drewnowski A (2010) Low micronutrient intake is associated with low socio-economic status and low diet cost. *FASEB J* **24**, lb 359.
38. Le Marchand L, Hankin JH, Carter FS *et al.* (1994) A pilot study on the use of plasma carotenoids and ascorbic acid as markers of compliance to a high fruit and vegetable dietary intervention. *Cancer Epidemiol Biomarkers Prev* **3**, 245–251.