

Modeling of the IR Light Curves of the Symbiotic Recurrent Nova T CrB

A. A. Tatarnikova, A. M. Tatarnikov, and V. I. Shenavrin

Lomonosov Moscow State University, Sternberg Astronomical Institute, Moscow, Russia
email: aat@sai.msu.ru

Abstract. We analyze the results of our IR photometric monitoring of T CrB during 1987-2003 and describe the ellipsoidal variability of the Roche lobe filling cool component. We obtain limits to the binary inclination of $i \in [50, 60]$ deg and binary mass ratio $q \in [0.4, 2]$ (90 per cent confidence). The mass of the hot component is therefore 1.3-3 M_{\odot} . If the hot component of T CrB is a white dwarf, its mass will be near the Chandrasekhar limit.

Keywords. binaries: symbiotic — stars: individual (T CrB)

1. Observations

T Coronae Borealis is a recurrent nova which underwent two outbursts, one in 1866 and one in 1946. The double bump visible in the optical and IR light curves indicates that the giant of T CrB is tidally distorted (Kraft 1958; Yudin & Munari 1993).

Photometric JK observations of T CrB have been carried out with the 1.25-m telescope at the Crimean Station of the Sternberg Astronomical Institute during 1987 - 2003. The IR standard is BS 5947 ($J = 2.09$ mag, $K = 1.30$ mag). The observational errors do not exceed 0.02 mag.

2. Calculations

A theoretical light curve for a tidally distorted red giant was calculated in the frames of the classical model of the ellipsoidal variability (Tjemkes *et al.* 1986). According to the radial-velocity analyses, we assumed that the orbit of T CrB is circular. Each local surface of the red giant radiates as a black body with a temperature determined by Lucy's law for stars with convective envelopes (the mean temperature is equal to the effective temperature $T_{eff} = 3300$ K). The J band is the most suitable band for approximation of a real red giant's energy distribution by black body radiation (Shahbaz *et al.* 1997). A non-linear limb-darkening law has been assumed and the coefficients have been interpolated from the tables of Claret for the temperatures on the local surfaces (Claret 2000). Figure 1 shows a convolution of the J light curve with a period $P(J) = 227.^d56$, which was searched by Kolpakov's program (<http://infra.sai.msu.ru/prog/kolpakov>).

In Figure 2 we plotted two areas of possible values for binary parameters (q, i) according to Fisher's distribution (dark grey – 75% confidence limit, light grey – 90% confidence limit). The lack of eclipses in the UV continuum observed with International Ultraviolet Explorer indicates that T CrB is not an eclipsing binary (Selvelli *et al.* 1992). The limits based on the lack of eclipses are shown in Figure 2 as a dashed line. The parameters (q, i) of T CrB must be below this curve. A reasonable limit to the mass of the red giant is $M_{giant} \geq 0.6 M_{\odot}$ (the secondary can evolve to a giant during the lifetime of our Galaxy). Using the mass function derived by Fekel *et al.* (2000) we plotted the line corresponding to $M_{giant} = 0.6 M_{\odot}$ (solid line). The dotted lines are the lines of the constant hot component masses (thick dotted line – $M_{hot} = 1.44 M_{\odot}$).




Figure 1. The J light curve of T CrB folded with orbital period $P = 227.^d56$ and the synthetic light curve for the model with $q = 0.5$, $i = 53$, $T_{eff} = 3300 K$




Figure 2. The parameter space diagram (q - i) for T CrB (dark grey – 75% confidence limit, light grey – 90% confidence limit). The solid line corresponds to $M_g = 0.6 M_\odot$. The dashed line is the limit based on the lack of eclipses. The dotted lines are the lines of the constant hot component masses (thick dotted line is $M_h = 1.44 M_\odot$).

From Figure 2 we conclude that, if the hot component of T CrB is a white dwarf its binary parameters will be $q \in [0.5, 0.8]$ and $i \in [52, 62]$. It should be noted that the hot component's mass is not less than $1.25 M_\odot$. Therefore, according to Iben & Tutukov (1996), T CrB can be considered as a real progenitor of a Type Ia supernova.

References

- Claret, A. 2000, *A&A*, 363, 1081
 Fekel, F. C., Joyce, R. R., & Hinkle, K. H. and Skrutskie, M. F. 2000, *AJ*, 119, 1375
 Iben, I. & Tutukov, A. V. 1996, *ApJS*, 105, 145
 Kraft, R. P. 1958, *ApJ*, 127, 620
 Selvelli, P. L., Cassatella, A., & Gilmozzi, R. 1992, *AJ*, 393, 289
 Shahbaz, T., Somers, M., Yudin, B. F., & Naylor, T. 1997, *MNRAS*, 288, 1027
 Tjemkes, S. J., van Paradijs, J. & Zuiderwijk, E. J. 1986, *A&A*, 154, 77
 Yudin, B. F. & Munari, U. 1993, *A&A*, 270, 165