

## RESONANT SCATTERING AND RECOMBINATION IN CAL 87

J. Greiner,<sup>1</sup> A. Iyudin,<sup>1,2</sup> M. Jimenez-Garate,<sup>3</sup> V. Burwitz,<sup>1</sup> R. Schwarz,<sup>4</sup> R. DiStefano,<sup>5</sup> and N. Schulz<sup>3</sup>

### RESUMEN

Se observó la binaria eclipsante de rayos X supersuave CAL 87 con Chandra el 13/14 de agosto de 2001, durante casi 100 ksec, completando dos ciclos orbitales totales y tres eclipses. La forma de la curva de luz del eclipse derivada de los fotones de orden cero indica que el tamaño de la región de emisión de rayos X es de alrededor de  $1.5 R_{\odot}$ . El espectro ACIS/LETG está dominado por completo por las líneas de emisión sin ningún continuo apreciable. Las líneas de emisión más brillantes presentan un corrimiento al rojo significativo y dos picos, sugiriendo que se producen en un viento de 2000 km/s. Realizamos un modelo del espectro de rayos X mezclando la dispersión de recombinación y resonancia. Esto nos permite deducir que la temperatura y la luminosidad de la fuente ionizante son  $kT \sim 50 - 100$  eV y  $L_X \sim 5 \times 10^{37}$  erg/s, respectivamente.

### ABSTRACT

The eclipsing supersoft X-ray binary CAL 87 has been observed with Chandra on August 13/14, 2001 for nearly 100 ksec, covering two full orbital cycles and three eclipses. The shape of the eclipse light curve derived from the zeroth-order photons indicates that the size of the X-ray emission region is about  $1.5 R_{\odot}$ . The ACIS/LETG spectrum is completely dominated by emission lines without any noticeable continuum. The brightest emission lines are significantly redshifted and double-peaked, suggestive of emanating in a 2000 km/s wind. We model the X-ray spectrum by a mixture of recombination and resonant scattering. This allows us to deduce the temperature and luminosity of the ionizing source to be  $kT \sim 50 - 100$  eV and  $L_X \sim 5 \times 10^{37}$  erg/s.

*Key Words:* BINARIES: CLOSE — STARS: INDIVIDUAL (CAL 87) — X-RAY: STARS

### 1. INTRODUCTION

CAL 87 was detected with the IPC onboard the Einstein X-ray observatory during the Columbia Astrophysical Laboratory survey of LMC (Long et al. 1981), and it was optically identified with an eclipsing binary in the Large Magellanic Cloud with an orbital period of 10.6 hrs (Pakull et al. 1988, Callanan et al. 1989, Cowley et al. 1990). A shallow X-ray eclipse was discovered with ROSAT (Schmidtke et al. 1993).

Early attempts to model the optical light curve of CAL 87 by Callanan et al. (1989) already indicated an elevated accretion disk rim. This model has been extended by Schandl et al. (1997), and successfully describes the optical light curve of the CAL 87 binary system, composed of a primary (WD) with a mass of  $M_1 = 0.75 M_{\odot}$ , placed at a distance of  $2.2 \times 10^{11}$  cm from the mass-donating secondary star with a mass of  $M_2 = 1.5 M_{\odot}$ . The model of Schandl et al. (1997) includes (1) optical emission from the secondary star which is irradiated by the emission of the WD and (2) emission of an accretion disk with a thick rim

and an optically thick, cold, clumpy spray produced by the high mass-flow rate of the accretion stream impinging on the disk (hot spot). This spray, moving around the disk, nicely reproduced the asymmetry in the optical light curve and the depth of the secondary dip.

Based on the pre-Chandra, low-resolution X-ray spectra and the detection of emission up to 1 keV, CAL 87 has long been considered as one of the two hottest known SSS, thus spurring observations with all X-ray satellites since ROSAT (Parmar et al. 1997, Asai et al. 1998, Dotani et al. 2000, Ebisawa et al. 2001).

We have exploited the unique spectral capability of the Chandra low-energy transmission grating (LETG) to study in detail the X-ray emission of the canonical supersoft source CAL 87.

### 2. CHANDRA OBSERVATION IN 2001

We observed CAL 87 with Chandra for 97 ksec on August 13/14, 2001, using the low-energy transmission grating (LETG/ACIS). This observation covered three consecutive eclipses of the binary system. The phase-folded light curve of the zeroth order photons (Fig. 1) shows a wide eclipse with triangular shape, about  $\pm 0.15$  phase units wide (30% of the orbit), no flat bottom and an eclipse depth of about

<sup>1</sup>MPE Garching, 85741 Garching, Germany.

<sup>2</sup>Moscow, Russia.

<sup>3</sup>MIT, Cambridge, MA 02139, USA.

<sup>4</sup>Universität Göttingen, 37083 Germany.

<sup>5</sup>CfA, Cambridge, MA 02138, USA.






Fig. 3. Preliminary model of the X-ray spectrum of CAL 87, composed of recombination (blue) and resonant scattering (black) with a white dwarf mass of  $0.8 M_{\odot}$  (top) and  $1.4 M_{\odot}$  (bottom), corresponding to different effective temperatures of the ionizing source. The dashed line in the lower panel is the primary white dwarf spectrum.

### 3. CONCLUSIONS

The shape of the eclipse light curve, the lack of any continuum emission and the dominance of emission lines due to resonant scattering and recombination suggests that CAL 87's X-ray emission comes from an extended wind (or outflow), similar to the accretion-disk-corona low-mass X-ray binary 4U 1822-37, as already suggested by Schmidtke (1993). However, the plasma in 4U 1822-37 (i) has a much higher ionization as evidenced by the H- and He-like ions of Ne, Mg, Si, S and Fe which are completely absent in CAL 87 and (ii) is much more dominated by recombination, indicative of larger column densities than those in CAL 87.

The wind is probably not spherical, but rather has a conical shape with an opening angle of  $120^{\circ}$ , extending on both sides of the accretion disk. During eclipse, the donor occults the wind on the southern disk side.

The primary X-ray emission region in CAL 87 certainly has a temperature substantially lower than  $\sim 1$  keV due to the lack of Fe L, Si XIII or Ne IX/Ne X lines. This rules out a neutron star or black hole (except for retrograde rotation) accretor, since they would exhibit a maximum temperature of the inner part of the accretion disk of 0.9–1.2 keV. Also, the outflow velocity of  $\sim 2000$  km/s is consistent with a white dwarf primary.

A corona with an ionization parameter and electron density ranging from  $>3000/10^{12}$   $\text{cm}^{-3}$  near the accretor to  $\sim 30/10^{11}$   $\text{cm}^{-3}$  near the outer boundary would scatter about 1% of the irradiated X-ray emission. This suggests that the intrinsic X-ray luminosity of CAL 87 is of order a few times  $10^{37}$  erg/s.

Though the earlier SSS classification was based on a wrong interpretation of the X-ray spectrum, the new kT and L estimates from our LETG spectrum re-confirm the supersoft source nature of CAL 87.

JG and RD gratefully acknowledge the *Chandra* Award No. GO1-2022X.

### REFERENCES

- Asai, K., Dotani, T., Nagase, F., Ebisawa, K., Mukai, K. et al. 1998, *ApJ* 503, L143  
 Callanan P.J., Machin G., Naylor T., Charles P.A. 1989, *MNRAS* 241, 37p  
 Cowley A.P., Schmidtke P.C., Crampton D., Hutchings J.B. 1990, *ApJ* 350, 288  
 Dotani T., Asai K., Ebisawa K., Greiner J., 2000, *COSPAR*, *Adv. Space Res.* 25, No. 3/4, 375  
 Ebisawa K., Mukai K., Kotani T., et al. 2001, *ApJ* 550, 1007  
 Long K.S., Helfand D.J., Grabelsky D.A. 1981, *ApJ* 248, 925  
 Pakull M.W., Beuermann K., van der Klis M., van Paradijs J., 1988, *A&A* 203, L27  
 Parmar A.N., Kahabka P., Hartmann H.W. et al., 1997, *A&A* 323, L33  
 Schandl S., Meyer-Hofmeister E., Meyer F., 1997, *A&A* 318, 73  
 Schmidtke P.C., McGrath T.K., Cowley A.P., Frattare L.M. 1993, *PASP* 105, 863

Vadim Burwitz, Jochen Greiner and Anatoli Iyudin: Max-Planck-Institut für extraterrestrische Physik, Giessenbachstraße, 85748 Garching, Germany (burwitz,jcg,ani@mpe.mpg.de).

Rosanne DiStefano: Center for Astrophysics, Cambridge, MA 02138. U.S.A. (rd@cfa.harvard.edu).

Mario Jimenez-Garate and Norbert Schulz: MIT, Cambridge, U.S.A. (mario,nss@space.mit.edu).

Robert Schwarz: Universität Göttingen, 37083 Göttingen, Germany, (rsc@uni-sw.gwdg.de).