INSTRUCTION TO CONTRIBUTORS

The Journal of Helminthology publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance, but only in exceptional circumstances will systematic or taxonomic studies be acceptable.

Manuscripts, which must be in English or French (with an English summary) should be accompanied by a letter signed by all the authors and should be addressed to:

The Editor, Journal of Helminthology, London School of Hygiene and Tropical Medicine, Winches Farm Field Station, 395 Hatfield Road, St Albans, Herts AL4 0XQ, England.

Two copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief Introduction; Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Research Notes should also be preceded by a brief abstract. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration will be printed is 12.0×20.0 cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space, should be indicated. Some indication of scale (preferably a scale bar) should normally be given on the figure. Photocopies of illustrations should be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints of the same size as they are to appear in the Journal (maximum size 12.0×20.0 cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

DUKE, B. O. L. (1971) The ecology of onchocerciasis in man and animals. In: *Ecology and Physiology of parasites* (editor, A. M. Fallis) pp. 213–222. Adam Hilger Ltd.: London.

JAMES, C. & WEBBE, G. (1973) A comparison of Egyptian and East African strains of Schistosoma haematobium. Journal of Helminthology, 47, 49-59.

25 offprints are provided free of charge; additional copies may be ordered at the proof stage.

Contents

	Pages
Litomosoides carinii: retardation of worm growth and of migration of challenge infections in jirds (Meriones unguiculatus). D. J. WEINER and B. Y. LEEDS	191–194
Immunological evidence of chemotherapeutic action of mebendazole against <i>Ancylostoma ceylanicum</i> (Looss, 1911) in hamsters (<i>Mesocricetus auratus</i>). V. R. KAMATH, M. K. BHOPALE and M. B. BHIDE	195–199
Retinol deficiency and <i>Dipetalonema viteae</i> infection in the hamster. D. STÜRCHLER, A. HANCK, H. WEISER, U. MANZ and N. WEISS	201–210
Anthelmintic effects of bithionol, paromomycin sulphate, flubendazole and mebendazole on mature and immature <i>Hymenolepis nana</i> in mice. J. MAKI and T. YANAGISAWA	211–216
The influence of temperature and osmotic stress on the development and eclosion of hookworm eggs. B. E. MATTHEWS	217–224
The cercarial emergence of <i>Schistosoma japonicum</i> from <i>Oncomelania quadrasi</i> under outdoor conditions in the Philippines. K. KAWASHIMA, B. L. BLAS and A. T. SANTOS	225–231
Prolonged explusion of adult <i>Trichinella spiralis</i> and eosinophil infiltration in mast cell-deficient W/WV mice. M. KAMIYA, Y. OKU, H. ITAYAMA and M. OHBAYASHI	233–239
A new filarial nematode (Onchocercidae) from warthogs (<i>Phacochoerus aethiopicus</i>) of the Kruger National Park. J. R. PALMIERI, J. M. PLETCHER, V. DE VOS and J. BOOMKER	241–245
Infection of dogs with <i>Echinococcus granulosus</i> protoscolices derived from serial passages in mice. (Research Note). P. H. DE RYCKE, O. VANPARIJS, E. PENNOIT-DE COOMAN and R. DECEUNYNCK	247–249
The suppression of homologous immunity by soluble adult antigens of <i>Nematospiroides dubius</i> . D. I. PRITCHARD and J. M. BEHNKE	251–256
Efficacy of ivermectin against <i>Syphacia obvelata</i> (Nematoda) in mice. D. A. OSTLIND, M. A. NARTOWICZ and W. G. MICKLE	257–261
5'-Nucleotidase activity and substrate affinity in digenetic trematodes. (Research Note) A. A. SIDDIQUI and W. A. NIZAMI	263–266
Observations on Toxocara pteropodis infections in mice. P. PROCIV	267–275
The domestic cat as a host for Brugian filariasis in South Kalimantan (Borneo), Indonesia. J. R. PALMIERI, S. MASBAR, PURNOMO, H. A. MARWOTO, S. TIRTOKUSUMO and F. DARWIS	277–281
Variable infectivity of third-stage larvae of <i>Brugia malayi</i> . M. L. EBERHARD and R. C. LOWRIE, Jr.	283–286
Book Review	276
Announcements 23	32, 240, 286