

LETTER TO THE EDITOR

doi:10.1017/S1041610220001428

Air pollution and geriatric mental health: perspectives on the COVID-19 pandemic

In the absence of any vaccine or effective pharmaceutical treatment options for COVID-19, the likelihood of long-term necessity of social restrictions and confinement practices like isolation and quarantine among older adults—the most visible victims of COVID-19—remains high (CBC News, 2020; *The Washington Post*, 2020). Such practices, however, contribute substantially to psychological distress, depression, anxiety, mood disorders, insomnia, stigmatization, reduced self-esteem and self-control, and post-traumatic stress disorder among those confined, and are more pronounced amongst the geriatric population (Holmes *et al.*, 2020; Hossain *et al.*, 2020). This necessitates insight and action into addressing various risk factors that may aggravate such adverse mental health consequences among older adults as they continue to self-isolate amid the COVID-19 pandemic (Holmes *et al.*, 2020).

A growing body of research has linked air pollution, both at ambient and household levels, with an array of mental health disorders such as anxiety, stress, depression, mood disorders, suicidal behavior, cognitive impairment, and dementia among older adults (Braithwaite *et al.*, 2019; Oudin *et al.*, 2018). While ambient air pollution levels have considerably plummeted globally due to reduction in outdoor anthropological activities during the pandemic, the levels of household air pollution (HAP) can be expected to have exponentiated as the majority of the populations are spending greater time indoors and engaging in activities such as cooking, heating, and lighting in greater proportions than the pre-COVID-19 era (Venter *et al.*, 2020). This is especially of concern as more than 3 billion people across the globe continue to rely on highly polluting solid fuels (wood, dung, agricultural residues, coal, charcoal) to meet their household energy demands (World Health Organization, 2018).

Experimental evidence suggests that air pollutants, particularly fine particulate matter (PM_{2.5}, particulate matter with an aerodynamic diameter less than 2.5- μm)—the most common and major health deterrent combustion product of solid fuels, can easily translocate to the central nervous system through the olfactory-neural pathway and the

cardio-respiratory system to induce neuroinflammation, oxidative stress, microglial activation, cerebrovascular dysfunction, and alterations in the blood–brain barrier (Genc *et al.*, 2012). Among epidemiological investigations, Liu and colleagues, by employing the propensity score matching method, compared the effects of HAP exposure on depression among solid and clean fuel (liquefied gas, natural gas, and electricity) users, using a nationally representative dataset of the older population in China. They found a significantly higher Center for Epidemiologic Studies Depression Scale (CES-D) score and depression among solid fuel users than clean fuel users; with a CES-D score of 0.59 (95% confidence interval [CI]: 0.31, 0.89) and the depression risk odds ratio of 1.26 (95% CI: 1.14, 1.41) among solid fuel users (Liu *et al.*, 2020). In another study, based out of Northern Sweden, Oudin and colleagues assessed the longitudinal association between PM_{2.5} emissions from local residential wood burning and dementia. They found a significant association between the PM_{2.5} exposure from woodsmoke and dementia incidence among older adults, with a hazard ratio of 1.55 (95% CI: 1.00, 2.41) for a 1- $\mu\text{g}/\text{m}^3$ increase in PM_{2.5} concentration (Oudin *et al.*, 2018).

Therefore, given the plausibility of exacerbation of adverse mental health outcomes due to HAP exposure among older adults, primary prevention should be enabled at the household level itself. The need for maintaining maximum ventilation at homes, particularly whilst undertaking activities involving solid fuel use, should be mass communicated along with improving awareness around HAP and associated mental health risks among older adults and their caregivers. Further, ensuring affordability and robust door-step-delivery mechanisms for clean fuels and systems for older adults would be critical, considering their limited mobility and financial capacities that have been worsened by the pandemic. This can be challenging, especially in developing countries that are highly populated and where social welfare is suboptimal. Strengthening of psychosocial care by optimizing existing mental health services for older adults, alongside preventing their HAP exposures by considering the drivers and barriers of sustained clean fuel use should be prioritized. This would require the engagement of multidisciplinary stakeholders, as it may help prioritize context-specific strategies to meet the mental health and clean fuel needs of the

geriatric populace. Lastly, from a research perspective, building psychogeriatric evidence in relation to COVID-19 should incorporate modeling for environmental risk factors, including household air quality and trends in fuel usage. Such research efforts should be contextualized by accounting for personal environmental exposures, socio-economic and ethnocultural differences, and the implications of various psychosocial support interventions extended for older adults during the global crisis.

Conflict of interest

We declare no conflicts of interest.

References

- Braithwaite, I. et al.** (2019) Air pollution (Particulate matter) exposure and associations with depression, anxiety, bipolar, psychosis and suicide risk: a systematic review and meta-analysis. *Environmental Health Perspectives*, 127(12), 126002. doi: [10.1289/EHP4595](https://doi.org/10.1289/EHP4595).
- CBC News** (2020) As Ontario gets set to reopen, self-isolation continues for millions of citizens, 18 May. Available at: <https://www.cbc.ca/news/canada/ottawa/ontario-isolation-seniors-covid-coronavirus-1.5572184>
- Genc, S. et al.** (2012) The adverse effects of air pollution on the nervous system. *Journal of Toxicology*, 2012(782462). doi: [10.1155/2012/782462](https://doi.org/10.1155/2012/782462).
- Holmes, E.A. et al.** (2020). Multidisciplinary research priorities for the COVID-19 pandemic: a call for action for mental health science. *The Lancet Psychiatry*, 7, 547–560. doi: [10.1016/S2215-0366\(20\)30168-1](https://doi.org/10.1016/S2215-0366(20)30168-1).
- Hossain, M.M., Sultana, A. and Purohit, N.** (2020). Mental health outcomes of quarantine and isolation for infection prevention: a systematic umbrella review of the global evidence. *Epidemiology and Health*. doi: [10.4178/epih.e2020038](https://doi.org/10.4178/epih.e2020038).
- Liu, Y., Chen, X. and Yan, Z.** (2020) Depression in the house: the effects of household air pollution from solid fuel use among the middle-aged and older population in China. *Science of the Total Environment*, 703(134706). doi: [10.1016/j.scitotenv.2019.134706](https://doi.org/10.1016/j.scitotenv.2019.134706).
- Oudin, A. et al.** (2018) Association between air pollution from residential wood burning and dementia incidence in a longitudinal study in Northern Sweden. *PLoS ONE*, 13(6), e0198283. doi: [10.1371/journal.pone.0198283](https://doi.org/10.1371/journal.pone.0198283).
- The Washington Post** (2020) Europe is starting to lift coronavirus restrictions. But might seniors be left out? 2 May. Available at: https://www.washingtonpost.com/world/europe/germany-eases-restrictions-coronavirus-old/2020/05/01/77ffa5a4-857c-11ea-81a3-9690c9881111_story.html
- Venter, Z. S. et al.** (2020). COVID-19 lockdowns cause global air pollution declines with implications for public health risk. *medRxiv*. doi: [10.1101/2020.04.10.20060673](https://doi.org/10.1101/2020.04.10.20060673).
- World Health Organization** (2018) Household air pollution and health. Available at: <https://www.who.int/en/news-room/fact-sheets/detail/household-air-pollution-and-health>; accessed 19 June 2020.
- RACHIT SHARMA,¹**  **MD MAHBUB HOSSAIN,²**
PRIYANKA PAWAR¹ AND **SONAM SHARMA³**
- ¹GRID Council, Delhi-NCR 201301, India
²Department of Health Promotion and Community Health Sciences, School of Public Health, Texas A&M University, TX 77843, USA
³Department of Pathology, Kalpana Chawla Government Medical College, Karnal, Haryana 132001, India
- Correspondence should be addressed to:* Rachit Sharma, 11-A, Building 9, The HIBISCUS, Sector 50, Gurugram, Haryana 122018, India. Phone: (+91) 8199021326; Fax: (+91) 124 227 8151. Email: drrachitsharma09@gmail.com