

Letters to the Editor

Preoperative Prophylactic Antibiotics: Brief Historical Note

To the Editor:

It is now well established that antibiotics administered perioperatively can effectively prevent incisional wound infections. Original data in animals suggested that the antibiotics should be administered prior to the time of the initial incision and that if administered four or more hours afterward there was a complete loss of protection.^{1,2} Recently, an observational study showed that administration of the preoperative antibiotic was most effective if given in the "window" period of 0 to 2 hours before the incision is made.³ The rate of infection increased on either side of the optimal window: administering the prophylactic antibiotic more than two hours before or at longer times after the incision progressively increased incisional wound infection rates.

Accompanying the New *England Journal of Medicine* article was an editorial that I wrote attempting to put the observational study in context.⁴ I subsequently received three references and correspondence from Gabriel P. Seley, MD, which he said "...will cover all the facts...[to help document]...that I was the one who first suggested preoperative antibiotic use in surgery and used it with good results."

Dr. Seley may in fact be right to claim priority, and I throw the historical gauntlet down for those who think otherwise. In 1939, "The

Use of Sulfanilamide in Surgery of the Colon and Rectum: Preliminary Report," by John H. Garlock and Gabriel P. Seley was published in *Surgery*.⁵ The work was conducted primarily by Dr. Seley at the Mt. Sinai Hospital in New York under the direction of Dr. Gregory Schwartzman. Bacterial cultures were taken of the retroperitoneal and pericolonic tissue of the bowel wall and from the mucosal surface of the neoplasm. Hemolytic streptococci and *Escherichia coli* were found commonly in untreated patients.

In a study with historical controls, Dr. Seley then administered sulfanilamide for three days preoperatively (15 grains every 4 hours). Two of 21 patients died, one major infection occurred, and some patients convalesced smoothly despite "gross spilling." Importantly, no *Streptococcus hemolyticus* was recovered in any operative cultures of the 21 cases. Garlock and Seley concluded cautiously, however, that "the number of cases reported herewith is not sufficient to warrant definite conclusions."

In a subsequent report in 1941, Seley and Colp extended their practice of perioperative antibiotics for colon surgery to gastric surgery.⁶ Two years later, in 1943, there was a proceedings document in which it was specifically reported that there was a "striking diminution of the numbers of streptococci and *B welchii* in cases treated preoperatively with sulfanilamide...The monthly rate from peritonitis in 123 colon and rectal operations thus treated was 4% against 10% mortality reported in the literature."⁷

Dr. Seley surely did not perform a randomized controlled clinical trial but appears to have had the idea of preoperative prophylactic antibiotics for colon surgery more than 50 years ago, only a few years after sulfanilamide was discovered. Moreover, his idea was scientifically based on the best microbiological evidence available and his observations carefully stated. At this point, it would seem reasonable that we recognize a pioneer of microbiology and surgery, Dr. Gabriel P. Seley.

REFERENCES

1. Burke JF. The effective period of preventive antibiotic action in experimental incisions and dermal lesions. *Surgery*. 1991;50:161-168.
2. Bernard HR, Cole WR. The prophylaxis of surgical infection: the effect of prophylactic antimicrobial drugs on the incidence of infection following potentially contaminated operations. *Surgery* 1964;56:151-157.
3. Classen DC, Evans RS, Pestotnik SL, Horn SD, Menlove RL, Burke JP. The timing of prophylactic administration of antibiotics and the risk of surgical-wound infection. *N Engl J Med*. 1992;326:281-286.
4. Wenzel P. Preoperative antibiotic prophylaxis. *N Engl J Med*. 1992;326:337-339.
5. Garlock JH, Seley GP. The use of sulfanilamide in surgery of the colon and rectum. Preliminary report. *Surgery*. 1939;5:787-790.
6. Seley GP, Colp R. The bacteriology of peptic ulcers and gastric malignancies: possible bearing on complications following gastric surgery. *Surgery*. 1941;10:369-380.
7. Seley GE. Surgical bacteriology. In: *Proceedings of Seminars on Recent Advances in Bacteriology and Immunology with Clinical Considerations*. Reprinted from *Journal of Mount Sinai Hospital*. 1943;9:923-924.

Richard P. Wenzel, MD, MSc
University of Iowa
Iowa City, Iowa


Submission of Abstracts
Deadline: January 8, 1993
Late Breakers before March 5

Submission of Abstracts
Deadline: January 8, 1993
Late Breakers before March 5

The Third Annual Meeting of

SHEA

The Society for Hospital Epidemiology of America



April 18-20, 1993

The Palmer House Hotel, Chicago, Illinois

15 hours of Category I/AMA Credit

**For Preliminary Program and Registration Information,
please contact SHEA at 609-845-1 720 or fax 609-853-0411**

University of Illinois College of Medicine is accredited by the Accreditation Council for CME to sponsor continuing medical education for physicians. University of Illinois College of Medicine designates this continuing medical education activity for 15 credit hours in Category 1 of Physicians Recognition Award of the American Medical Association.

VACUTAINER Blood Collection Needles

CAUTION

Biologic samples and blood collection "sharps" (lancets, needles, and blood collection sets) require special handling and procedures of your facility. Seek immediate medical attention in the event of any exposure to biologic samples (for example, those that may transmit HBV (HEPATITIS), HIV (AIDS), or other infectious diseases). Always use the built-in needle protector, if the blood collection device provides one. Becton Dickinson VACUTAINER needles, but the policies and procedures of your facility may differ and should always be followed. Dispose of blood collection "sharps" in biohazard containers approved for their disposal.

Dickinson VACUTAINER Systems
Franklin Lakes, NJ 07417-1885
VACUTAINER is a trademark of Becton Dickinson and Company.

Made in U.S.A.

WE'RE BRINGING SAFETY TO THE SURFACE

As the world's leading manufacturer of sharps products, Becton Dickinson is committed to helping make today's healthcare workplace safer. Developing products that comply with current regulations and setting new safety standards is a large part of that commitment, but an equally important factor is promoting correct and conscientious usage of all sharps products. As a step toward accomplishing this goal, Becton Dickinson VACUTAINER Systems is now including special safety messages on VACUTAINER Brand Needle packaging as an extra reminder to "handle with care."

Promoting adherence to recommended procedures is one important way to prevent dangerous needlestick injuries and exposure to hazardous biologic specimens—and we at Becton Dickinson VACUTAINER Systems take this responsibility seriously.

For detailed information on all of our safety products, contact your Becton Dickinson VACUTAINER Systems Sales Representative.



BECTON DICKINSON

Becton Dickinson VACUTAINER Systems, Franklin Lakes, New Jersey 07417
PRECISIONGLIDE and VACUTAINER are registered trademarks of Becton Dickinson and Company.
© 1993