

## Micronutrient and anthropometric status indicators are associated with physical fitness in Colombian schoolchildren

Joanne E. Arsenault<sup>1</sup>, Mercedes Mora-Plazas<sup>2</sup>, Yibby Forero<sup>3</sup>, Sandra Lopez-Arana<sup>2</sup>, Germán Jáuregui<sup>4</sup>, Ana Baylin<sup>5</sup>, Paul M. Gordon<sup>6</sup> and Eduardo Villamor<sup>1,5,7\*</sup>

<sup>1</sup>Department of Nutrition, Harvard School of Public Health, Boston, MA, USA

<sup>2</sup>Department of Nutrition, National University of Colombia, Bogotá, Colombia

<sup>3</sup>The Nutrition Group, National Institute of Health, Bogotá, Colombia

<sup>4</sup>Department of Physical Education, INCCA University of Colombia, Bogotá, Colombia

<sup>5</sup>Department of Epidemiology, University of Michigan School of Public Health, Ann Arbor, MI, USA

<sup>6</sup>Laboratory for Physical Activity and Exercise Intervention Research, University of Michigan, Ann Arbor, MI, USA

<sup>7</sup>Department of Environmental Health Sciences, University of Michigan School of Public Health, 1420 Washington Heights, SPH II, Room M6017, Ann Arbor, MI 48109, USA

(Received 19 July 2010 – Revised 7 December 2010 – Accepted 7 December 2010 – First published online 1 February 2011)

### Abstract

Poor physical fitness is associated with increased health-related risks in children. The association of nutritional status indicators and physical fitness in children residing in developing countries is not well characterised. We conducted a cross-sectional study among 1945 children of age 5–12 years in Bogotá, Colombia, to assess whether anthropometric and micronutrient status indicators were associated with performance in the shuttle run and standing long jump tests. Stunted children scored significantly lower in the run (0.4 s;  $P=0.0002$ ) and jump (6 cm; boys only;  $P=0.003$ ) tests than non-stunted children, after adjustment for age and other factors. Children who were thin, overweight or obese ran slower than normal-weight children ( $P<0.01$ ). Lower jump scores were associated with overweight or obesity and greater arm fat area in boys only ( $P<0.0001$ ). Girls with low ferritin concentrations ran 0.6 s slower than girls with normal ferritin concentrations ( $P=0.02$ ). Erythrocyte folate concentrations were linearly related to higher run ( $P<0.0001$ ) and long jump scores ( $P=0.0001$ ). Boys with marginal or low vitamin B<sub>12</sub> status had 4 cm lower long jump scores than children with normal status ( $P=0.01$ ). Suboptimal anthropometric and micronutrient status are related to poorer performance in fitness tests. The effects of improving nutritional status on physical fitness of children warrant investigation.

**Key words:** Anaerobic performance; Micronutrients; Stunting; Overweight

Physical fitness is associated with decreased risks of high blood pressure, dyslipidaemia and insulin resistance in children<sup>(1)</sup>, which in turn result in lowered risk of CVD and metabolic disorders during adulthood. Physical fitness during adolescence has also been related to a healthier cardiovascular profile in adulthood<sup>(2)</sup>.

National standardised physical fitness tests for children are conducted in many countries<sup>(3)</sup>. A battery of tests is often included to evaluate both aerobic and anaerobic performance-related fitness. According to reviews of studies in children worldwide, performance in these tests has declined during the past three decades<sup>(4,5)</sup>. This may be partly attributed to the increased prevalence of overweight and obesity that many countries are experiencing.

In several countries that are undergoing the nutritional transition, the increasing prevalence of overweight and obesity

coexists with substantial rates of malnutrition<sup>(6,7)</sup>. The role that marginal malnutrition could play in performance-related fitness of children has not been studied extensively. Stunted children have short lower extremities that may have a negative impact on performance in run or jump tests, but they may also have low muscle mass or poor micronutrient status that could negatively affect performance. Lower performance scores in fitness tests have been reported among stunted children compared with non-stunted children<sup>(8,9)</sup>. Fe-deficiency anaemia results in reduced performance<sup>(10,11)</sup>, and physical training can decrease body Fe stores<sup>(12)</sup>. However, few studies have examined other micronutrients that are associated with anaemia, such as folate or vitamin B<sub>12</sub>, in relation to performance-related fitness. In adolescents, elevated homocysteine, which is associated with folate and vitamin B<sub>12</sub> deficiencies, was related to reduced cardiovascular fitness<sup>(13)</sup>.

\*Corresponding author: E. Villamor, fax +1 734 936 7283, email villamor@umich.edu

Children receiving a fortified beverage that provided Fe, folate, vitamin B<sub>12</sub> and other micronutrients showed significantly improved fitness compared with children receiving a beverage without micronutrients<sup>(14)</sup>.

We conducted a cross-sectional study to examine the associations between anaerobic performance-related fitness and anthropometric and micronutrient status indicators in schoolchildren from Bogotá, Colombia. Colombia is a country undergoing a nutritional transition, from high rates of undernutrition to increasing rates of overweight. Non-negligible rates of stunting, overweight or obesity and micronutrient deficiencies in this cohort of children have previously been reported<sup>(15–18)</sup>. We hypothesised that children with suboptimal nutritional status (stunting, overweight or low micronutrient status) would have lower scores on anaerobic fitness tests than children with adequate nutritional status.

## Experimental methods

### Study population

The present study is part of a project on children's health and nutritional status in primary public schools of Bogotá, Colombia, which was initiated in 2006. Details of the study design have been reported previously<sup>(19)</sup>. In brief, we randomly selected a representative sample of 3202 primary schoolchildren of age 5–12 years, using a cluster sampling strategy in which all primary public schools in the city were included. The sampling units were the classrooms. Recruitment occurred at the beginning of the school year, in February 2006. The sample is representative of low- and middle-income families from Bogotá, since the public primary school system enrolls the majority of school-age children, and close to 90% of them belong to low- and middle-socioeconomic strata. The present study was conducted according to the guidelines laid down in the Declaration of Helsinki, and all procedures involving human subjects were approved by the Ethics Committee of the National University of Colombia Medical School. The Human Subjects Committee at the Harvard School of Public Health approved the use of data from the study. Written informed consent was obtained from the children's primary care providers before enrolment.

### Field procedures

At the time of enrolment, information was obtained on the parents' sociodemographic characteristics, including age, marital status, education level and indicators of the household socio-economic status, using a self-administered questionnaire that was completed and returned by 2466 households, representing 81% of enrolled children's families. During the 3 weeks following recruitment, teams of trained research assistants visited the schools to obtain a fasting blood specimen by venepuncture from enrolled children and to collect anthropometric measurements using standardised techniques<sup>(20)</sup>. Weight was measured to the nearest 0.1 kg on Tanita HS301 electronic scales (Tanita, Tokyo, Japan), while height was measured to the nearest 1 mm using wall-mounted portable

Seca 202 stadiometers (Seca, Hamburg, Germany). Arm circumference was measured to the nearest 0.1 cm using a non-extensible tape. Triceps skinfolds were measured to the nearest 0.5 mm using Slim Guide Skinfold Calipers (Creative Health Products, Inc., Plymouth, MI, USA).

Also during this period, teams of physical trainers administered two standardised physical fitness tests in a subgroup of 1945 students: the shuttle run and the standing long jump. These tests are part of the 'Eurofit' battery of fitness tests that include endurance, strength, speed, flexibility and balance<sup>(21)</sup>. The shuttle run is a test of speed and agility, and the standing long jump is a test of explosive power and leg strength; both are measures of anaerobic performance-related fitness primarily. For the shuttle run, children were instructed to stand on a line, run a distance of 9 m at maximum speed to a reference point and pick up a block, turn and run back to the starting line, place the block down, and repeat this course a second time (for a total distance of 36 m). The run time was recorded to the nearest 0.01 s using digital stopwatches when the child returned to the starting line and placed the second block down. For the standing long jump, children were instructed to stand on a marked line and jump as far as possible from the standing position, using their arms for balance. The jump distance was recorded to the nearest 1.0 cm. Each test was administered three times, and results from the three attempts were recorded.

### Laboratory methods

Approximately 4 ml of blood were collected in EDTA tubes following a protocol to avoid haemolysis. On the same day of collection, the aliquots were protected from sunlight and transported on ice to the National Institute of Health in Bogotá, where all biochemistry analyses took place. Complete blood counts were obtained, and Hb concentrations were determined using the haemoglobinocyanide method. An aliquot was centrifuged at 1500 g for 15 min, and plasma was separated. Plasma ferritin and vitamin B<sub>12</sub> concentrations were measured using competitive chemiluminescent immunoassay in an ADVIA Centaur analyser (Bayer Diagnostics, Tarrytown, NY, USA). C-reactive protein was measured using a turbidimetric immunoassay on an ACS180 analyser (Bayer Diagnostics). The packed erythrocyte volume was haemolysed by dilution in a hypotonic aqueous solution of 1% ascorbic acid. Erythrocyte folate was measured on the erythrocyte lysates using chemiluminescent immunoassay.

### Data analyses

Outcomes were the shuttle run score (s) and the standing long jump score (cm). We chose the best result from each child's three attempts for analyses. Both outcomes were treated as continuous variables. Henceforth, a 'higher' score refers to a lower run time or a further jump distance.

The main exposures of interest were anthropometric and micronutrient status indicators. Anthropometric indicators included stunting, thinness, overweight, obesity, and arm muscle and fat area. Child stunting was defined as

height-for-age  $< -2$  standard deviations from the sex- and age-specific median of the National Center for Health Statistics/WHO reference population<sup>(22)</sup>. Thinness was defined based on the BMI-for-age and sex cut-off point corresponding to  $< 18.5 \text{ kg/m}^2$  in adults as proposed by Cole *et al.*<sup>(23)</sup>. Overweight or obese was defined according to BMI cut-off points for age and sex corresponding to  $\geq 25 \text{ kg/m}^2$  in adults, following the International Obesity Task Force recommendations<sup>(24)</sup>. Mid-upper arm muscle area (an index of total body muscle mass) and arm fat area (an index of body fat) were calculated using equations that include the arm circumference and triceps skinfold measures<sup>(25)</sup>. Age- and sex-adjusted *z*-scores for the arm muscle and fat areas were calculated using the study population as the reference and then were categorised into quartiles. Indicators of micronutrient status were anaemia, ferritin (an indicator of Fe stores), vitamin B<sub>12</sub> serostatus and erythrocyte folate concentrations. Anaemia was defined as Hb  $< 115 \text{ g/l}$ , after subtracting  $12 \text{ g/l}$  from the individual values as an adjustment for an altitude of  $2500 \text{ m}$ <sup>(26)</sup>. Low ferritin was defined as  $< 15 \text{ } \mu\text{g/l}$  if C-reactive protein  $\leq 10 \text{ mg/l}$  or  $< 30 \text{ } \mu\text{g/l}$  if C-reactive protein  $> 10$ <sup>(27)</sup>. Vitamin B<sub>12</sub> status was defined as normal ( $> 221 \text{ pmol/l}$ ) or marginal or low ( $\leq 221 \text{ pmol/l}$ )<sup>(28)</sup>. Erythrocyte folate concentrations were categorised into quartiles since only two children were below  $305 \text{ nmol/l}$ , a cut-off point suggested by the Institute of Medicine<sup>(29)</sup> to define deficiency.

Child and maternal characteristics were considered as covariates in the analysis. Child characteristics included age, sex and time spent playing outdoors as a measure of habitual physical activity. Maternal characteristics included age, education level and parity. Household socio-economic indices included the daily income per capita (household income divided by the household size) and the household socio-economic stratum according to the city's classification of the neighbourhoods' public service fees.

We first compared the distribution of run and jump scores by categories of each sociodemographic characteristic using univariate linear regression models, in which the physical performance test score was introduced as the outcome and each characteristic as the predictor. For ordinal predictors, a test for trend was conducted by introducing a variable representing the ordinal categories of the predictor in the linear model as a continuous covariate. In order to identify socio-demographic covariates that were independently associated with fitness scores and thus could potentially confound their associations with nutritional status indicators, we constructed a multivariate model including the variables that were significantly associated with each test at  $P < 0.10$  in univariate analyses. Predictors were retained in the model if they remained statistically significant at  $P < 0.05$  or were considered to be relevant from a mechanistic viewpoint.

We then examined the associations between each of the anthropometric or micronutrient status indicators and the physical performance tests. Based on established sex differences in physical performance<sup>(8,9)</sup>, which we verified with the present results, the analyses were stratified by sex. Differences in test scores and 95% CI were calculated between

categories of each nutritional predictor using multivariate linear models, which included significant sociodemographic correlates of each test as adjustment covariates. Micronutrient status indicators were included simultaneously into the same model to obtain estimates that were independent of each other. Finally, interactions were assessed among micronutrients on test scores by introducing into the models cross-product terms between the indicators and testing them with the likelihood-ratio test. CI were constructed with robust estimates of the variance, which do not require normality assumptions. In all models, an exchangeable correlation matrix (PROC GENMOD, SAS Institute, Cary, NC, USA) was specified to account for potential correlations within households among siblings. The effect of clustering from the sampling strategy was also considered in the models, but it was negligible and excluded for parsimony. All tests were double-sided and considered to be statistically significant if  $P \leq 0.05$ . Analyses were performed using the Statistical Analysis System (release 9.1; SAS Institute, Cary, NC, USA).

## Results

Among the 1945 children in whom fitness tests were conducted, shuttle run and long jump scores were available in 1894 and 1939, respectively. Some of the tests could not be administered in all schools for logistical reasons. On average, children in whom tests were performed were 0.3 years older ( $P < 0.0001$ ), played outside 0.9 h/week less ( $P = 0.01$ ) and had mothers who were 1 year older ( $P = 0.0005$ ) than children in whom performance tests were not done. There were no differences regarding sex or socio-economic factors. The mean age of the children was 8.9 years. Each of the test scores was positively associated with age ( $P$  for trend  $< 0.0001$ ). Boys ran faster ( $P < 0.0001$ ) and jumped further ( $P < 0.0001$ ) than girls (Table 1). Performance in both tests was positively associated with indicators of socio-economic status. In addition, jump scores were positively associated with time spent playing outdoors ( $P$  for trend = 0.02). The scores from the two tests were positively correlated with each other, independent of age and sex (partial  $r = 0.36$ ,  $P < 0.0001$ ; coefficient term is negative because a lower run time is a 'higher' score).

Stunted children of both the sexes ran significantly slower than non-stunted children (0.4 s;  $P = 0.0002$ ), after adjustment for age, socio-economic status, BMI and arm muscle area (Table 2). Stunted boys scored 6 cm lower in the standing long jump test than non-stunted boys ( $P = 0.003$ ; Table 3). Children who were either thin, overweight or obese ran 0.3 s slower than normal-weight children ( $P < 0.01$ ). Thin girls scored significantly lower in the jump test than normal-weight children of the same sex ( $P = 0.01$ ). Arm fat area was negatively associated with run performance in both girls ( $P$  for trend = 0.002) and boys ( $P$  for trend = 0.0004). Overweight or obesity was negatively associated with jump scores in boys ( $P < 0.0001$ ). Boys in the highest quartile of arm fat area had lower jump scores than those in the lowest quartile ( $P < 0.0001$ ).

**Table 1.** Sociodemographic correlates of shuttle run and standing long jump scores in schoolchildren from Bogotá, Colombia (Mean values, standard deviations, number of children and 95 % confidence intervals)

	Shuttle run time (s)					Standing long jump distance (cm)				
	<i>n</i>	Unadjusted mean	SD	Adjusted differences*	95 % CI	<i>n</i>	Unadjusted mean	SD	Adjusted differences*	95 % CI
Overall	1894	12.75	1.62	–	–	1939	123	24	–	–
Age of child (1-year change)	1870	–0.54	0.02†	–0.54	–0.58, –0.50	1915	8	0.3†	8	7, 8
Sex of child										
Girls	978	13.14	1.66	Ref	–	1008	116	23	Ref	–
Boys	916	12.33	1.46	–0.75	–0.86, –0.64	931	130	22	12	11, 14
Hours played outdoors last week										
Q1 (<1.5)	341	12.94	1.59	–	–	348	119	24	Ref	–
Q2 (1.5–4.49)	353	12.76	1.48	–	–	363	122	22	1	–1, 4
Q3 (4.5–9.99)	358	12.73	1.64	–	–	363	124	24	3	0, 6
Q4 (≥10)	352	12.60	1.71	–	–	361	125	25	3	0, 6
<i>P</i> for trend		0.009		–	–		0.0004		0.02	
Maternal education (years)										
Incomplete primary (1–4)	129	12.53	1.53	–	–	133	125	22	–	–
Complete primary (5)	321	12.70	1.71	–	–	331	123	22	–	–
Incomplete secondary (6–10)	427	12.59	1.52	–	–	438	125	24	–	–
Complete secondary (11)	704	12.83	1.61	–	–	718	122	23	–	–
University (≥12)	110	12.92	1.53	–	–	111	120	27	–	–
<i>P</i> for trend		0.02		–	–		0.09		–	–
Mother's parity										
1	188	13.08	1.82	–	–	193	117	24	–	–
2	611	12.77	1.60	–	–	626	123	23	–	–
3	520	12.65	1.53	–	–	531	125	24	–	–
4	211	12.60	1.64	–	–	213	123	23	–	–
5	151	12.51	1.51	–	–	157	124	24	–	–
<i>P</i> for trend		0.0007		–	–		0.01		–	–
Household income/person per d										
Q1 (median 1880 pesos)	348	12.75	1.64	–	–	362	121	24	Ref	–
Q2 (median 3289 pesos)	392	12.73	1.63	–	–	397	123	22	2	0, 5
Q3 (median 4386 pesos)	372	12.69	1.60	–	–	381	122	23	2	0, 5
Q4 (median 6579 pesos)	378	12.76	1.59	–	–	386	124	25	4	1, 6
<i>P</i> for trend		0.99		–	–		0.09		0.01	
Household socio-economic stratum										
1 (lowest)	158	12.89	1.76	Ref	–	162	119	27	Ref	–
2	634	12.75	1.71	–0.03	–0.25, 0.19	666	124	23	3	0, 7
3	875	12.67	1.47	–0.40	–0.61, –0.19	881	123	23	5	2, 8
4	48	12.65	1.32	–0.32	–0.70, 0.05	47	123	25	4	–1, 10
<i>P</i> for trend		0.07		<0.0001			0.69		0.003	

Q, quartile; Ref, reference.

\*Estimates are from a multivariate linear generalised estimating equation model. Only estimates for variables retained in the final model are presented. Note: Negative estimates for the shuttle run time correspond to 'faster' run times and are considered as 'higher' scores.

† Standard error for a 1-year change in age.

Nutritional status and physical fitness

**Table 2.** Anthropometric correlates of shuttle run scores in schoolchildren from Bogotá, Colombia (Mean values, standard deviations, number of children and 95 % confidence intervals)

	Shuttle run time (s)									
	Girls					Boys				
	<i>n</i>	Unadjusted mean	SD	Adjusted differences	95% CI*	<i>n</i>	Unadjusted mean	SD	Adjusted differences	95% CI*
Child is stunted†										
No	889	13.09	1.61		Ref	817	12.29	1.43		Ref
Yes	76	13.60	2.10	0.56	0.20, 0.92	78	12.58	1.66	0.32	0.03, 0.61
BMI-for-age status‡ (kg/m <sup>2</sup> )										
Thin	107	13.09	1.50	0.30	0.03, 0.57	91	12.45	1.32	0.26	0.00, 0.52
Normal	769	13.09	1.67		Ref	705	12.27	1.47		Ref
Overweight or obese	88	13.55	1.71	0.33	0.02, 0.65	98	12.56	1.45	0.33	0.03, 0.63
MUAMA z-score quartiles§										
Q1	194	12.94	1.39		Ref	198	12.25	1.32		Ref
Q2	217	13.04	1.49	0.04	-0.18, 0.25	182	12.15	1.43	-0.11	-0.32, 0.10
Q3	200	12.98	1.57	0.07	-0.16, 0.30	193	12.16	1.16	-0.13	-0.34, 0.08
Q4	202	13.16	1.57	0.25	-0.02, 0.52	190	12.52	1.54	0.08	-0.19, 0.36
<i>P</i> for trend		0.21			0.25		0.09			0.84
MUAFA z-score quartiles§										
Q1	214	12.97	1.40		Ref	182	12.14	1.13		Ref
Q2	183	12.96	1.61	-0.01	-0.23, 0.21	215	12.14	1.34	0.08	-0.11, 0.27
Q3	201	12.93	1.43	0.06	-0.15, 0.28	186	12.27	1.48	0.21	-0.02, 0.44
Q4	215	13.24	1.57	0.40	0.17, 0.64	180	12.56	1.48	0.50	0.23, 0.78
<i>P</i> for trend		0.09			0.002		0.002			0.0004

Ref, reference; MUAMA, mid-upper arm muscle area; MUAFA, mid-upper arm fat area; HAZ, height-for-age.

\* Estimates are from linear generalised estimating equation models with covariates that included age and household socio-economic stratum. Other covariates included in models with each anthropometric status indicator were as follows: BMI status and MUAMA z-score in the model with 'child is stunted'; HAZ and MUAMA z-score in the model with 'BMI-for-age status'; HAZ and MUAFA z-score in the model with 'MUAMA z-score quartiles'; HAZ and MUAMA z-score in the model with 'MUAFA z-score quartiles'. Note: negative estimates for the shuttle run time correspond to 'faster' run times and are considered as 'higher' scores.

† Stunted is defined as HAZ < -2 from the sex- and age-specific median of the National Center for Health Statistics/WHO reference population<sup>(22)</sup>.

‡ Thinness is defined as BMI-for-age and sex below the cut-off point corresponding to < 18.5 kg/m<sup>2</sup> in adults<sup>(23)</sup>; overweight and obese are defined according to BMI cut-off points for age and sex corresponding to ≥ 25 kg/m<sup>2</sup> in adults<sup>(24)</sup>.

§ MUAMA and MUAFA were calculated based on arm circumference and triceps skinfolds and standardised for age and sex using the study population.

**Table 3.** Anthropometric correlates of standing long jump scores in schoolchildren from Bogotá, Colombia (Mean values, standard deviations, number of children and 95 % confidence intervals)

	Standing long jump distance (cm)									
	Girls					Boys				
	<i>n</i>	Unadjusted mean	SD	Adjusted differences	95 % CI*	<i>n</i>	Unadjusted mean	SD	Adjusted differences	95 % CI*
Child is stunted†										
No	913	117	23	Ref		831	131	22	Ref	
Yes	82	113	23	2	−31, 35	79	124	23	−6	−10, −2
BMI-for-age status‡ (kg/m <sup>2</sup> )										
Thin	110	119	19	−24	−42, −6	90	133	23	3	−1, 6
Normal	790	117	24	Ref		719	131	22	Ref	
Overweight or obese	93	109	19	−33	−80, 14	100	124	20	−11	−14, −7
MUAMA z-score quartiles§										
Q1	204	116	22	Ref		200	129	21	Ref	
Q2	220	115	22	−3	−23, 18	185	129	21	0	−3, 3
Q3	209	118	21	2	−19, 23	198	129	22	1	−2, 5
Q4	210	116	23	3	−20, 27	195	129	23	4	0, 7
<i>P</i> for trend		0.91		0.96			0.90		0.73	
MUAFA z-score quartiles§										
Q1	221	117	21	Ref		184	132	21	Ref	
Q2	188	118	22	11	−6, 28	218	131	22	−2	−5, 2
Q3	213	117	22	26	−1, 52	192	129	20	−5	−8, −1
Q4	221	113	22	8	−7, 22	184	123	22	−13	−17, −9
<i>P</i> for trend		0.02		0.84			<0.0001		0.22	

Ref, reference; MUAMA, mid-upper arm muscle area; MUAFA, mid-upper arm fat area; HAZ, height-for-age.

\* Estimates are from linear generalised estimating equation models with covariates that included age, household socio-economic stratum, hours spent playing outdoors per week and household per capita income. Other covariates included in models with each anthropometric status indicator were as follows: BMI status and MUAMA z-score in the model with 'child is stunted'; HAZ and MUAMA z-score in the model with 'BMI-for-age status'; HAZ and MUAFA z-score in the model with 'MUAMA z-score quartiles'; HAZ and MUAMA z-score in the model with 'MUAFA z-score quartiles'.

† Stunted is defined as HAZ < −2 from the sex- and age-specific median of the National Center for Health Statistics/WHO reference population<sup>(22)</sup>.

‡ Thinness is defined as BMI-for-age and sex below the cut-off point corresponding to <18.5 kg/m<sup>2</sup> in adults<sup>(23)</sup>; overweight and obese are defined according to BMI cut-off points for age and sex corresponding to ≥25 kg/m<sup>2</sup> in adults<sup>(24)</sup>.

§ MUAMA and MUAFA were calculated based on arm circumference and triceps skinfolds and standardised for age and sex using the study population.



Anaemic children did not differ significantly from non-anaemic children in test scores (Table 4). Girls with low ferritin, an indicator of Fe stores, scored significantly lower in the run test than girls with normal ferritin (0.6 s;  $P=0.02$ ), but in boys the difference did not reach significance (0.5 s;  $P=0.06$ ). Boys with low ferritin had 7 cm lower jump scores than boys with normal ferritin ( $P=0.03$ ; Table 5). Erythrocyte folate concentrations were significantly, independently and linearly related to higher run scores in both girls ( $P$  for trend=0.0005) and boys ( $P$  for trend=0.009). Erythrocyte folate concentrations were also linearly associated with jump scores in both girls ( $P$  for trend=0.01) and boys ( $P$  for trend=0.003). Children in the highest quartile of folate jumped 4 cm further than children in the lowest quartile of the same sex (females,  $P=0.02$ ; males,  $P=0.007$ ). Boys who had marginal or low vitamin B<sub>12</sub> status had 4 cm lower jump scores ( $P=0.01$ ) than boys with normal B<sub>12</sub> status.

There were no significant interactions between micronutrients for any of the outcomes among all the children; there were too few children in some of the categories to test for interactions by sex. In supplemental analysis, we examined whether the associations between stunting and the test scores were confounded by micronutrient status, given that micronutrient deficiencies could cause linear growth retardation. After controlling for micronutrient concentrations, the associations between stunting and fitness scores remained unchanged. In addition, there were no interactions between stunting and the micronutrients.

## Discussion

In the present study, we examined cross-sectional associations of anthropometric indices and biomarkers of micronutrient status with performance-related fitness indicators in a large representative sample of low- and middle-income schoolchildren from Bogotá, Colombia. We found negative associations between stunting, overweight or obesity, and arm fat area and anaerobic performance. With respect to micronutrients, there were significant trends in higher run and jump scores with higher erythrocyte folate concentrations, despite negligible folate deficiency. In male children, marginal or low vitamin B<sub>12</sub> status was associated with lower long jump scores than children with normal status.

Our finding of a positive association between folate status and anaerobic performance-related fitness is novel. The few studies that have been reported on folate status and exercise performance have focused on aerobic performance. Higher serum folate concentrations were reported in more highly trained athletes compared with less trained athletes<sup>(30)</sup>, yet folate supplementation of folate-deficient female marathon runners had no effect on aerobic capacity or running time<sup>(31)</sup>. Despite the lack of folate deficiency in the present study population, presumably due to folic acid fortification of wheat flour in Colombia, we found a linear association between erythrocyte folate and anaerobic performance indicators. Although it is unclear how folate could affect anaerobic performance, one possible mechanism may be via creatine, a reservoir for high-energy phosphate bonds that are needed

for ATP synthesis at the onset of intense exercise, because folate is needed for the methylation of *S*-adenosylmethionine, which serves as a methyl donor for creatine biosynthesis.

Vitamin B<sub>12</sub> could also potentially affect both aerobic and anaerobic performance-related fitness through the same mechanisms as folate. We found a positive association between vitamin B<sub>12</sub> status and jump scores among boys only but no associations with shuttle run scores. The jump test is a test of explosive power and may require more muscle strength than the run, consistent with a possible role of vitamin B<sub>12</sub> in creatine synthesis and muscle function. Supplementation studies conducted so far do not support a role of vitamin B<sub>12</sub> on fitness tests. A study of 18–21-year-old men receiving vitamin B<sub>12</sub> or placebo injections for 6 weeks found no effects on aerobic capacity, standing jump, hand grip or coordination tasks<sup>(32)</sup>. Montoye *et al.*<sup>(33)</sup> supplemented boys aged 12–17 years with vitamin B<sub>12</sub> or placebo for 7 weeks and found no effects on a half-mile run or a step test. These studies did not assess baseline vitamin B<sub>12</sub> status and were of short duration. Whether longer supplementation studies in populations with high prevalence of vitamin B<sub>12</sub> deficiency may yield positive effects on performance-related fitness deserves future investigation.

Our finding that stunted children scored lower in both the run and long jump tests than non-stunted children is consistent with previous studies<sup>(8,9)</sup>. For the long jump test, this relationship was only found in boys, consistent with another study in schoolchildren from Mozambique<sup>(9)</sup>. Stunting is thought to be caused by impaired growth of long leg bones during later infancy<sup>(34)</sup>. We found that stunting was negatively associated with both performance tests independent of muscle mass, suggesting that shorter leg length rather than less muscle mass may be responsible for the performance deficits. We did not find associations between arm muscle area and performance; however, Davies had found a linear relationship between lean leg volume and standing long jump distance in 12–13-year-old children<sup>(35)</sup>. We did not measure lean leg volume, and it is possible that arm muscle area estimated from the arm circumference and triceps skinfold is not as valid an indicator of leg muscle mass as lean leg volume. We also found that the effect of stunting was independent of micronutrient status.

We found that children who were classified as thin ran slightly but significantly slower than children with normal weight, independent of height. This finding is in agreement with another study in 12–15-year-old children from the Republic of Seychelles, where children with low BMI ran slower than those with normal BMI<sup>(36)</sup>. Thin children in the Republic of Seychelles also had lower jump scores, which agrees with our findings among girls only. Thin boys in our population may have differed in other nutritional characteristics, such as micronutrient status, from the children in the Republic of Seychelles. The negative effects of thinness may be due to less muscle mass, which we may have measured with some error by using arm muscle area.

Our finding that overweight or obese children had lower scores in the shuttle run and long jump than normal-weight children is consistent with previous observational studies<sup>(36,37)</sup>.

**Table 4.** Micronutrient status correlates of shuttle run scores in schoolchildren from Bogotá, Colombia  
(Mean values, standard deviations, number of children and 95 % confidence intervals)

	Shuttle run time (s)									
	Girls					Boys				
	<i>n</i>	Unadjusted mean	SD	Adjusted differences*	95 % CI	<i>n</i>	Unadjusted mean	SD	Adjusted differences*	95 % CI
Anaemia (Hb < 115 g/l)†										
No	939	13.13	1.66		Ref	877	12.30	1.45		Ref
Yes	37	13.43	1.69	-0.09	-0.60, 0.43	37	12.82	1.57	0.30	-0.11, 0.71
Plasma ferritin (µg/l)‡										
Normal	941	13.12	1.64		Ref	875	12.29	1.41		Ref
Low	31	13.86	1.95	0.63	0.12, 1.15	32	13.18	2.30	0.54	-0.01, 1.10
Erythrocyte folate (nmol/l)										
Q1 (<700.5)	242	13.39	1.59		Ref	199	12.48	1.36		Ref
Q2 (700.5–824.4)	226	13.15	1.53	-0.04	-0.28, 0.20	230	12.42	1.47	0	-0.22, 0.22
Q3 (824.5–976.0)	244	13.11	1.86	-0.17	-0.41, 0.07	217	12.22	1.38	-0.27	-0.49, -0.05
Q4 (>976.0)	216	12.75	1.44	-0.40	-0.63, -0.16	234	12.22	1.53	-0.22	-0.44, 0
<i>P</i> for trend		< 0.0001			0.0005		0.02			0.009
Plasma vitamin B <sub>12</sub> (pmol/l)										
Normal (>221)	825	13.08	1.59		Ref	752	12.32	1.45		Ref
Marginal or low (≤221)	108	13.51	2.00	0.28	-0.01, 0.57	138	12.21	1.44	-0.02	-0.21, 0.18

Q, quartile; Ref, reference; HAZ, height-for-age; CRP, C-reactive protein.

\* Estimates are from a multivariate linear general estimating equation model with covariates that include age, household socio-economic stratum, HAZ, BMI-for-age status, plasma ferritin, folate quartiles and plasma vitamin B<sub>12</sub>.

Note: Negative estimates for shuttle run time correspond to 'faster' run times and are considered as 'higher' scores.

† After subtracting 12 g/l from the individual values as an adjustment for an altitude of 2500 m<sup>(26)</sup>.

‡ Low ferritin was defined as < 15 µg/l if the concentration of CRP was ≤ 10 mg/l or < 30 µg/l if CRP > 10 mg/l<sup>(27)</sup>.



**Table 5.** Micronutrient status correlates of standing long jump scores in schoolchildren from Bogotá, Colombia (Mean values, standard deviations, number of children and 95 % confidence intervals)

	Standing long jump distance (cm)									
	Girls					Boys				
	<i>n</i>	Unadjusted mean	SD	Adjusted differences*	95 % CI	<i>n</i>	Unadjusted mean	SD	Adjusted differences*	95 % CI
Anaemia (Hb < 115 g/l)†										
No	968	116	23	Ref		891	130	22	Ref	
Yes	38	114	26	4	−3, 11	38	124	20	−1	−8, 5
Plasma ferritin‡ (µg/l)										
Normal	971	116	23	Ref		889	130	22	Ref	
Low	31	114	28	−2	−9, 5	32	119	25	−7	−12, −1
Erythrocyte folate (nmol/l)										
Q1 (<700.5)	246	113	24	Ref		199	127	20	Ref	
Q2 (700.5–824.4)	233	118	23	3	−1, 6	230	130	21	2	−1, 5
Q3 (824.5–976.0)	243	117	23	3	0, 7	220	131	22	4	0, 7
Q4 (>976.0)	236	119	22	4	1, 8	245	132	24	4	1, 8
<i>P</i> for trend		0.003			0.01		0.01			0.003
Plasma vitamin B <sub>12</sub> (pmol/l)										
Normal (>221)	837	117	23	Ref		756	131	22	Ref	
Marginal or low (≤221)	125	114	26	0	−4, 3	148	128	22	−4	−7, −1

Q, quartile; Ref, reference; HAZ, height-for-age; CRP, C-reactive protein.

\*Estimates are from a multivariate linear general estimating equation model with covariates that include age, household socio-economic stratum, hours spent playing outdoors per week, household per capita income, HAZ, BMI-for-age status, plasma ferritin, folate quartiles and plasma vitamin B<sub>12</sub>.

† After subtracting 12 g/l from the individual values as an adjustment for an altitude of 2500 m<sup>(26)</sup>.

‡ Low ferritin was defined as <15 µg/l if the concentration of CRP was ≤10 mg/l or <30 µg/l if CRP >10 mg/l<sup>(27)</sup>.

The reason why overweight children tend to perform less well on physical tasks that require projection of the body through space, such as runs and jumps, may be related with the fact that excess fat represents an additional load that must be moved<sup>(3)</sup>. The finding that arm fat area was associated with poorer performance independent of muscle area suggests that excess fat mass among overweight children impairs performance. It is unclear why the negative relationship between arm fat area and long jump scores was only present in boys; however, another study in schoolchildren found that subcutaneous fat in boys only was associated with lower long jump scores<sup>(8)</sup>. Weight reduction interventions in obese children and adolescents have resulted in improvements of both anaerobic and aerobic performance<sup>(38–40)</sup>.

Differences in fitness scores among anaemic *v.* non-anaemic children in the present study did not reach statistical significance. The prevalence of anaemia in our population was low (4%), which may have limited statistical power. Although Fe status would be expected to affect aerobic more than anaerobic performance due to Hb's role in oxygen transport, we found a significant association between low Fe stores assessed by ferritin and anaerobic performance-related fitness. Some studies have demonstrated improved aerobic capacity and endurance performance in non-anaemic women with low ferritin concentrations who received Fe supplements<sup>(10,11)</sup>. The effect of Fe supplementation on anaerobic performance-related fitness tests such as those in the present study is not known and deserves consideration in future studies.

A limitation of the present study is its cross-sectional design, which limits the possibility of drawing causal inferences. We were not able to assess motivation of the children, which is a major determinant of performance in fitness tests<sup>(41)</sup>. It is conceivable that Fe and vitamin B<sub>12</sub> deficiencies could affect motivation via their effects on cognition, and thus their potential effects on these tests would not be direct but mediated through improved motivation. We were also not able to assess the children's prior physical training as a predictor of performance in the fitness tests; however, we did find a positive association between the children's jump scores and reported time spent playing outdoors, a factor that has been correlated with children's physical activity<sup>(42,43)</sup>. Our finding of associations between long jump scores and anthropometric and micronutrient status indicators were independent of the association with time spent playing outdoors.

In conclusion, the present study suggests that there might be negative effects of suboptimal anthropometric and micronutrient status on anaerobic performance-related fitness of children from settings undergoing the nutritional transition. Further studies are needed to determine the effect of nutritional interventions on performance-related fitness of children in these settings.

### Acknowledgements

The present study was supported by the Secretary of Education of Bogotá, the David Rockefeller Center for Latin American Studies at Harvard University, the National University of Colombia and the National Institute of Health of Colombia.

J. E. A. was supported by training grant T32DK07703 from the National Institutes of Health. None of the authors has any conflicts of interest. The authors' contributions were as follows: J. E. A. was involved in the data analyses, interpretation of the results and writing of the first draft of the manuscript; M. M.-P. contributed to the study design and implementation of tests in the field; Y. F. contributed to the study implementation and laboratory analyses; S. L.-A. contributed to the study implementation and data management; G. J. was responsible for the implementation of physical performance tests in the field; A. B. participated in the study design and interpretation of the data; P. M. G. contributed to the interpretation of the data; E. V. contributed to the study design, data analyses, interpretation of the results and preparation of the manuscript. All authors critically reviewed the manuscript.

### References

1. Froberg K & Andersen LB (2005) Mini review: physical activity and fitness and its relations to cardiovascular disease risk factors in children. *Int J Obes* **29**, S34–S39.
2. Twisk JW, Kemper HC & van Mechelen W (2002) The relationship between physical fitness and physical activity during adolescence and cardiovascular disease risk factors at adult age. The Amsterdam Growth and Health Longitudinal Study. *Int J Sports Med* **23**, Suppl. 1, S8–S14.
3. Malina RM & Katzmarzyk PT (2006) Physical activity and fitness in an international growth standard for preadolescent and adolescent children. *Food Nutr Bull* **27**, S295–S313.
4. Tomkinson GR (2007) Global changes in anaerobic fitness test performance of children and adolescents (1958–2003). *Scand J Med Sci Sports* **17**, 497–507.
5. Tomkinson GR & Olds TS (2007) Secular changes in pediatric aerobic fitness test performance: the global picture. *Med Sport Sci* **50**, 46–66.
6. Doak CM, Adair LS, Monteiro C, *et al.* (2000) Overweight and underweight coexist within households in Brazil, China and Russia. *J Nutr* **130**, 2965–2971.
7. Duran P, Caballero B & de Onis M (2006) The association between stunting and overweight in Latin American and Caribbean preschool children. *Food Nutr Bull* **27**, 300–305.
8. Benefice E, Fouere T & Malina RM (1999) Early nutritional history and motor performance of Senegalese children, 4–6 years of age. *Ann Hum Biol* **26**, 443–455.
9. Prista A, Ribeiro Maia JA, Damasceno A, *et al.* (2003) Anthropometric indicators of nutritional status: implications for fitness, activity, and health in school-age children and adolescents from Maputo, Mozambique. *Am J Clin Nutr* **77**, 952–959.
10. Hinton PS, Giordano C, Brownlie T, *et al.* (2000) Iron supplementation improves endurance after training in iron-depleted, nonanemic women. *J Appl Physiol* **88**, 1103–1111.
11. Rowland TW, Deisroth MB, Green GM, *et al.* (1988) The effect of iron therapy on the exercise capacity of nonanemic iron-deficient adolescent runners. *Am J Dis Child* **142**, 165–169.
12. Beard J & Tobin B (2000) Iron status and exercise. *Am J Clin Nutr* **72**, Suppl., 594S–597S.
13. Ruiz JR, Sola R, Gonzalez-Gross M, *et al.* (2007) Cardiovascular fitness is negatively associated with homocysteine levels in female adolescents. *Arch Pediatr Adolesc Med* **161**, 166–171.

14. Solon FS, Sarol JN, Bernardo ABI, *et al.* (2003) Effect of a multiple-micronutrient-fortified fruit powder beverage on the nutrition status, physical fitness, and cognitive performance of schoolchildren in the Philippines. *Food Nutr Bull* **24**, Suppl. 4, S129–S140.
15. Arsenault JE, Mora-Plazas M, Forero Y, *et al.* (2009) Hemoglobin concentration is inversely associated with erythrocyte folate concentrations in Colombian school-age children, especially among children with low vitamin B<sub>12</sub> status. *Eur J Clin Nutr* **63**, 842–849.
16. McDonald CM, Baylin A, Arsenault JE, *et al.* (2009) Overweight is more prevalent than stunting and is associated with socioeconomic status, maternal obesity, and a snacking dietary pattern in school children from Bogotá, Colombia. *J Nutr* **139**, 370–376.
17. Maslova E, Mora-Plazas M, Forero Y, *et al.* (2009) Are vitamin A and iron deficiencies re-emerging in urban Latin America? A survey of schoolchildren in Bogota, Colombia. *Food Nutr Bull* **30**, 103–111.
18. Dekker LH, Mora-Plazas M, Marín C, *et al.* (2010) Stunting associated with poor socioeconomic and maternal nutrition status and respiratory morbidity in Colombian schoolchildren. *Food Nutr Bull* **31**, 242–250.
19. Arsenault JE, Mora-Plazas M, Forero Y, *et al.* (2009) Provision of a school snack is associated with vitamin B-12 status, linear growth, and morbidity in children from Bogota, Colombia. *J Nutr* **139**, 1744–1750.
20. Lohman T, Roche A & Martorell R (1988) *Anthropometric Standardization Reference Manual*. Champaign, IL: Human Kinetics Books.
21. Council of Europe & Committee for the Development of Sport (1988) *EUROFIT: European Test of Physical Fitness*. Rome: Committee for the Development of Sport within the Council of Europe.
22. World Health Organization (1983) *Measuring Change in Nutritional Status*. Geneva: WHO.
23. Cole TJ, Flegal KM, Nicholls D, *et al.* (2007) Body mass index cut offs to define thinness in children and adolescents: international survey. *BMJ* **335**, 194.
24. Cole TJ, Bellizzi MC, Flegal KM, *et al.* (2000) Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ* **320**, 1240–1245.
25. Gibson RS (2005) *Principles of Nutritional Assessment*, 2nd ed. New York, NY: Oxford University Press.
26. Nestle P (2002) *Adjusting Hemoglobin Values in Program Surveys*. Washington, DC: INACG.
27. Zimmermann MB & Hurrell RF (2007) Nutritional iron deficiency. *Lancet* **370**, 511–520.
28. Jones KM, Ramirez-Zea M, Zuleta C, *et al.* (2007) Prevalent vitamin B-12 deficiency in twelve-month-old Guatemalan infants is predicted by maternal B-12 deficiency and infant diet. *J Nutr* **137**, 1307–1313.
29. Food and Nutrition Board & Institute of Medicine (1998) *Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B<sub>6</sub>, Folate, Vitamin B<sub>12</sub>, Pantothenic Acid, Biotin, and Choline*. Washington, DC: National Academy Press.
30. Konig D, Bisse E, Deibert P, *et al.* (2003) Influence of training volume and acute physical exercise on the homocysteine levels in endurance-trained men: interactions with plasma folate and vitamin B<sub>12</sub>. *Ann Nutr Metab* **47**, 114–118.
31. Matter M, Stittfall T, Graves J, *et al.* (1987) The effect of iron and folate therapy on maximal exercise performance in female marathon runners with iron and folate deficiency. *Clin Sci* **72**, 415–422.
32. Tin-May-Than, Ma-Win-May, Skin-Sann-Aung, *et al.* (1978) The effect of vitamin B<sub>12</sub> on physical performance capacity. *Br J Nutr* **40**, 269–273.
33. Montoye HJ, Spata PJ, Pinckney V, *et al.* (1955) Effects of vitamin B<sub>12</sub> supplementation on physical fitness and growth of young boys. *J Appl Physiol* **7**, 589–592.
34. Cole TJ (2000) Secular trends in growth. *Proc Nutr Soc* **59**, 317–324.
35. Davies BN (1990) The relationship of lean limb volume to performance in the handgrip and standing long jump tests in boys and girls, aged 11.6–13.2 years. *Eur J Appl Physiol* **60**, 139–143.
36. Bovet P, Auguste R & Burdette H (2007) Strong inverse association between physical fitness and overweight in adolescents: a large school-based survey. *Int J Behav Nutr Phys Act* **4**, 24.
37. Malina RM, Beunen GP, Claessens AL, *et al.* (1995) Fatness and physical fitness of girls 7 to 17 years. *Obes Res* **3**, 221–231.
38. Dao HH, Frelut ML, Peres G, *et al.* (2004) Effects of a multidisciplinary weight loss intervention on anaerobic and aerobic aptitudes in severely obese adolescents. *Int J Obes* **28**, 870–878.
39. Deforche B, de Bourdeaudhuij I, Deboode P, *et al.* (2003) Changes in fat mass, fat-free mass and aerobic fitness in severely obese children and adolescents following a residential treatment programme. *Eur J Pediatr* **162**, 616–622.
40. Knöpfli BH, Radtke T, Lehmann M, *et al.* (2008) Effects of a multidisciplinary inpatient intervention on body composition, aerobic fitness, and quality of life in severely obese girls and boys. *J Adolesc Health* **42**, 119–127.
41. Cairney J, Hay JA, Faught BE, *et al.* (2008) Generalized self-efficacy and performance on the 20-metre shuttle run in children. *Am J Hum Biol* **20**, 132–138.
42. Sallis JF, Prochaska JJ & Taylor WC (2000) A review of correlates of physical activity of children and adolescents. *Med Sci Sports Exerc* **32**, 963–975.
43. Cleland V, Crawford D, Baur LA, *et al.* (2008) A prospective examination of children's time spent outdoors, objectively measured physical activity and overweight. *Int J Obes* **32**, 1685–1693.