

The gut microbiome of kittens is affected by dietary protein:carbohydrate ratio and associated with blood metabolite and hormone concentrations

Seema Hooda¹, Brittany M. Vester Boler¹, Katherine R. Kerr¹, Scot E. Dowd² and Kelly S. Swanson^{1*}

¹Department of Animal Sciences, University of Illinois, 1207 West Gregory Drive, Urbana, IL 61801, USA

²MR DNA Molecular Research LP, 503 Clovis Road, Shallowater, TX 79363, USA

(Submitted 24 February 2012 – Final revision received 28 May 2012 – Accepted 5 July 2012 – First published online 31 August 2012)

Abstract

High-protein, low-carbohydrate (HPLC) diets are common in cats, but their effect on the gut microbiome has been ignored. The present study was conducted to test the effects of dietary protein:carbohydrate ratio on the gut microbiota of growing kittens. Male domestic short-hair kittens were raised by mothers fed moderate-protein, moderate-carbohydrate (MPMC; n 7) or HPLC (n 7) diets, and then weaned at 8 weeks onto the same diet. Fresh faeces were collected at 8, 12 and 16 weeks; DNA was extracted, followed by amplification of the V4–V6 region of the 16S rRNA gene using 454 pyrosequencing. A total of 384 588 sequences (average of 9374 per sample) were generated. Dual hierarchical clustering indicated distinct clustering based on the protein:carbohydrate ratio regardless of age. The protein:carbohydrate ratio affected faecal bacteria. Faecal Actinobacteria were greater ($P < 0.05$) and Fusobacteria were lower ($P < 0.05$) in MPMC-fed kittens. Faecal *Clostridium*, *Faecalibacterium*, *Ruminococcus*, *Blautia* and *Eubacterium* were greater ($P < 0.05$) in HPLC-fed kittens, while *Dialister*, *Acidaminococcus*, *Bifidobacterium*, *Megasphaera* and *Mitsuokella* were greater ($P < 0.05$) in MPMC-fed kittens. Principal component analysis of faecal bacteria and blood metabolites and hormones resulted in distinct clusters. Of particular interest was the clustering of blood TAG with faecal Clostridiaceae, Eubacteriaceae, Ruminococcaceae, Fusobacteriaceae and Lachnospiraceae; blood ghrelin with faecal Coriobacteriaceae, Bifidobacteriaceae and Veillonellaceae; and blood glucose, cholesterol and leptin with faecal Lactobacillaceae. The present results demonstrate that the protein:carbohydrate ratio affects the faecal microbiome, and highlight the associations between faecal microbes and circulating hormones and metabolites that may be important in terms of satiety and host metabolism.

Key words: Kittens: Faecal microbiome: Protein:carbohydrate ratio

The significance of the microbiome in the gut and overall health has been well established, as it affects metabolism and modulation of the immune system, and provides protection against intestinal pathogens⁽¹⁾. Various prenatal and post-natal factors, such as the initial exposure to the environment, including the mother⁽²⁾, composition of milk⁽³⁾ and diet after weaning, contribute to the development of the core intestinal microbiota. The initial colonisation of the gastrointestinal tract has been shown to have significance in human infants, as it might define the lifelong composition of the gut microbiota and consequently contribute to host health⁽⁴⁾. The composition of the gut microbiota during early age also has been reported to affect the onset of metabolic diseases such as obesity in humans⁽⁵⁾.

A small number of studies have characterised the gastrointestinal microbiome of healthy adult cats^(6,7) and how it may be modified by dietary fibre⁽⁸⁾. Others have reported the gut microbial dysbiosis that occurs during inflammatory

bowel disease in cats⁽⁹⁾. To our knowledge, however, evaluation of the faecal microbiome of kittens around weaning has not been performed. Because weaning is a stressful period that often results in gut microbiota shifts and gastrointestinal stress, there is a great interest to characterise the gut microbial composition around weaning and design nutritional strategies that minimise these negative outcomes.

Cats are obligate carnivores, having evolved on diets rich in protein and fat. Thus, cats have a higher minimal crude protein requirement (18.0% for growing kittens and 16.0% for adult cats; NRC, 2006) and recommended allowance (22.5% for growing kittens and 20.0% for adult cats; NRC, 2006) than commonly studied animals (e.g. rodents, dogs) and human subjects. Most commercial cat foods contain a much higher amount of dietary protein than that minimally required, often containing 30–40% or more of diet dry weight. The macronutrient profile and the processing of such diets (high protein and fat; low fibre) not only make them highly

Abbreviations: 16S rRNA, 16S ribosomal RNA; HPLC, high protein, low carbohydrate; MPMC, moderate protein, moderate carbohydrate; PCA, principal component analysis.

* **Corresponding author:** Dr K. S. Swanson, fax +1 217 333 7861, email ksswanso@illinois.edu

palatable, but lead to high digestibility and low stool volume. While this may be beneficial from an owner's standpoint, the effects on hindgut fermentation have been poorly studied. Moreover, although cats and humans differ metabolically in many ways and cats do not live nearly as long as humans, the effects of diet on long-term intestinal health outcomes (e.g. disease incidence), if any, are unknown and require further study.

Using quantitative PCR, a recent study in adult cats in our laboratory demonstrated reduced faecal *Bifidobacterium* and *Lactobacillus* in those fed a high-protein, low-carbohydrate (HPLC) *v.* moderate-protein, moderate-carbohydrate (MPMC) diet⁽¹⁰⁾. Using quantitative PCR and denaturing gradient gel electrophoresis, a similar response in faecal microbiology was observed in kittens fed these same diets⁽¹¹⁾. Moreover, diet-induced changes in blood lipids and hormones, physical activity and body fat percentage were observed in these kittens⁽¹²⁾. Given the targeted microbiological and physiological response observed in these kittens previously, we hypothesised that a HPLC diet will promote the growth of proteolytic bacteria, whereas intake of a MPMC diet will increase the abundance of saccharolytic bacteria. Thus, the study was conducted to characterise the entire faecal microbiome of growing kittens fed a HPLC and MPMC diet using 16S ribosomal RNA (16S rRNA) gene pyrosequencing, and to identify the associations between the microbiota and host physiological data.

Materials and methods

Animals and diets

The animal protocol was approved by the University of Illinois Animal Care and Use Committee and was conducted at the Edward R. Madigan Laboratory of the University of Illinois. For the present study, eight domestic shorthair female cats (3.5 (SEM 0.24) kg body weight and 1.3 (SEM 0.02) years of age) and one male domestic shorthair cat (5.8 kg and 1.3 years of age) were used to produce kittens. All female cats were randomly assigned to two test diets, 1 month before mating, and continued throughout gestation and lactation. Kittens from mothers fed the MPMC diet (*n* 7) or the HPLC diet (*n* 7) were studied herein. The kittens were housed with dams until 8 weeks of age, weaned and then fed the same diets as mothers. After weaning, the kittens were twin- or triple-housed within the dietary group in cages (1 × 0.76 × 0.7 m) in temperature-controlled rooms with *ad libitum* food and water intake to allow for adequate growth. The kittens were allowed out of their cage to play with people and each other at least once per d to minimise post-weaning stress and to improve their socialisation. Fresh faecal samples (within 15 min of defecation) were collected from all kittens at 8, 12 and 16 weeks of age. All faecal samples were stored immediately at −80°C.

The diets were formulated and manufactured by Natura Manufacturing, Inc. The ingredients and chemical composition of both diets are given in Table 1. Both diets were formulated to meet or exceed all nutrient requirements for growth

Table 1. Ingredients and chemical composition of the moderate-protein, moderate-carbohydrate (MPMC) or high-protein, low-carbohydrate (HPLC) diet fed to kittens

Items	Diets	
	MPMC	HPLC
Ingredients (% as fed)		
Chicken meal	30.84	64.79
Dried potato product	40.38	10.69
Chicken fat	12.79	8.57
Dried egg	5.00	5.00
Herring meal	5.00	5.00
Beet pulp	3.00	3.00
Natural flavours	1.00	1.00
Herring oil	0.34	0.66
Mineral premix*†	0.65	0.65
Salt	0.25	0.25
Vitamin premix*‡	0.15	0.15
Potassium chloride	0.39	0.10
Dried chicory root	0.10	0.10
Dried natural antioxidant	0.03	0.05
DL-Met	0.07	0.00
Chemical composition (% DM)		
DM	94.22	94.83
Organic matter	91.28	89.19
Crude protein	34.34	52.88
Acid hydrolysed fat	19.23	23.55
Total dietary fibre	6.88	2.01
Gross energy (kJ/g)	21.80	23.20

*Trouw Nutrition (Highland, IL, USA).

†Composition of mineral premix (g/kg mix): CaCO₃, 360.2; ZnSO₄, 208.3; OPTiMIN (Trouw Nutrition) Zn proteinate (15% Zn; Trouw Nutrition), 166.7; FeSO₄, 77.4; OPTiMIN Fe proteinate (15% Fe), 53.3; CuSO₄, 35.7; OPTiMIN Cu proteinate (10% Cu), 3.0; MnSO₄, 23.4; OPTiMIN Mn proteinate (15% Mn), 16.7; Se, 12.0; carrier (soyabean oil), 10.0; OPTiMIN Co proteinate (2.5% Co), 3.8; I, 1.8; CoCO₃, 0.6.

‡Composition of vitamin premix (g/kg mix): carrier (pea fibre), 728.4; CaCO₃, 170.9; vitamin E (50% adsorbate), 40.0; betaine (source of choline), 26.0; carrier (soyabean oil), 10.0; nicotinic acid, 9.6; vitamin A, 4.0; D-calcium pantothenate, 2.7; vitamin B₁ (thiamin mononitrate), 2.7; vitamin B₂ (riboflavin), 1.25; β-carotene, 1.0; vitamin B₁₂, 1.0; vitamin D₃, 0.8; biotin, 0.7; vitamin B₆ (pyridoxine), 0.7; folic acid, 0.2.

according to the Association of American Feed Controls Officials (2007). Chemical analyses of the diets were done according to the procedures described by Vester *et al.*⁽¹¹⁾.

Faecal DNA extraction and pyrosequencing

Bacterial DNA was extracted using a QIAamp DNA stool mini kit (Qiagen) using the repeated bead beating plus column method⁽¹³⁾. Faecal DNA was quantified using a NanoDrop ND-1000 spectrophotometer (NanoDrop Technologies). Extracted DNA was diluted to 20 ng/μl and genomic DNA quality was assessed using electrophoresis using precast E-Gel[®] EX Gel 1% (Invitrogen). Amplification of a 600 bp sequence of the variable region V4–V6 of the 16S rRNA gene was done using barcoded primers according to Cephas *et al.*⁽¹⁴⁾. PCR amplicons of all samples were further purified using AMPure XP beads (Beckman-Coulter, Inc.) to remove smaller fragments. Further DNA concentration and quality was measured using a NanoDrop ND-1000 spectrophotometer and electrophoresis, respectively. Finally, the amplicons were combined in equimolar ratios to create a DNA pool that was used for pyrosequencing. DNA quality was assessed before

pyrosequencing using a 2100 Bioanalyzer (Agilent). Pyrosequencing of the PCR amplicons was performed at the W. M. Keck Center for Biotechnology at the University of Illinois using a 454 Genome Sequencer and FLX titanium reagents (Roche Applied Science). After sequencing was completed, all reads were scored for quality and any poor-quality reads and primer dimers were removed.

Bioinformatics

The sequences were selected to estimate total bacterial diversity of DNA samples and were trimmed to remove barcodes, primers, chimeras, plastid, mitochondrial, any non-16S bacterial reads and sequences <350 bp. A total of 4500 ± 100 sequences from each sample were selected based upon the highest average quality score, sequences trimmed to 350 bp and aligned with MUSCLE (Multiple Sequence Comparison by Log-Expectation)⁽¹⁵⁾, and then the distance matrix was calculated from the alignment with PHYLIP (PHYLogeny Inference Package). Operational taxonomical units was assigned by MOTHUR⁽¹⁶⁾. Bacterial identification community structure was evaluated using Phred20 quality reads, including both 530F and 1100R oriented (each analysed separately), trimmed to remove tags and primer sequence collections, and then depleted of chimera, plastid, mitochondrial and any non-16S reads (<70% identity to any known high-quality 16S sequence) and sequences <250 bp. The final sequence data were evaluated using Kraken (www.krakenblast.com) against a 10-21-10 version database curated from the NCBI to include >300 000 high-quality 16S bacterial sequences as well as quality-control screening sequences for mitochondria, plastid and chloroplast screening sequences. Blast output based upon top hit designations was compiled to generate percentage files at each taxonomic level as described previously^(17,18).

Statistical analyses

Sequence percentages at each taxonomic level were analysed as repeated measures using the Mixed Models procedure of SAS (version 9.2; SAS Institute, Inc.). The statistical model included kitten as a random effect and diet, age and diet \times age as fixed effects. Means were separated for treatments using a Fisher-protected least significant difference with Tukey's adjustment. Results are reported as least-squares means with $P < 0.05$ defined as significant and $0.05 \leq P < 0.10$ as trends. Principal component analysis (PCA) of bacterial families and blood metabolites and hormones at 8 weeks was done using JMP software of SAS (SAS Institute Inc.).

Results

Kitten faecal microbiota

Pyrosequencing of 16S rRNA gene barcoded amplicons resulted in a total of 384 588 sequences, with an average of 9374 (range 5204–13 456) per sample. The operational taxonomical unit estimates of diversity were not different

($P > 0.05$) among the groups (data not shown), which indicates that faecal bacterial diversity was not different among the dietary groups or ages. The dual hierarchical clustering dendrogram of the fifty most abundant bacterial genera indicated distinct clustering of samples based on the protein:carbohydrate ratio regardless of age (Fig. 1). Faecal *Collinsella*, *Faecalibacterium*, *Eubacterium*, *Anaerotruncus*, *Ruminococcus*, *Fusobacterium*, *Blautia* and *Clostridium* were most abundant in HPLC-fed kittens. In contrast, kittens fed the MPMC diet had higher abundance of *Megasphaera*, *Bifidobacterium* and *Mitsuokella*. Further analysis using Uni Frac PCA also showed that kittens fed the HPLC diet were clustered distinct from MPMC-fed kittens regardless of age (Fig. 2).

Firmicutes was the most abundant bacterial phylum in all kitten faeces at all ages, followed by Actinobacteria, Fusobacteria and Proteobacteria (Table 2). A diet \times age interaction was observed for faecal Firmicutes ($P < 0.05$) and Actinobacteria ($P < 0.01$). Kittens fed the MPMC and HPLC diets had similar faecal Firmicutes at 8 weeks of age. Kittens fed the MPMC diet had lower faecal Firmicutes at 12 and 16 weeks of age compared with 8 weeks of age ($P < 0.05$), whereas no change was observed with age for kittens fed the HPLC diet. Faecal Actinobacteria abundance was greater ($P < 0.05$) in MPMC-fed *v.* HPLC-fed kittens at 8 weeks. The disparity in faecal Actinobacteria grew with age, as this population did not change over time in kittens fed the HPLC diet, but continued to increase ($P < 0.05$) with age in MPMC-fed kittens. In contrast, the proportion of faecal Fusobacteria and Proteobacteria was greater ($P < 0.0001$) in HPLC-fed *v.* MPMC-fed kittens. Fusobacteria were almost absent in MPMC-fed kittens, but the second most prevalent phylum in HPLC-fed kittens, regardless of age.

Among the Firmicutes, Veillonellaceae, Ruminococcaceae, Clostridiaceae, Erysipelotrichaceae, Lachnospiraceae, Lactobacillaceae and Eubacteriaceae were predominant families in kitten faeces (Table 2). Veillonellaceae was the most abundant bacterial family present in MPMC-fed kittens and greater ($P < 0.0001$) than in HPLC-fed kittens at all time points. However, faecal Clostridiaceae, Eubacteriaceae and Lachnospiraceae were greater ($P < 0.001$) in HPLC-fed kittens. A significant ($P < 0.001$) diet \times age interaction was observed for faecal Ruminococcaceae and Lactobacillaceae. Faecal Ruminococcaceae was not affected by diet at 8 weeks of age. After weaning, however, faecal Ruminococcaceae increased in HPLC-fed kittens and decreased in MPMC-fed kittens. Faecal Lactobacillaceae abundance was greater ($P < 0.05$) in kittens fed the MPMC *v.* HPLC diet at 8 weeks of age, but were decreased to abundance similar to that of HPLC-fed kittens after weaning. The abundance of faecal Erysipelotrichaceae was only affected by age, being quite low at 8 weeks of age and then increased ($P < 0.05$) with age in both dietary treatment groups.

Within the phylum Actinobacteria, the proportion of faecal Coriobacteriaceae and Bifidobacteriaceae demonstrated a diet \times age interaction ($P < 0.05$). At 8 weeks, MPMC-fed kittens had greater faecal Bifidobacteriaceae abundance compared with HPLC-fed kittens. In MPMC-fed kittens, the abundance of faecal Bifidobacteriaceae increased ($P < 0.05$) with age. Faecal Bifidobacteriaceae, however, remained very low in




Fig. 1. Dual hierarchical clustering dendrogram of the fifty most abundant bacterial genera among the kitten samples collected at 8, 12 and 16 weeks of age and fed the moderate-protein, moderate-carbohydrate (M) or high-protein, low-carbohydrate (H) diet. This double dendrogram is based on the Wards clustering and Manhattan distance methods. The heat map depicts the relative percentage of each genus for each sample. The relative distance scale for the left y-axis is provided in the lower left corner of the figure. The colour scale for the heat map is shown in the upper left corner of the figure.




Fig. 2. Principal component analysis of Uni Frac distance metric. Light and dark blue, and black balls represent high-protein, low-carbohydrate (HPLC; H)-fed kittens and orange, red and purple balls represent moderate-protein, moderate-carbohydrate (MPMC; M)-fed kittens. Kittens fed the HPLC diet were clustered distinct from MPMC-fed kittens regardless of age.

Table 2. Bacterial families (expressed as a percentage of sequences) in the faeces of kittens fed a moderate-protein, moderate-carbohydrate (MPMC) or high-protein, low-carbohydrate (HPLC) diet at 8, 12 and 16 weeks of age as determined by 16S rRNA gene pyrosequencing (Mean values with their pooled standard errors, *n* 7)

	Diets						SEM	<i>P</i>		
	MPMC			HPLC				Diet	Age	Diet × age
	Age (weeks)			Age (weeks)						
	8	12	16	8	12	16				
Firmicutes	80.10 ^b	72.69 ^a	70.99 ^a	75.27 ^{a,b}	79.14 ^b	78.28 ^b	2.88	0.34	0.49	0.04
Clostridiaceae	8.34	4.20	5.77	28.28	20.69	26.12	1.48	<0.0001	0.0002	0.27
Clostridiales*	4.89 ^a	3.81 ^a	3.96 ^a	15.17 ^b	17.93 ^b	14.71 ^b	1.14	<0.0001	0.10	0.02
Veillonellaceae	42.38	50.83	43.28	1.43	2.39	2.89	2.43	<0.0001	0.05	0.07
Ruminococcaceae	11.63 ^b	5.91 ^a	7.16 ^{a,b}	16.89 ^b	23.73 ^c	21.14 ^c	1.78	<0.0001	0.88	0.0002
Eubacteriaceae	0.72 ^a	0.08 ^a	0.07 ^a	6.80 ^b	7.98 ^b	6.19 ^b	0.74	<0.0001	0.04	0.04
Lachnospiraceae	0.89 ^a	1.11 ^a	1.93 ^a	2.84 ^b	2.31 ^{a,b}	2.15 ^{a,b}	0.31	0.001	0.53	0.02
Erysipelotrichaceae	1.18	5.18	5.85	0.82	2.66	3.66	1.05	0.18	<0.0001	0.37
Lactobacillaceae	6.97 ^b	0.11 ^a	0.78 ^a	1.50 ^a	0.08 ^a	0.05 ^a	1.03	0.09	<0.0001	0.004
Actinobacteria	18.21 ^b	26.49 ^c	28.45 ^c	7.92 ^a	4.59 ^a	5.24 ^a	2.67	<0.0001	0.32	0.001
Coriobacteriaceae	6.18 ^{a,b}	9.03 ^b	7.71 ^{a,b}	7.78 ^{a,b}	4.51 ^a	5.11 ^a	1.27	0.20	0.86	0.02
Bifidobacteriaceae	12.02 ^b	17.62 ^c	20.74 ^c	0.06 ^a	0.04 ^a	0.10 ^a	2.32	<0.0001	0.03	0.03
Fusobacteria	0.10	0.17	0.07	12.45	12.33	12.94	1.40	<0.0001	0.98	0.96
Fusobacteriaceae	0.11	0.10	0.09	12.46	12.34	12.96	1.40	<0.0001	0.97	0.97
Bacteroidetes	0.55	0.59	0.20	0.55	0.65	0.16	0.19	0.91	0.01	0.87
Proteobacteria	1.06	0.08	0.28	3.68	3.28	3.34	0.35	<0.0001	0.06	0.11

^{a,b,c} Mean values with unlike superscript letters within a row were significantly different (*P*<0.05).

* Unknown family within the order Clostridiales.

HPLC-fed kittens throughout the study. Faecal Coriobacteriaceae were not affected by diet at 8 weeks. After weaning, however, faecal Coriobacteriaceae decreased in HPLC-fed kittens and increased in MPMC-fed kittens. Faecal Fusobacteriaceae were only affected by diet, with kittens fed the HPLC diet having greater ($P < 0.0001$) abundance than those fed the MPMC diet.

Among the well-characterised genera within the phylum Firmicutes, a significant ($P < 0.05$) diet \times age interaction was observed for faecal *Blautia*, *Dialister*, *Acidaminococcus*, *Megasphaera*, *Phascolarctobacterium*, *Ruminococcus*, *Faecalibacterium*, *Subdoligranulum*, *Eubacterium*, *Catenibacterium* and *Lactobacillus* (Table 3). Faecal *Dialister*, *Acidaminococcus* and *Catenibacterium* were lower ($P < 0.001$) in HPLC-fed kittens than in MPMC-fed kittens. Faecal *Megasphaera* also was lower ($P < 0.0001$) in HPLC-fed kittens, but greater at 12 weeks of age compared with 8 weeks of age in MPMC-fed kittens. In contrast, faecal *Blautia*, *Ruminococcus*, *Faecalibacterium* and *Eubacterium* were lower ($P < 0.001$) in MPMC-fed kittens than in HPLC-fed kittens at all ages. Faecal *Clostridium* and *Mitsuokella* were affected ($P < 0.0001$) by diet; *Clostridium* being higher and *Mitsuokella* being lower in HPLC-fed kittens. Faecal *Olsenella* and *Collinsella* were affected ($P < 0.001$) by diet. Faecal *Olsenella* abundance was greater ($P < 0.0001$) and faecal *Collinsella* was lower ($P < 0.001$) in MPMC-fed kittens. Faecal *Bifidobacterium* had significant ($P < 0.05$) diet \times age interactions. Faecal *Bifidobacterium* abundance was greater ($P < 0.05$) in MPMC-fed kittens at

8 weeks of age, and increased ($P < 0.05$) with age. Faecal *Bifidobacterium* remained low in HPLC-fed kittens throughout the study. In contrast, faecal *Fusobacterium* was greater ($P < 0.0001$) in HPLC-fed kittens. The abundance of faecal *Lactobacillus* was highest at 8 weeks in MPMC-fed kittens and then decreased ($P < 0.05$) after weaning to abundance similar to that of HPLC-fed kittens.

The abundance of faecal *Clostridium hiranonis*, *Clostridium citroniae*, *Blautia glucerosea* and *Eubacterium coprostanoligenes* was higher ($P < 0.001$) in HPLC-fed *v.* MPMC-fed kittens (Table 4). In contrast, faecal *Megasphaera elsdenii* was affected by age \times diet interaction ($P < 0.0001$), being highest at 12 weeks of age in MPMC-fed kittens. The abundance of faecal *Catenibacterium mitsuokai* was greater ($P < 0.05$) at 12 and 16 weeks than at 8 weeks of age in MPMC-fed kittens. Faecal *Olsenella profusa* was only affected by diet and was predominant ($P < 0.001$) in HPLC-fed kittens.

The PCA of faecal bacterial families, blood metabolites and hormones resulted in four distinct clusters (Fig. 3). One cluster included blood TAG and faecal Clostridiaceae, Eubacteriaceae, Ruminococcaceae, Fusobacteriaceae and Lachnospiraceae. A second cluster consisted of blood ghrelin and faecal Coriobacteriaceae, Bifidobacteriaceae and Veillonellaceae. A third cluster consisted of blood glucose, cholesterol and leptin, and faecal Lactobacillaceae. The last cluster included body weight, blood creatinine, urea N, protein and albumin. The PCA loading plot indicated that blood TAG are negatively correlated with faecal Veillonellaceae and faecal Bifidobacter-

Table 3. Bacterial genera (expressed as a percentage of sequences) in the faeces of kittens fed a moderate-protein, moderate-carbohydrate (MPMC) or high-protein, low-carbohydrate (HPLC) diet at 8, 12 and 16 weeks of age as determined by 16S rRNA gene pyrosequencing (Mean values with their pooled standard errors, n 7)

	Diets						SEM	<i>P</i>		
	MPMC			HPLC				Diet	Age	Diet \times age
	Age (weeks)			Age (weeks)						
	8	12	16	8	12	16				
Firmicutes										
<i>Clostridium</i>	6.25	3.41	4.36	18.05	14.38	17.42	1.16	<0.0001	0.008	0.58
Clostridiaceae*	2.04 ^a	0.68 ^a	1.34 ^a	10.11 ^c	6.24 ^b	8.58 ^{b,c}	0.58	<0.0001	<0.0001	0.04
<i>Blautia</i>	2.78 ^a	2.24 ^a	2.99 ^a	7.20 ^b	9.75 ^b	8.46 ^b	0.74	<0.0001	0.13	0.01
<i>Dialister</i>	9.38 ^c	5.44 ^b	5.42 ^b	0.02 ^a	0.01 ^a	0.01 ^a	0.81	<0.0001	0.001	0.001
<i>Acidaminococcus</i>	5.05 ^c	1.93 ^b	3.75 ^{b,c}	0.01 ^a	0.01 ^a	0.01 ^a	0.26	<0.0001	<0.0001	<0.0001
<i>Megasphaera</i>	17.85 ^b	32.99 ^c	23.87 ^{b,c}	0.01 ^a	0.10 ^a	0.10 ^a	1.80	<0.0001	<0.0001	<0.0001
<i>Mitsuokella</i>	8.79	7.64	10.12	0.01	0.01	0.02	1.23	<0.0001	0.57	0.58
<i>Phascolarctobacterium</i>	0.02 ^a	0.04 ^a	0.02 ^a	1.48 ^b	2.30 ^b	2.56 ^b	0.24	<0.0001	0.0006	0.001
<i>Ruminococcus</i>	3.00 ^a	2.71 ^a	3.76 ^a	10.74 ^b	15.26 ^c	13.47 ^{b,c}	0.71	<0.0001	0.001	0.0004
<i>Faecalibacterium</i>	2.46 ^a	1.21 ^a	0.11 ^a	4.96 ^{a,b}	7.25 ^b	6.68 ^b	1.33	0.0004	0.55	0.02
<i>Subdoligranulum</i>	5.89 ^b	1.92 ^a	3.19 ^{a,b}	0.25 ^a	0.11 ^a	0.12 ^a	0.97	0.004	0.02	0.03
<i>Eubacterium</i>	1.43 ^a	0.53 ^a	0.95 ^a	6.86 ^b	8.38 ^b	6.79 ^b	0.78	<0.0001	0.43	0.02
<i>Catenibacterium</i>	0.05 ^a	4.39 ^b	4.33 ^b	0.06 ^a	0.01 ^a	0.01 ^a	0.61	0.0002	0.001	0.0008
<i>Lactobacillus</i>	6.97 ^b	0.11 ^a	0.77 ^a	1.49 ^a	0.08 ^a	0.05 ^a	1.03	0.09	<0.0001	0.004
Actinobacteria										
<i>Coriobacterium</i>	0.65	0.63	0.67	0.41	0.38	0.48	0.12	0.08	0.79	0.94
<i>Olsenella</i>	3.18	6.76	5.57	0.01	0.05	0.06	0.83	<0.0001	0.06	0.07
<i>Collinsella</i>	2.29	1.56	1.40	6.74	3.88	4.39	0.72	0.0009	0.01	0.24
<i>Bifidobacterium</i>	12.02 ^b	17.61 ^c	20.78 ^c	0.06 ^a	0.04 ^a	0.10 ^a	2.31	<0.0001	0.03	0.03
Fusobacteria										
<i>Fusobacterium</i>	0.12	0.11	0.11	10.42	10.76	11.06	0.83	<0.0001	0.06	0.07

^{a,b,c} Mean values with unlike superscript letters within a row were significantly different ($P < 0.05$).

* Unknown genus within the family Clostridiaceae.

Table 4. Bacterial species (expressed as a percentage of sequences) in the faeces of kittens fed a moderate-protein, moderate-carbohydrate (MPMC) or high-protein, low-carbohydrate (HPLC) diet at 8, 12 and 16 weeks of age as determined by 16S rRNA gene pyrosequencing (Mean values with their pooled standard errors, *n* 7)

	Diets						SEM	<i>P</i>		
	MPMC			HPLC				Diet	Age	Diet × age
	Age (weeks)			Age (weeks)						
	8	12	16	8	12	16				
<i>Clostridium hiranonis</i>	1.33	0.62	0.95	2.86	2.55	3.49	0.65	0.0002	0.66	0.79
<i>Clostridium hathewayi</i>	0.05	0.03	0.02	0.26	0.20	0.26	0.03	0.0001	0.42	0.47
<i>Clostridium citroniae</i>	0.36	0.03	0.00	1.44	0.69	0.86	0.15	0.0004	0.001	0.20
<i>Blautia glucerasea</i>	1.58	0.60	1.40	3.94	4.87	4.41	0.51	0.0001	0.89	0.05
<i>Blautia schinkii</i>	0.37 ^a	0.09 ^a	0.09 ^a	1.71 ^{b,c}	2.2 ^c	1.58 ^b	0.16	0.0001	0.06	0.01
<i>Eubacterium coprostanoligenes</i>	0.63	0.01	0.01	2.96	1.78	1.91	0.44	0.004	0.002	0.56
<i>Eubacterium eligens</i>	0.14 ^a	0.14 ^a	0.14 ^a	0.77 ^b	4.75 ^c	2.92 ^d	0.47	0.0005	0.0001	0.0001
<i>Megasphaera elsdenii</i>	17.85 ^b	32.99 ^c	23.87 ^{b,c}	0.01 ^a	0.10 ^a	0.10 ^a	1.80	<0.0001	<0.0001	<0.0001
<i>Catenibacterium mitsuokai</i>	0.05 ^a	3.88 ^b	3.86 ^b	0.06 ^a	0.01 ^a	0.01 ^a	0.54	0.0002	0.001	0.0009
<i>Olsenella profuse</i>	2.31	5.03	4.31	0.01	0.05	0.01	0.70	0.0002	0.08	0.09

a,b,c,d Mean values with unlike superscript letters within a row were significantly different (*P*<0.05).

iaceae (angle close to 180° among variables). Similarly, faecal Lactobacillaceae had a negative association with body weight and positive with blood leptin.

Discussion

There has been a recent surge in research studying the impact of macronutrient ratios, especially protein:carbohydrate, on human and animal metabolism. In feline nutrition, HPLC diets are often marketed based on the carnivorous nature of cats. Moreover, a HPLC diet has been suggested to prevent metabolic diseases such as obesity and type 2 diabetes⁽¹⁹⁾. Although HPLC diets have been promoted to improve the metabolic health of humans and companion animals, their effect on the gut microbiome is not known. The commensal microbes are important in the development of the gut immune system and pathogen resistance, and thus the prevention of gastrointestinal diseases⁽²⁰⁾. A few studies have provided a detailed view of the gut microbiome of healthy adult cats using next-generation pyrosequencing techniques^(6,7). However, to our knowledge, the present study is the first to describe the dietary and age effects in growing kittens using 16S rRNA gene pyrosequencing.

The predominant bacterial phylum present in our kittens (i.e. Firmicutes) was similar to previous reports in adult cats^(6,21) and dogs⁽²²⁾. Interestingly, the abundance of faecal Firmicutes decreased with age slightly, with approximately 10% of sequences being replaced by faecal Actinobacteria in MPMC-fed kittens. In contrast, no changes were observed for faecal Firmicutes, Actinobacteria and Fusobacteria with increasing age in HPLC-fed kittens.

The colonisation of the newborn gut usually starts immediately after birth. Various internal and external factors contribute to the development of the core microbiota. In humans, the gut microbiota has been considered stable at 2 years of age and almost identical to adults⁽²³⁾. This has not been adequately studied in cats. The initial colonisation of the gastrointestinal tract is believed to be quite important in

human infants, as it might define the lifelong composition of the gut microbiota and consequently contribute to host health⁽⁴⁾. Specifically, the genus *Bifidobacterium* has been linked to weight gain. Bifidobacteria numbers were shown to be higher in normal-weight *v.* overweight children, implying that high *Bifidobacterium* in infancy may provide protection against metabolic diseases later in life⁽⁵⁾. Bifidobacteria are also positively linked with gastrointestinal health. Lower faecal *Bifidobacterium*, for example, has been associated with the incidence of inflammatory bowel disease in




Fig. 3. Principal component analysis loading plot of the primary faecal bacterial families and blood metabolites and hormones of interest in seven healthy growing kittens fed the moderate-protein, moderate-carbohydrate or high-protein, low-carbohydrate diet. The loading plots indicate the relationships among variables. 1, Blood leptin; 2, blood ghrelin; 3, body weight; 4, blood creatinine; 5, blood urea nitrogen; 6, blood protein; 7, blood albumin; 8, blood glucose; 9, blood cholesterol; 10, blood TAG; 11, faecal Eubacteriaceae; 12, faecal Ruminococcaceae; 13, faecal Coriobacteriaceae; 14, faecal Fusobacteriaceae; 15, faecal Clostridiaceae; 16, faecal Veillonellaceae; 17, faecal Lachnospiraceae; 18, faecal Bifidobacteriaceae; 19, faecal Lactobacillaceae.

adult cats⁽²⁴⁾. The vast differences in *Bifidobacterium* in the present study, with an almost absence of *Bifidobacterium* in HPLC-fed kittens, are of great interest. High protein intake may lead to an increased flow of undigested protein to the colon⁽²⁵⁾, which is subjected to fermentation and the production of harmful nitrogenous metabolites⁽²⁶⁾. Fermentative end products, such as branched-chain fatty acids, NH₃, indoles and phenols, have been associated with increased colon cancer⁽²⁷⁾ and ulcerative colitis⁽²⁸⁾ in humans. Although cats differ from humans metabolically, are obligate carnivores that have evolved consuming high protein concentrations, and do not have a high incidence of colon cancer, they do develop inflammatory bowel diseases that may be contributed to by diet. Thus, the effects of diet on the feline gut microbiome and its potential role in disease should be studied in the future. The present data agree with those of earlier studies in adult cats⁽¹⁰⁾ and kittens⁽¹¹⁾ fed these same diets in which a HPLC diet led to reduced faecal *Bifidobacterium*. The near absence of *Bifidobacterium* in HPLC-fed kittens could have been due to the lack of carbohydrate-based substrates, the negative effects of protein-fermentative metabolites or competitive exclusion by protein-fermenting microbes in the gut. While these data would indicate detrimental effects of high protein intake in kittens, all kittens remained healthy throughout the study, without any signs of gastrointestinal distress (e.g. loose stools, excessive gas), and are still maintained in our colony in adulthood.

The presence of faecal Fusobacteria in HPLC-fed kittens is not surprising, as this group has been reported to be proteolytic and to ferment various amino acids and be only weakly saccharolytic^(29,30). In healthy adult cats, a lower abundance of faecal Fusobacteria has been reported previously^(6,7). Although this bacterial group has been associated with liver cirrhosis, appendicitis⁽³¹⁾ and various oral cavity infections⁽³²⁾ in humans, all of the kittens in the present study were healthy and had Fusobacteria populations similar to those of dogs in recent studies^(22,33).

Among the Firmicutes, faecal Clostridiaceae, Ruminococcaceae, Eubacteriaceae, which include the genera *Clostridium*, *Blautia*, *Ruminococcus* and *Faecalibacterium*, were dominant in HPLC-fed kittens. *C. bيرانonis* was one species detected in greater amounts in HPLC-fed kittens. This species exhibits bile acid 7 α -dehydroxylation activity, which results in the 7 α -dehydroxylation of bile primary salts to secondary salts⁽³⁴⁾. The presence of secondary bile acids is correlated with the increased risk of colon cancer in humans⁽³⁵⁾. Similarly, *Clostridium bathewayi* and *C. citroniae* have been reported to contribute to acute cholecystitis, hepatic abscess and bacteraemia⁽³⁶⁾, acute appendicitis⁽³⁷⁾, and clinical infections⁽³⁸⁾. In contrast, the families Ruminococcaceae, Lachnospiraceae and Eubacteriaceae including genera such as *Ruminococcus*, *Faecalibacterium*, *Eubacterium* and *Phascolarctobacterium* were greater in HPLC-fed kittens. These families are known to degrade dietary fibre and produce SCFA⁽³⁹⁾. In particular, *E. coprostanoligenes* has been shown to have hypocholesterolaemic effects⁽⁴⁰⁾.

Another lactate-utilising butyrate producer, *M. elsdenii*, abundance was significantly greater in MPMC-fed kittens.

This species has been used as a probiotic to promote recovery from diarrhoea in rats⁽⁴¹⁾, and is known to improve mucosal atrophy in the small and large intestine of weaning pigs⁽⁴²⁾. Moreover, *Megasphaera* is one of the major butyrate producers in the gut and thus promotes intestinal health⁽⁴³⁾. The lactate producer, *Mitsuokella*, may support *Megasphaera* (lactate utiliser) in MPMC-fed kittens and highlight the significance of cross-feeding. Although the complete absence of these two bacterial groups in HPLC-fed kittens may suggest detrimental effects of excess protein intake on gut microbes, there was no sign of weaning stress in any kittens.

The abundance of faecal *Lactobacillus* was higher in MPMC-fed kittens and decreased after weaning (8 weeks). Various studies have reported significant decreases of *Lactobacillus* after weaning^(43,44), often explained by unavailability of substrates (lactate) and change in gastric pH. Studies in pigs have indicated that early colonisation of *Lactobacillus* influences the development of the immune system, intestinal structure and integrity, and decreased pathogen load⁽⁴⁵⁾, thus improving intestinal and overall health^(46,47). A higher number of probiotic *Lactobacillus* also have been reported to prevent gastrointestinal disturbances related to weaning stress in piglets⁽⁴⁸⁾. Considering the significance of *Lactobacillus* at the time of weaning, a diet containing a moderate balance of protein and carbohydrates may provide greater stability than one containing very high concentrations of protein.

The PCA of faecal microbial data indicated distinct clustering and separation of samples of kittens fed the HPLC diet at all ages. However, no distinct clustering was observed for MPMC-fed kittens, indicating that the HPLC-based effects were more significant than the age effects on gut microbial communities. Similar separation due to the diet has been reported using denaturing gradient gel electrophoresis banding patterns previously, which further confirms that protein intake significantly affects the composition of gut microbial communities⁽¹¹⁾.

The PCA of abundant faecal bacterial families and blood hormones and metabolites indicated some interesting associations. One association to note was that of the well-known probiotic bacteria, *Lactobacillus*, with blood leptin concentrations. Leptin has an important role in satiety, interacting with hypothalamic receptors to control appetite and energy storage⁽⁴⁹⁾. Other interesting associations involved the bacterial family Bifidobacteriaceae, which was positively correlated with fasting blood ghrelin and negatively correlated with blood TAG. Human studies have reported the positive effects of probiotics on blood lipids through increased deconjugation of bile salts, resulting in greater excretion, which in turn leads to lower enterohepatic circulation⁽⁵⁰⁾. Although faecal bile salts were not measured in the present study, the present results seem to agree with previous reports and highlight the significance of the gut microbiome to host metabolism and satiety.

In conclusion, the present study demonstrated a significant impact of dietary protein:carbohydrate ratio on the composition of the gut bacterial microbiome in growing kittens. These data clearly demonstrate that differences in dietary protein and carbohydrate affect the populations of bacteria,



especially within the Firmicutes, Actinobacteria and Fusobacteria phyla. In general, high protein intake appeared to have a potential negative impact on the amount of health-promoting bacteria, such as *Bifidobacterium*, *Megasphaera* and *Lactobacillus*. Furthermore, strong associations between blood hormones and metabolites and well-known probiotic bacterial families indicate potential significance as it pertains to satiety and host metabolism. Further research is required to identify any health implications of having such different gut microbiota populations early in life, namely how long-term metabolism and health of the host may be affected.

Acknowledgements

This study received no specific grant from any funding agency in the public, commercial or not-for-profit sectors. S. H., B. M. V. B. and K. S. S. designed the research; S. H., K. R. K. and B. M. V. B. conducted the research; S. H. and S. E. D. analysed the data; S. H. and K. S. S. wrote the paper; K. S. S. had primary responsibility for the final content. All authors read and approved the final manuscript. The authors declare that there is no conflict of interest.

References

- Hooper LV, Wong MH, Thelin A, *et al.* (2001) Molecular analysis of commensal host–microbial relationships in the intestine. *Science* **291**, 881–884.
- Mackie RI, Sghir A & Gaskins HR (1999) Developmental microbial ecology of the neonatal gastrointestinal tract. *Am J Clin Nutr* **69**, 1035S–1045S.
- Gronlund MM, Gueimonde M, Laitinen K, *et al.* (2007) Maternal breast-milk and intestinal bifidobacteria guide the compositional development of the *Bifidobacterium* microbiota in infants at risk of allergic disease. *Clin Exp Allergy* **37**, 1764–1772.
- Adlerberth I (2008) Factors influencing the establishment of the intestinal microbiota in infancy. *Nestle Nutr Workshop Ser Pediatr Program* **62**, 13–29.
- Kalliomaki M, Salminen S & Isolauri E (2008) Positive interactions with the microbiota: probiotics. *Adv Exp Med Biol* **635**, 57–66.
- Ritchie LE, Steiner JM & Suchodolski JS (2008) Assessment of microbial diversity along the feline intestinal tract using 16S rRNA gene analysis. *FEMS Microbiol Ecol* **66**, 590–598.
- Handl S, Dowd SE, Garcia-Mazcorro JF, *et al.* (2011) Massive parallel 16S rRNA gene pyrosequencing reveals highly diverse fecal bacterial and fungal communities in healthy dogs and cats. *FEMS Microbiol Ecol* **76**, 301–310.
- Barry KA (2010) Indices of gut health and intestinal microbial ecology of the cat as affected by ingestion of select carbohydrates varying in fermentative capacity. PhD Thesis, University of Illinois
- Janeczko S, Atwater D, Bogel E, *et al.* (2008) The relationship of mucosal bacteria to duodenal histopathology, cytokine mRNA, and clinical disease activity in cats with inflammatory bowel disease. *Vet Microbiol* **128**, 178–193.
- Lubbs DC, Vester BM, Fasting N, *et al.* (2009) Dietary protein concentration affects intestinal microbiota of adult cats: a study using DGGE and qPCR to evaluate differences in microbial populations in the feline gastrointestinal tract. *J Anim Physiol Anim Nutr* **93**, 113–121.
- Vester BM, Dalsing BL, Middelbos IS, *et al.* (2009) Faecal microbial populations of growing kittens fed high- or moderate-protein diets. *Arch Anim* **63**, 254–265.
- Vester BM, Liu KJ, Keel TL, *et al.* (2009) *In utero* and post-natal exposure to a high-protein or high-carbohydrate diet leads to differences in adipose tissue mRNA expression and blood metabolites in kittens. *Br J Nutr* **102**, 1136–1144.
- Yu Z & Morrison M (2004) Improved extraction of PCR-quality community DNA from digesta and fecal samples. *Biotechniques* **36**, 808–812.
- Cephas KD, Kim J, Mathai RA, *et al.* (2011) Comparative analysis of salivary bacterial microbiome diversity in edentulous infants and their mothers or primary care givers using pyrosequencing. *PLoS One* **6**, e23503.
- Edgar RC (2004) MUSCLE: a multiple sequence alignment method with reduced time and space complexity. *BMC Bioinformatics* **5**, 113.
- Thompson JD, Gibson TJ, Plewniak F, *et al.* (1997) The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* **25**, 4876–4882.
- Gontcharova V, Youn E, Wolcott RD, *et al.* (2010) Black box chimera check (B2C2): a windows-based software for batch depletion of chimeras from bacterial 16S rRNA gene datasets. *Open Microbiol J* **4**, 47–52.
- Smith DM, Snow DE, Rees E, *et al.* (2010) Evaluation of the bacterial diversity of pressure ulcers using bTEFAP pyrosequencing. *BMC Med Genomics* **3**, 41.
- Kushner RF & Doerfler B (2008) Low-carbohydrate, high-protein diets revisited. *Curr Opin Gastroenterol* **24**, 198–203.
- Cebra JJ (1999) Influences of microbiota on intestinal immune system development. *Am J Clin Nutr* **69**, 1046S–1051S.
- Ritchie LE, Burke KF, Garcia-Mazcorro JF, *et al.* (2010) Characterization of fecal microbiota in cats using universal 16S rRNA gene and group-specific primers for *Lactobacillus* and *Bifidobacterium* spp. *Vet Microbiol* **144**, 140–146.
- Middelbos IS, Vester Boler BM, Qu A, *et al.* (2010) Phylogenetic characterization of fecal microbial communities of dogs fed diets with or without supplemental dietary fiber using 454 pyrosequencing. *PLoS One* **5**, e9768.
- Thompson-Chagoyan OC, Maldonado J & Gil A (2007) Colonization and impact of disease and other factors on intestinal microbiota. *Dig Dis Sci* **52**, 2069–2077.
- Inness VL, McCartney AL, Khoo C, *et al.* (2007) Molecular characterisation of the gut microflora of healthy and inflammatory bowel disease cats using fluorescence *in situ* hybridisation with special reference to *Desulfovibrio* spp. *J Anim Physiol Anim Nutr (Berl)* **91**, 48–53.
- Magée EA, Richardson CJ, Hughes R, *et al.* (2000) Contribution of dietary protein to sulfide production in the large intestine: an *in vitro* and a controlled feeding study in humans. *Am J Clin Nutr* **72**, 1488–1494.
- Gill CI & Rowland IR (2002) Diet and cancer: assessing the risk. *Br J Nutr* **88**, S73–S87.
- Norat T & Riboli E (2001) Meat consumption and colorectal cancer: a review of epidemiologic evidence. *Nutr Rev* **59**, 37–47.
- Ramakrishna BS, Roberts-Thomas IC, Pannall PR, *et al.* (1991) Impaired sulphation of phenol by the colonic mucosa in quiescent and active colitis. *Gut* **32**, 46–49.
- Loesche WJ & Gibbons RJ (1968) Amino acid fermentation by *Fusobacterium nucleatum*. *Arch Oral Biol* **3**, 191–202.
- Potrykus J, White RL & Bearne SL (2008) Proteomic investigation of amino acid catabolism in the indigenous gut anaerobe *Fusobacterium varium*. *Proteomics* **8**, 2691–2703.

31. Swidsinski A, Dorffel Y, Loening-Baucke V, *et al.* (2011) Acute appendicitis is characterised by local invasion with *Fusobacterium nucleatum/necrophorum*. *Gut* **60**, 34–40.
32. Bennett KW & Eley A (1993) Fusobacteria: new taxonomy and related diseases. *J Med Microbiol* **39**, 246–254.
33. Suchodolski JS (2011) Intestinal microbiota of dogs and cats: a bigger world than we thought. *Vet Clin North Am Small Anim Pract* **41**, 261–272.
34. Kitahara M, Takamine F, Imamura T, *et al.* (2001) *Clostridium biranonis* sp. nov., a human intestinal bacterium with bile acid 7 alpha-dehydroxylating activity. *Int J Syst Evol Microbiol* **51**, 39–44.
35. Reddy BS, Simi B, Patel N, *et al.* (1996) Effect of amount and types of dietary fat on intestinal bacterial 7 alpha-dehydroxylase and phosphatidylinositol-specific phospholipase C and colonic mucosal diacylglycerol kinase and PKC activities during stages of colon tumor promotion. *Cancer Res* **56**, 2314–2320.
36. Elsayed S & Zhang K (2004) Human infection caused by *Clostridium batthewayi*. *Emerg Infect Dis* **10**, 1950–1952.
37. Woo PC, Lau SK, Woo GK, *et al.* (2004) Bacteremia due to *Clostridium batthewayi* in a patient with acute appendicitis. *J Clin Microbiol* **42**, 5947–5949.
38. Warren YA, Tyrrell KL, Citron DM, *et al.* (2006) *Clostridium aldenense* sp. nov. and *Clostridium citroniae* sp. nov. isolated from human clinical infections. *J Clin Microbiol* **44**, 2416–2422.
39. Louis P, Duncan SH, McCrae SI, *et al.* (2004) Restricted distribution of the butyrate kinase pathway among butyrate-producing bacteria from the human colon. *J Bacteriol* **186**, 2099–2106.
40. Freier TA, Beitz DC, Li L, *et al.* (1994) Characterization of *Eubacterium coprostanoligenes* sp. nov., a cholesterol-reducing anaerobe. *Int J Syst Bacteriol* **44**, 137–142.
41. Hashizume K, Tsukahara T, Yamada K, *et al.* (2003) *Megasphaera elsdenii* JCM1772T normalizes hyperlactate production in the large intestine of fructooligosaccharide-fed rats by stimulating butyrate production. *J Nutr* **133**, 3187–3190.
42. Yoshida Y, Tsukahara T & Ushida K (2009) Oral administration of *Lactobacillus plantarum* Lq80 and *Megasphaera elsdenii* iNP-001 induces efficient recovery from mucosal atrophy in the small and the large intestines of weaning piglets. *Anim Sci J* **80**, 709–715.
43. Su Y, Yao W, Perez-Gutierrez ON, *et al.* (2008) Changes in abundance of *Lactobacillus* spp. and *Streptococcus suis* in the stomach, jejunum and ileum of piglets after weaning. *FEMS Microbiol Ecol* **66**, 546–555.
44. Konstantinov SR, Awati AA, Williams BA, *et al.* (2006) Post-natal development of the porcine microbiota composition and activities. *Environ Microbiol* **8**, 1191–1199.
45. Scharek L, Guth J, Reiter K, *et al.* (2005) Influence of a probiotic *Enterococcus faecium* strain on development of the immune system of sows and piglets. *Vet Immunol Immunopathol* **105**, 151–161.
46. Hacin B, Rogelj I & Matijasic BB (2008) *Lactobacillus* isolates from weaned piglets' mucosa with inhibitory activity against common porcine pathogens. *Folia Microbiol (Praba)* **53**, 569–576.
47. Spencer RJ & Chesson A (1994) The effect of *Lactobacillus* spp. on the attachment of enterotoxigenic *Escherichia coli* to isolated porcine enterocytes. *J Appl Bacteriol* **77**, 215–220.
48. Siggers RH, Siggers J, Boye M, *et al.* (2008) Early administration of probiotics alters bacterial colonization and limits diet-induced gut dysfunction and severity of necrotizing enterocolitis in preterm pigs. *J Nutr* **138**, 1437–1444.
49. Banks WA (2001) Leptin transport across the blood-brain barrier: implications for the cause and treatment of obesity. *Curr Pharm Des* **7**, 125–133.
50. Pereira DI & Gibson GR (2002) Effects of consumption of probiotics and prebiotics on serum lipid levels in humans. *Crit Rev Biochem Mol Biol* **37**, 259–281.