

Background: Psychotropic drugs including some of the new generation of antidepressants and antipsychotics can have important effects on the cardiovascular system including changes in blood pressure and effects on the QTc interval. It is good practice to check ECG and blood pressure (BP) before the administration of certain psychotropic agents. It has been suggested that Psychiatrists should be able to interpret ECG's.

Aims and Method: The aim of our study was to assess the facilities available in the Psychiatric clinic to check blood pressure and arrange ECG's. We were also interested to find out whether psychiatrists were confident in interpreting ECG's and clarify any associated training requirements. So, we carried out an anonymous postal survey of 260 consultant psychiatrists in the North West of England. Data were analysed with the Statistical Package of Social Sciences (SPSS) version 13 for windows.

Results: 132 consultants returned the completed questionnaires giving a response rate of 50.7%. A majority of respondents (59%) felt that it was difficult to arrange for ECG in the clinic and worryingly an even higher percentage (61.4%) lacked facilities to check blood pressure. Only a small minority (12.9%) felt confident about identifying QT prolongation on ECG. An overwhelming percentage of respondents (81.8%) respondents felt that doctors working in psychiatry should have regular training in interpreting ECG's.

Conclusions: This survey highlights the lack of facilities in mental health clinics to check blood pressure and arrange simple medical procedures like ECG. It also highlights the need for regular ECG training for psychiatrists

P218

Coercion and antipsychotic medication for voluntary out-patients: Depot versus oral

M.X. Patel, N. De Zoysa, M. Bernadt, J. Bindman, A.S. David. *Division of Psychological Medicine, Institute of Psychiatry, London, United Kingdom*

Background: Some clinicians consider depot antipsychotics to be stigmatizing and coercive. Former coercion studies have predominantly considered hospital admission rather than medication. This cross-sectional study investigated patients' perspectives of coercion for depot and oral antipsychotics.

Methods: 72 participants, with schizophrenia or schizoaffective disorder on voluntary maintenance medication were randomly selected for further in-depth interviews as a sub-sample from an antipsychotic attitudinal study. The MacArthur Admission Experience (short form) was adapted to explore coercion regarding medication. Scores were compared for formulation groups (depot versus oral).

Results: Only 9 (12.5%) had no concerns about coercion. Coercion scores were higher for depot than oral in terms of total score (mean 4.39 vs 2.80, $p=0.027$), perceived coercion (2.52 vs 1.73, $p=0.041$) and negative pressures subscales (1.17 vs 0.33, $p=0.009$). No significant differences were found for the "voice" subscale (0.70 vs 0.73) and affective reactions. Specifically, more participants on depot felt that people try to force them to take medication (30% vs 2%, $p<0.001$).

Conclusions: To our knowledge, this study is unique in that it reports specifically on coercion regarding both depot and oral antipsychotics, using systematic quantitative methodology. Participants felt that treatment with depots was more coercive than with oral antipsychotics and was associated with a relative lack of true autonomy. One reason for this might be that depots are "given" rather than "taken"; thus the "power of others" may be seen as more potent. Greater

perceived coercion may explain why some consider depots to be a more stigmatizing form of treatment.

P219

Depression - Causes and remedies

M. Popa. *Ecology University, Bucharest, Romania*

The events of life are present in all forms of depression. The major cause of the events who determine depression is the behavior of the others, whether or not it is a reaction to the subject's behavior. In parallel with biological cures and psychotherapy interventions, the author uses a form of re-socialization by religious education, with three objectives: 1) the initiation in the study of human behavior, insisting on the instincts and their actions; 2) The Decalogue- the first step of the greatest importance towards the education of the instincts 3) The Christian belief- the human aspiration to perfection, a maximum of (re)socialization of the human being. Human ontogenesis repeats the evolution of Humanity, but not everyone becomes an adult. Immaturity with moral retardation affects millions of ours contemporary.

Keywords: DEPRESSION, DECALOGUE, CHRISTIANITY, MORAL RETARDATION.

P220

Ziprasidone in hospitalized patients with schizophrenia: Evidence supporting rapid dose titration

F. Rappard, D. Vanderburg, L. Warrington, R. Yang. *Pfizer Inc., New York, NY, USA*

Optimal dosing of psychotherapeutic agents has implications for both symptom control and patient compliance. Trials of ziprasidone in bipolar mania and schizophrenia suggest a target dose of 120-160 mg/d and that rapid titration to this level provides maximum symptom improvement. In this report, data from 2 similarly designed fixed-dose placebo-controlled studies of ziprasidone (rapidly titrated to target doses of 40, 80, 120, or 160 mg/d) in patients with acute schizophrenia were pooled. 369 patients received ziprasidone and 171 patients received placebo. Efficacy was assessed using PANSS at Weeks 1 and 6 (LOCF endpoint) of treatment. Tolerability was assessed by discontinuations (all-cause and due to adverse events). There was a significant linear dose-response relationship between ziprasidone dose and PANSS total score ($F = 12.32$, $P \leq 0.001$). All ziprasidone doses produced statistically significant improvement in PANSS total score; the largest effect size (0.52) was observed for the 160 mg/d group. At Week 6, least-squares mean PANSS total score decreases from baseline were 9.98, 9.54, 11.71, and 14.87 in 40, 80, 120, and 160 mg/d groups, respectively. The corresponding placebo decrease was 2.79. At Week 1, decreases from baseline were 6.18, 5.70, 7.80, and 8.96 in 40, 80, 120, and 160 mg/d groups, respectively. The corresponding placebo decrease was 0.84. Tolerability of ziprasidone 160 mg/d (all-cause/AE discontinuations at week 6: 22%/15% versus 35%/0% for placebo) was comparable with that of lower doses. Rapid titration of ziprasidone to 160 mg/d was associated with greater efficacy compared with lower doses and was well tolerated.

P221

Use of long lasting risperidone in hospitalized patients

M.N. Rodriguez, F. Romero Marmaneu, I. Zaera Cuadrado, C. Ramos Vidal, M. Villar Garcia. *Unidad de Enfermos Agudos, Consorcio Hospitalario de Castellón, Castellón de la Plana, Spain*