

Effects of vitamin D on calcium regulation in vitamin-D-deficient pigs given a phytate-phosphorus diet

BY A. POINTILLART AND NICOLE FONTAINE

Station de Recherches de Nutrition, INRA, 78 350 Jouy-en-Josas, France

AND MONIQUE THOMASSET

INSERM, Unité 120, 78 110 Le Vesinet, France

(Received 23 January 1986 – Accepted 26 June 1986)

1. Vitamin-D-deficient pigs were fed on a phytate-phosphorus diet and treated with vitamin D₃ (+D) to examine the time-course of adaptative changes in plasma minerals, vitamin D metabolites, parathyroid hormone (PTH) and calcium balance and intestinal Ca-binding protein (CaBP).
2. The 5-week vitamin D repletion (25 µg cholecalciferol/kg diet) regimen restored plasma Ca, P and alkaline phosphatase (EC 3.1.3.1) to normal, decreased PTH and markedly and rapidly increased plasma 25-hydroxycholecalciferol (25-OHD, sevenfold after 4 d) and 1,25-dihydroxycholecalciferol (1, 25(OH)₂D₃, 1.8-fold after 4 d).
3. CaBP concentrations were markedly elevated all along the digestive tract, especially in the distal regions.
4. Ca absorption and retention were enhanced (fourfold and sixfold respectively) by the +D diet.
5. The improved Ca absorption, coupled with increased CaBP and 1,25(OH)₂D₃ levels, suggest that vitamin D metabolism in phytate-P-fed pigs is sensitive to the depressed Ca availability due to phytate feeding. It also indicates that CaBP may play an important role in the adaptation of Ca absorption.
6. Persistent hypercalciuria indicates that mineral metabolism was still affected by the phytate nature of the dietary P in spite of the vitamin D treatment.

Phosphorus depletion induces increases in circulating 1,25-dihydroxycholecalciferol (1,25(OH)₂D₃), intestinal calcium-binding protein (CaBP) and Ca absorption (for review, see Sommerville *et al.* 1985*b*). Phytate-P, which is the predominant P source in pig feeds, leads to P deficiency if no additional inorganic P is provided (Pointillart *et al.* 1984, 1985), and phytate-P absorption is improved by vitamin D supplementation (Fontaine *et al.* 1985) demonstrating the importance of this form of P for pig husbandry. Diets in which the bulk of the phosphate is in the form of phytate also give rise to major disturbances of Ca metabolism (hypocalcaemia, hypercalciuria) and cause significant reductions in the utilization of Ca by the formation of poorly absorbed insoluble Ca phytate (Pointillart *et al.* 1985). These disturbances are aggravated by vitamin D deficiency (Pointillart *et al.* 1985). In the present study, we have used vitamin-D-deficient pigs to examine the capacity of vitamin D repletion to overcome the disturbances of Ca metabolism caused by consumption of a diet whose phosphate content, while normal, is largely (80%) in the form of phytate.

We have also examined the changes in the principal plasma variables linked to plasma phosphate and Ca regulation, 25-hydroxycholecalciferol (25-OHD), 1,25(OH)₂D₃, alkaline phosphatase (EC 3.1.3.1; AP) and parathyroid hormone (PTH) levels, throughout the experimental period. They were measured before and during the period of vitamin D supplementation and compared with those of non-supplemented vitamin-D-deficient pigs. At the end of the experimental period the pigs were killed and the CaBP contents of a number of areas of the small intestine were measured to determine what, if any, correlation existed between the values and the Ca absorption measured immediately before death.

Table 1. *Composition of experimental diets (g/kg)*

Dehulled oats*	564
Soya-bean meal 50*	200
Rapeseed meal*	200
Molasses†	20
Calcium carbonate	10
Sodium chloride	4
Mineral-vitamin mix‡	1.5
Calcium	5.9
Total phosphorus	5.8
Phytate-P	4.6

* From INRA (78280 Guyancourt, France).

† Sugar-beet.

‡ Providing (mg/kg diet): 125 Cu as $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 150 Zn as $\text{ZnSO}_4 \cdot \text{H}_2\text{O}$, 80 Fe as $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 30 Mn as $\text{MnSO}_4 \cdot \text{H}_2\text{O}$, 2 Co as $\text{CoSO}_4 \cdot 7\text{H}_2\text{O}$, 0.6 I as KI; vitamins (/kg diet): vitamin A 1500 μg as retinyl palmitate, vitamin E 20 mg as α -tocopheryl acetate, thiamin 2 mg as thiamin hydrochloride, pyridoxine 10 mg as pyridoxine hydroxide, riboflavin 10 mg, ascorbic acid 20 mg, cyanocobalamin 50 μg , Ca pantothenate 40 mg, choline chloride 750 mg, niacin 20 mg, vitamin D₃: no supplementation except +D group 25 μg as cholecalciferol.

MATERIAL AND METHODS

Animals and diets

Fourteen newly-weaned 4-week-old Large White pigs, weighing 7 (SE 0.2) kg, were placed in an u.v.-light-free environment and fed on a standard weaning diet for 2 months. This diet, as previously described (Fontaine *et al.* 1985), contained no vitamin D supplement. The pigs were then placed on the experimental diet (Table 1) for a period of 2 weeks. Neither vitamin D nor inorganic P were added to the diet, which contained 6 g Ca and 6 g P (of which 800 mg/g was phytate-P)/kg. At the end of this period the pigs were divided into two groups (seven pigs per group). One group was fed on the experimental diet without vitamin D (–D) and the other was given the same diet supplemented with 25 μg vitamin D₃/kg (+D). Animals were killed after 5 weeks on these (+D or –D) diets. Blood samples were taken at time 0, i.e. when first placed on the experimental diet, and 4 d and 2 and 5 weeks after the beginning of the vitamin D supplementation. Thus, the initial 2-week feeding of the unsupplemented diet was the phytate-feeding period and the following 5 week period was the vitamin-D-treatment period.

Measurements

Intestinal CaBP. At slaughter, segments (200 mm) of proximal and distal duodenum, distal ileum, mid-jejunum, proximal caecum and colon were excised, everted and washed in cold saline (9 g sodium chloride/l). Mucosal tissue was scraped off in an ice-bath and the CaBP contents determined using a radioimmunoassay employing specific antibodies to purified intestinal pig CaBP as previously described (Pointillart *et al.* 1984). Maximum sensitivity was 1 ng CaBP per assay tube. The results were expressed as μg CaBP/mg soluble mucosal protein (100000 g supernatant fraction).

Vitamin D metabolites. Plasma vitamin D metabolites, 25-OHD and 1,25(OH)₂D were extracted from 1–2 ml plasma samples and separated by a modification of the method of Adams *et al.* (1981). This modification consisted of ethanol–diethyl ether–hexane (2:5:2, by vol.) lipid extraction instead of using methanol–methylene chloride to obtain an upper organic phase. After elution on Sep-Pak cartridge, fraction II (25-OHD) was measured by a competitive protein-binding assay (Preece *et al.* 1974) and fraction III (1,25(OH)₂D) subjected to high-performance liquid chromatography (Waters Microporasil column,

300 × 3.9 mm, eluted with hexane–isopropanol (90:10, v/v; flow-rate 2 ml/min). The fraction containing 1,25(OH)₂D was evaporated under nitrogen and radioimmunoassay performed using the Buhlman Laboratories' (CH 4024 Basel, Switzerland) kit, based on the method of Bouillon *et al.* (1977). The sensitivities were 1.25 pmol for 25-OHD and 0.01 pmol for 1,25(OH)₂D per ml plasma. Variabilities did not exceed 15% for either assay. As the diet contained vitamin D₃ almost exclusively, plasma 25-OHD and 1,25(OH)₂D should be considered as 25-hydroxycholecalciferol and 1,25-dihydroxycholecalciferol respectively.

Plasma indices. Plasma Ca was measured by atomic absorption spectrophotometry, phosphate was determined as inorganic P by the method of Chen *et al.* (1956). Plasma immunoreactive PTH levels were measured with PTH-RIA 200 kits (CEA, ORIS, BP 21 Gif, France) using an antibody directed against the carboxyl terminal of bovine PTH. The lower limit of detectability was 0.37 µg/l plasma. Variability did not exceed 15%. Results are expressed as µg equivalent bovine PTH/l plasma. AP activity was analysed according to Bessey *et al.* (1946) and expressed as Bessey-Lowry units/l plasma (one unit corresponding to 1 mmol *p*-nitrophenyl phosphate hydrolysed/h per l at 37°).

Ca balance. A 10 d Ca balance was carried out on four pigs of each group over the 2 weeks just before slaughter. Pigs were individually housed in appropriate cages.

Statistical analysis. Student's *t* test was used to compare two means, studentized range *Q* test (Newman–Keuls including Chi² test) for multiple-mean comparison and in some cases paired *t* test for plasma value increases or decreases (Snedecor & Cochran, 1971).

RESULTS

Plasma kinetics (Table 2)

Inorganic P. Plasma P levels decreased in both groups until 4 d after the animals had been placed on the experimental (+D or –D) diets. However, this decrease was less marked in +D pigs since, at day 4, the plasma P level of this group was significantly higher than that in the –D group. Thereafter, plasma P significantly increased in +D pigs, reaching a maximum by 2 weeks with no further significant increase, while it remained low in –D pigs. The difference between –D and +D values increased throughout the experimental period. The values for +D pigs were 30, 40 and 50% greater than the corresponding –D values on day 4 and weeks 2 and 5 respectively.

Ca. Plasma Ca concentrations initially decreased in both groups of animals until vitamin D supplementation. They then increased throughout the experiment in the +D group while remaining low and stable in –D group. The value at slaughter did not differ from the initial value in the +D pigs. Thus, vitamin D supplementation restored calcaemia to normal.

AP. Plasma AP activity significantly decreased after 2 weeks on the vitamin-D-supplemented diet, but remained high in –D pigs. At slaughter, –D AP values were threefold higher than the corresponding +D values and approximately twice that of the 2-week –D values.

Vitamin D metabolites. Plasma 25-OHD levels increased (seven times the initial value) rapidly and markedly after 4 d of vitamin D supplementation with no further significant increase. They significantly decrease (by 77% of initial value on day 4) to reach negligible values in –D pigs. The level at slaughter was below the detection limit in many of the –D animals, thus confirming the vitamin-D-deficient status of the –D group.

Plasma 1,25(OH)₂D₃ increased progressively in +D pigs since maximal values were observed by 2 weeks. The percentage initial value increases were respectively 80, 180 and 170 (rounded values) by day 4 and weeks 2 and 5 following feeding of the +D diet. Thus, these increments were relatively less striking than those obtained for 25-OHD: sevenfold,

Table 2. Plasma calcium, inorganic phosphorus, alkaline phosphatase (EC 3.1.3.1; AP; Bessey-Lowry units/l) vitamin D metabolites and parathyroid hormone (PTH) of pigs fed on a phytate-P diet with (+D) or without (-D) vitamin D₃

(Results are expressed as means with their standard errors for fourteen (2 weeks before starting vitamin D supplementation) or seven (4 d, 2 and 5 weeks after starting vitamin D supplementation) pigs per group)

Vitamin D supplementation...	2 weeks before (time 0)	4 d after		2 weeks after		5 weeks after	
		+D	-D	+D	-D	+D	-D
Ca (mmol/l)							
Mean	3.0	2.5†	2.2***†	2.7†‡	2.1***†	2.9†	2.0***†
SE	0.04	0.07	0.04	0.05	0.07	0.08	0.04
P (mmol/l)							
Mean	1.9	1.4†	1.1***†	2.2†‡	1.6**	2.4†	1.6**
SE	0.1	0.1	0.1	0.2	0.2	0.2	0.1
AP (units/l)							
Mean	6.3	5.1	5.2	2.6†‡	4.3***†	2.9†	9.1***†‡
SE	0.2	0.4	0.5	0.2	0.6	0.2	1.0
25-OHD (nmol/l)							
Mean	4.3	29.8†	1.0***†	38.0†	1.3***†	42.2†	0.05***†
SE	0.3	18.8	0.5	6.0	0.8	10.3	0.2
1,25(OH) ₂ D ₃ (pmol/l)							
Mean	47.6	85.0†	15.6***†	132.5†‡	15.9***†	126.6†	13.0***†
SE	2.6	3.5	2.2	14.9	2.9	12.5	1.7
PTH (µg/l)							
Mean	1.1	0.8†	1.1**	0.9	1.4**	0.8†	1.7**
SE	0.1	0.05	0.1	0.05	0.1	0.1	0.3

25-OHD, 25-hydroxycholecalciferol; 1,25(OH)₂D₃, 1,25-dihydroxycholecalciferol.

** Mean values for +D were significantly different ($P < 0.01$) from those for -D at the same time period (Student's t test).

†‡ Mean values were significantly different ($P < 0.05$) from those for † initial (time 0) values or ‡ the immediately preceding value within the same group (i.e. +D or -D). For Ca, inorganic P and AP, multiple mean comparison (MMC) test was used. For 25-OHD, 1,25(OH)₂D₃ and PTH, chi² values prevented the use of the MMC test, therefore paired t test was used.




Fig. 1. Relation between plasma calcium and parathyroid hormone (PTH) of pigs fed on a phytate-phosphorus diet with (+D; Δ) or without (-D; \bullet) vitamin D₃. $y = 3.17 - 0.86x$, $r = -0.72$, $P < 0.01$.

ninefold and tenfold respectively by day 4 and weeks 2 and 5. In -D pigs, plasma 1,25(OH)₂D₃ fell significantly by day 4 with no further significant decrease.

PTH. The experimental +D and -D diets significantly modified plasma PTH levels. They were elevated in -D pigs and lowered in +D pigs. Compared with initial values, the first significant decrease occurred by day 4 with no further decrease in +D pigs, while




Fig. 2. Intestinal calcium-binding protein (CaBP) concentrations of pigs fed on a phytate-phosphorus diet with (+D; ▨) or without (-D ■) vitamin D₃. Results are expressed as means with their standard errors represented by vertical bars from seven pigs per group. ** $P < 0.01$, +D v. -D, Student's t test.




Fig. 3. Relation between pooled duodenal (proximal ○, □; distal ●, ■) calcium-binding protein (CaBP) and plasma 1,25-dihydroxycholecalciferol (1,25(OH)₂D₃) in pigs fed on a phytate-phosphorus diet with (+D; □, ■) or without (-D; ○, ●) vitamin D₃. Two samples were unavailable for assay in each of the -D groups and the +D distal duodenum, one from the +D proximal duodenum was not assayed. $y = 0.12x + 15.4$, $r = 0.76$, $P < 0.01$.

in -D pigs, increases appeared by weeks 2 and 5 but these were not significant. However, compared with +D values, plasma PTH levels were systematically and significantly greater in -D pigs with a rather important twofold difference at slaughter. In addition, a significant negative relation between plasma Ca and PTH was found in both groups for the values over the final 5-week period (Fig. 1).

Ca and CaBP

CaBP (Fig. 2). Mucosal CaBP contents were markedly increased by vitamin D supplementation along the whole intestine. The relative effect was greater in the jejunum (twelfold),




Fig. 4. Calcium balance in pigs fed on a phytate-phosphorus diet with (+D; ▨) or without (-D; ▩) vitamin D₃. Results are expressed as means with their standard errors represented by vertical bars from four pigs per group. ** $P < 0.01$, +D *v.* -D, Student's *t* test.




Fig. 5. Relation between proximal duodenal calcium-binding protein (CaBP) and apparently absorbed Ca in pigs fed on a phytate-phosphorus diet with (+D; Δ) or without (-D; \bullet) vitamin D₃. $y = 0.13x - 0.98$, $r = 0.96$, $P < 0.01$.

ileum (fourfold), caecum (sevenfold) and colon (fourfold) than in the proximal or distal duodenum (twofold, +D *v.* -D). Proximal and distal duodenal CaBP contents were also much greater than those of the more distal segments whatever the diet. The jejunal values were greater than those of the more distal intestinal segments in +D pigs only and they were not negligible when compared with duodenal values (jejunum 12.5 (SE 1.2) *v.* distal duodenum 37.1 (SE 6) $\mu\text{g}/\text{mg}$ protein). This was not the case in -D pigs where most of the intestinal values, with the exception of the duodenum, were negligible (from 0.5 to 1 $\mu\text{g}/\text{mg}$ protein). A significant relation between plasma 1,25(OH)₂D₃ (slaughter values) and duodenal CaBP (both segments pooled) was also found for both groups of pigs (Fig. 3).

Ca balance (Fig. 4). The vitamin D supplementation dramatically improved apparent Ca absorption and retention (expressed as a percentage of intake or daily amount). These were about four and six times greater respectively, than those observed in non-supplemented pigs. Calciuria was high and comparable for both groups when expressed as daily amounts (-D 0.4 (SE 0.1) *v.* +D 0.6 (SE 0.2) g/d) or percentage intake (-D 6 (SE 1.7), +D 9.5 (SE 3);

$P > 0.10$). However, when expressed as relative to apparently absorbed Ca, it was greater in $-D$ than in $+D$ pigs ($-D$ 0.37 (SE 0.01) *v.* $+D$ 0.18 (SE 0.07); $P < 0.05$). A significant and close relation was found between proximal duodenal CaBP content and apparently absorbed Ca for both groups (Fig. 5).

DISCUSSION

Vitamin D treatment produced a twofold increase (4 d *v.* initial values) in circulating $1,25(\text{OH})_2\text{D}_3$ within 4 d. At the end of the same period (4 d) there was also a significant difference between the plasma P of $+D$ and $-D$ pigs. This could be interpreted as the initial effect of vitamin D on P absorption since this vitamin markedly enhances phytate-P availability in pigs (Fontaine *et al.* 1985). Nevertheless, 2 weeks were necessary for plasma P to become significantly greater than initial values, corresponding to a threefold rise of $1,25(\text{OH})_2\text{D}_3$ (2-week *v.* initial values). Similar results were obtained by Engstrom *et al.* (1984) working on vitamin D-depleted but not P-depleted pigs given approximately the same amount of vitamin D_3 (25 $\mu\text{g}/\text{d}$). In the present study the plasma P of $+D$ pigs increased from 60 to 75 mg/l (from 1.9 to 2.4 mmol/l) while that in the vitamin-D-fed pigs in the study of Engstrom *et al.* (1984) increased from 59 to 78 mg/l over a similar period. The vitamin-D-deficient pigs studied by Engstrom *et al.* (1984) developed hyperphosphataemia while the vitamin-D-deficient pigs in the present study became hypophosphataemic (Table 2) although they received the same amounts of total P (5–6 g/kg). The difference is probably the consequence of the two kinds of P given, since Engstrom *et al.* (1984) provided inorganic P (CaHPO_4) instead of phytate-P, which is known to cause hypophosphataemia and other signs of P deficiency (Pointillart *et al.* 1984, 1985; Fontaine *et al.* 1985). Such a P deficiency will stimulate renal $1,25(\text{OH})_2\text{D}$ production via $1-\alpha$ -hydroxylase as demonstrated in P-depleted pigs by Engstrom *et al.* (1985). However, the hypophosphataemia observed by Engstrom *et al.* (1985) (1.2 mmol/l) was much more acute than that of our pigs (1.6 mmol/l at slaughter) probably because their P-deficient diet (0.85 g P/kg diet) caused a greater P deficiency than the phytate-P diet which contained a normal amount of total P (5.8 g/kg). This could also explain the difference between $1,25(\text{OH})_2\text{D}_3$ values (1100 pmol/l) reported by these authors after 6 weeks of P deficiency and our final values for the $+D$ phytate-P diet (Table 2).

The time-course of the vitamin D effect on plasma Ca was much slower than its effect on plasma P, since it returned to its initial value only after 5 weeks. Littledike *et al.* (1976) found a prompt reversal of the hypocalcaemia after 12 d of vitamin D_3 (25 $\mu\text{g}/\text{d}$) supplementation in pigs which were only vitamin-D-deficient. However, phytate-P diets decrease calcaemia (Pointillart *et al.* 1984, 1985) in pigs and thus could contribute to the delayed normalization of plasma Ca. Both hypophosphataemia and hypocalcaemia could also result from simultaneous Ca and P unavailability of phytate-P diets because formation of unavailable Ca-phytate takes place in the digestive tract as indicated by our previous findings on faecal Ca and P (Pointillart *et al.* 1985). Even if the phytate-based diets are responsible for the reduction in digestible P, and consequently for the phosphataemia, vitamin D deficiency itself may, in pigs, cause hyperphosphataemia (Engstrom *et al.* 1984). This would help to explain the tendency for the plasma P levels to rise, often between 4 d and 2 weeks (and remain stable until the end of the experimental period) (Table 2) in pigs given no vitamin D supplement. In these same pigs the extreme vitamin D deficiency, which was confirmed by the low plasma 25-OHD and $1,25(\text{OH})_2\text{D}$ values would make it impossible for the metabolism to be stimulated by lack of P as has been suggested (Anon., 1974) and recently confirmed in pigs (Engstrom *et al.* 1985; Fox & Ross, 1985). We ourselves have confirmed this in our earlier experiments on pigs fed on a diet containing phytate as the major form of P (Pointillart *et al.* 1984, 1985).

Vitamin D₃ supplementation increased mucosal CaBP contents along the whole intestine and dramatically enhanced Ca absorption. Moreover the significant correlations found between duodenal CaBP and apparently absorbed Ca and between duodenal CaBP and circulating 1,25(OH)₂D₃ suggest that vitamin D repletion leads to increased Ca absorption and a parallel increase in intestinal CaBP concentration. These responses are further stimulated by the secondary P deficiency resulting from the high-phytate intake. The elevated vitamin-D-dependent Ca absorption, coupled with increased intestinal CaBP, has also been shown in P-deficient rats (Thomasset *et al.* 1976), birds (Rosenberg *et al.* 1983; Bar & Hurtwitz, 1984; Sommerville *et al.* 1985*a*) and pigs (Fox *et al.* 1978). The results are also in agreement with those previously obtained in pigs fed on phytate-rich diets without vitamin D supplement (Pointillart *et al.* 1985) and with the findings reported for vitamin-D-deficient rats (Wasserman *et al.* 1974) which was confirmed in pigs (Thomasset *et al.* 1979) treated with vitamin D. The increased CaBP contents of distal intestine, especially jejunum, caecum and colon, also demonstrate that these regions are able to respond to 1,25(OH)₂D₃ as was observed recently in rats (Perret *et al.* 1985). There is, thus, considerable support for the hypothesis that intestinal CaBP can play an important role in Ca absorption.

The increased Ca retention by +D pigs can lead to better bone mineralization, as previously described (Fontaine *et al.* 1985). Both vitamin D and P deficiencies may increase plasma AP activity in pigs (Pointillart *et al.* 1984, 1985; Engstrom *et al.* 1985). Thus, simultaneous vitamin D and P deficiencies may have a synergistic effect on plasma AP since its final value in the -D group was 1.5 the initial value and three times the +D group final value.

Plasma PTH concentrations rapidly decreased in the +D group. They were invariably greater in the -D pigs than in the +D pigs. This is in agreement with the PTH rises found in pigs with hereditary vitamin-D-dependent rickets (Wilke *et al.* 1979), with simultaneous P and vitamin D deficiency (Pointillart *et al.* 1985) or with dietary vitamin-D-deficiency alone (Pointillart *et al.* 1978). However, the observation that P deficiency alone decreases plasma PTH in vitamin-D-replete pigs (Sommerville *et al.* 1985*b*) may explain why the PTH increase of the -D group which was also P-deficient, was not significantly different from its initial value. Such an explanation is also supported by the unchanged PTH levels described for pigs fed on hypophosphataemic diets with no dietary vitamin D supplementation but normal circulating 25-OHD (50–125 nmol/l, Pointillart *et al.* 1984) levels as compared with the very low levels observed in -D pigs (< 1 nmol/l). In addition, the significant negative correlation between plasma Ca and PTH levels suggests that hypocalcaemia was probably responsible for the PTH stimulation of the -D pigs, while the absence of true hyperparathyroidism (maximal PTH increase was only 0.6 µg/l, final *v.* initial values) was probably a consequence of the persistent hypophosphataemia resulting from dietary P unavailability. Nevertheless, hypercalciuria persisted in the +D pigs in spite of considerably enhanced Ca retention. Raising the dietary vitamin D of baby pigs from 0 to 25 µg/kg diet increases urinary Ca from 10 to 50 mg/d (Miller *et al.* 1965) but these values are negligible when expressed as a percentage of Ca intake (1%) or compared with the 400 and 600 mg urinary Ca excreted in the growing pigs fed on the phytate diet, with or without vitamin D respectively. Thus, it is suggested that vitamin D supplementation restored normal calcaemia and phosphataemia by increasing mineral absorption although some mineral disorders still persisted. Their persistence could be due to the phytate nature of the dietary P (Pointillart *et al.* 1984, 1985; Fontaine *et al.* 1985) or to the vitamin D treatment being too brief, or both. The second hypothesis is not supported by the normal circulating levels of 1,25(OH)₂D₃ (127–132 *v.* 140–160 pmol/l in normal growing pigs, De Vernejoul *et al.* 1984) and the normal 25-OHD (30–42 nmol/l *v.* normal pigs 55 nmol/l,

Thomasset *et al.* 1979; 32–82 nmol/l, Engstrom *et al.* 1985). Thus, the secondary P deficiency could be an explanation for the hypercalciuria. The rise in urinary Ca, in terms of percentage Ca absorbed, found in the –D pigs supports this hypothesis. This persistent hypercalciuria suggests that mineral metabolism was still affected by the phytic nature of the dietary P in spite of the normalized plasma Ca and P levels obtained with the vitamin D supplementation. This relation is supported by persistent hypophosphaturia (urinary P 50 mg/d, Fontaine *et al.* 1985). Thus vitamin D treatment considerably enhances Ca utilization (the present results) and P utilization (Fontaine *et al.* 1985) of pigs fed on a phytate-P diet but did not completely overcome the negative effects of phytate feeding on mineral metabolism.

REFERENCES

- Adams, J., Clemens, T. L. & Holick, M. F. (1981). *Journal of Chromatography* **226**, 198–201.
- Anon. (1974). *Nutrition Review* **38**, 247–249.
- Bar, A. & Hurtwitz, S. (1984). *Poultry Science* **63**, 1975–1979.
- Bessey, O., Lowry, O. & Brock, M. J. (1946). *Journal of Biological Chemistry* **164**, 321–329.
- Bouillon, R., Van Baelen, H. & De Moor, P. (1977). *Journal of Clinical Endocrinology and Metabolism* **45**, 225–231.
- Chen, P. S., Toribara, T. Y. & Warner, H. (1956). *Analytical Chemistry* **28**, 1756–1758.
- De Vernejoul, M. C., Pointillart, A., Cywiner Golenzer, C., Morieux, C., Bielakoff, J., Modrowski, D. & Miravet, L. (1984). *American Journal of Pathology* **116**, 377–384.
- Engstrom, G. W., Horst, R. L., Reinhardt, T. A. & Littledike, E. T. (1984). *Journal of Nutrition* **114**, 119–126.
- Engstrom, G. W., Horst, R. L., Reinhardt, T. A. & Littledike, E. T. (1985). *Journal of Animal Science* **60**, 1005–1010.
- Fontaine, N., Fourdin, A. & Pointillart, A. (1985). *Reproduction, Nutrition, Développement* **25**, 717–727.
- Fox, J., Pickard, D. W., Care, A. D. & Murray, T. M. (1978). *Journal of Endocrinology* **78**, 379–387.
- Fox, J. & Ross, R. (1985). *Journal of Endocrinology* **105**, 169–173.
- Littledike, E., Hughes, M., Haussler, M. & Arnaud, S. (1976). *Proceedings of the International Pig Veterinary Society Congress*, Ames, Iowa.
- Miller, E. R., Ullrey, D. E., Zutaut, C. L., Hoefler, J. A. & Luecke, R. W. (1965). *Journal of Nutrition* **85**, 255–259.
- Perret, C., Desplan, C. & Thomasset, M. (1985). *European Journal of Biochemistry* **150**, 211–217.
- Pointillart, A., Fontaine, N. & Thomasset, M. (1984). *Nutrition Reports International* **29**, 473–483.
- Pointillart, A., Fontaine, N., Thomasset, M. & Jay, M. E. (1985). *Nutrition Reports International* **32**, 155–167.
- Pointillart, A., Garel, J. M. & Guéguen, L. (1978). *Annales de Biologie Animale, Biochimie, Biophysique* **18**, 167–174.
- Preece, M. A., O'Riordan, J. L. H., Lawson, D. E. M. & Kodicek, E. A. (1974). *Clinica Chimica Acta* **54**, 235–242.
- Rosenberg, J., Ben-gal, I., Pines, M., Hurtwitz, S. & Bar, A. (1983). *Nutrition Reports International* **28**, 335–343.
- Sommerville, B. A., Blahos, J., Harvey, S., Chadwick, A. & Spencer, G. S. G. (1985a). *Hormone and Metabolic Research* **17**, 245–250.
- Sommerville, B. A., Maunder, E., Ross, R., Care, A. D. & Brown, R. C. (1985b). *Hormone and Metabolic Research* **17**, 78–81.
- Snedecor, G. & Cochran, W. (1971). *Méthodes Statistiques*, pp. 100–107, 302–307. Paris: ACTA.
- Thomasset, M., Cuisinier-Gleizes, P. & Mathieu, H. (1976). *Biomedicine* **25**, 345–349.
- Thomasset, M., Pointillart, A., Cuisinier-Gleizes, P. & Guéguen, L. (1979). *Annales de Biologie animale, Biochimie, Biophysique* **19**, 769–773.
- Wasserman, R., Taylor, A. & Fullmer, C. (1974). *Biochemical Society Special Publications* **3**, 55–74.
- Wilke, R., Harmeyer, J., von Grabe, C., Hehrmann, R. & Hesch, R. D. (1979). *Acta Endocrinologica* **92**, 295–308.