

## Gender differences in retinol metabolism are independent of $\beta$ -carotene bioconversion

Anthony Oxley<sup>1</sup>, Philip Berry<sup>2</sup>, John Hesketh<sup>3</sup>, Alan V Boddy<sup>2</sup> and Georg Lietz<sup>1</sup>

<sup>1</sup>Human Nutrition Research Centre, School of Agriculture, Food and Rural Development, <sup>2</sup>Northern Institute for Cancer Research and <sup>3</sup>Institute for Cell and Molecular Biosciences; Newcastle University, UK

Gender-related differences in retinol and provitamin A carotenoid metabolism have previously been detected, with higher retinol concentrations in men, and on the contrary, higher  $\beta$ -carotene concentrations in women<sup>(1)</sup>. This reversed relationship suggests a higher  $\beta$ -carotene to retinol conversion rate in males<sup>(1)</sup>. Furthermore, lower conversion efficiency is correlated with BMI in women, but not in men<sup>(2)</sup>. We investigated retinol metabolism and  $\beta$ -carotene conversion efficiency in 23 female and 19 male volunteers by co-administering 2 mg [<sup>13</sup>C<sub>10</sub>]- $\beta$ -carotene and 1 mg [<sup>13</sup>C<sub>10</sub>]-retinyl acetate. Relative absorption of  $\beta$ -carotene in the first 24 hours represented 11% of total ingested [<sup>13</sup>C<sub>10</sub>]- $\beta$ -carotene, with an inter-individual coefficient of variation of 49%. [<sup>13</sup>C<sub>10</sub>]- $\beta$ -carotene plasma concentration within the first 24 hours post-dose were significantly inversely related to the ability to convert  $\beta$ -carotene into retinyl-palmitate ( $r = -0.89$ ;  $p < 0.001$ ). More importantly, significantly higher plasma concentrations of preformed [<sup>13</sup>C<sub>10</sub>]-retinol and bio-converted [<sup>13</sup>C<sub>5</sub>]-retinol were found in men compared to women (Figure 1), although plasma [<sup>13</sup>C<sub>10</sub>]- $\beta$ -carotene concentrations were similar between the genders. Interestingly, differences in retinoid concentrations are independent from the ability to cleave [<sup>13</sup>C<sub>10</sub>]- $\beta$ -carotene, since both retinyl palmitate/ $\beta$ -carotene and the newly-formed retinoid reference dose ratios are not significantly different between men and women (Figure 1).




Fig. 1. Effect of gender on plasma retinol concentrations and provitamin A conversion efficiency (AUC = Area under the curve for the first 24 hours post-dose).

In summary, our isotope dilution technique revealed that men have higher circulating retinol concentrations compared to women, and that this difference is independent of provitamin A conversion efficiency. We hypothesize that a gender specific effect on metabolic flux between different tissues could explain this difference.

This work was supported by the BBSRC.

1. Faure H, Preziosi P *et al.* (2006) *Eur J Clin Nutr* **60**(6), 706–17.
2. Tang G, Qin J *et al.* (2003) *Am J Clin Nutr* **78**(2), 259–66.