

Objective To characterize conflict records made by students, teachers/others and parents/guardians in a Brazilian high school institution.

Method Qualitative research, exploratory documental. Studied 113 records from 2014 to 2016.

Results “Indiscipline and Incivility”–75 records (66 by teachers/others and 9 by students) of students: improper use of clothes and accessories, cell phone use in class, not bringing material, dating in school, conversations during class, do not respect the timetables, inappropriate jokes, disrupting activities, theft of materials and disrespect toward authority figures. “School violence”–22 records (12 by students, 9 by teachers/others and 1 by parent/guardian) of school violence: physical violence between students, psychological/verbal between students/teachers/others, and a match of sexual abuse. Three records (by teachers/others) of violence against the school: students destroyed teaching and cleaning materials and caused damage to the patrimony. Fifteen records (12 by students, 2 by parents/guardians and 1 by teacher) of school’s violence: teachers’ harassment (excessive rigor in regard to school performance, clutter in the ratings, refusal to clarify doubts and inappropriate criticism on student’s behavior) and institutional negligence (teachers’ delay, lack of clarification on teaching organization and supervision in practical activities).

Conclusion Some students’ acts of indiscipline and incivility can be protests against the social control of the school. Assistance in case of conflicts and violence as well as preventive measures must be based on interdisciplinary and inter-sectorial articulation practice.

Disclosure of interest The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.01.2217>

EW0348

Mothers mental health of children with attention deficit/hyperactivity disorder (ADHD)

Z. Sepehrmanesh

Kashan University of Medical Sciences, psychiatry department-medical faculty, Kashan, Iran

Background Attention deficit hyperactivity disorder (ADHD) is the most common neuropsychiatric disorder in pediatric populations with an early onset. Mental health of mother can effect on child mental health and relation between mother and child is very important. The aim of this study was investigation about mental health of mothers with children suffering from attention deficit/hyperactivity disorder.

Materials and methods This cross sectional study was carried out on 100 mothers of children with ADHD diagnosis (Kashan, Iran). The data collection instruments included the Goldberg’s General Health Questionnaire, and the questionnaire form includes the personal information and the variables associated to mental health.

Results The total means score of GHQ in the mothers’ was $26/6 \pm 11/78$. Fifty-eight percent of the mothers have mental health problem. The highest mean score in the sub scales was related to the anxiety subscale (7.73) and the most common mental health problem in these mothers was anxiety problems (11%). Mental health problem is most common in mothers with low socio-economic status, younger age than 30 years old, education lower than diploma degree, householder mothers, single, having boy child with ADHD, having child more than 9 years old.

Conclusion According to the results of this study, 58% of the mothers have mental health problem. So with screening and on time diagnosis and treatment, we can prevent disadvantage effects of these problems on social and mental health of their children.

Disclosure of interest The author has not supplied his/her declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.01.2218>

EW0349

Factors associated with depression severity in adolescence

G. Serra^{1,*}, M.P. Casini¹, V. Maggiore¹, R.J. Baldessarini², S. Vicari¹

¹ Pediatric Hospital Bambino Gesù, Department of Neuroscience, Roma, Italy

² Harvard Medical School, Department of Psychiatry, Boston, USA

* Corresponding author.

Introduction Severe depression is greatly impairing during adolescence and involves a high risk for suicidal behaviors.

Objectives and aims Identify clinical and demographic factors associated with severity of depression in adolescents diagnosed with a major mood disorder so as to improve clinical treatment and prevent suicidal behaviors.

Methods We analyzed factors associated with depression severity in 145 severely ill adolescents diagnosed with a major affective disorder using the K-SADS (Kiddie-Schedule for Affective Disorders and Schizophrenia) at the Mood Disorder Outpatient Program of Bambino Gesù Children’s Hospital (Rome). Depressive and manic symptoms were rated with the CDRS-R (Children’s Depression Rating Scale-Revised) and K-SADS-MRS (Mania Rating Scale), respectively. Bivariate comparisons were followed by multivariable linear regression modeling.

Results Depression severity was greater among females than males (mean CDRS scores: 53.0 vs. 42.8; $P < 0.0001$) and with major depressive versus bipolar disorder diagnosis (50.4 vs. 45.4; $P = 0.001$). Manic symptoms, including irritability, mood lability, crowded thoughts, delusions, and insomnia, were more likely with more severe depression; their number and severity correlated with CDRS-R total score (respectively, $\beta = 1.53$ and 5.44 ; both $P < 0.0001$). Factors independently and significantly associated with CDRS-R depression score in multivariate modeling were:

- presence of suicidal ideation;
- absence of ADHD;
- female sex;
- greater number of manic symptoms.

Conclusions Severe depression was associated with manic symptoms and with suicidal ideation among adolescents diagnosed with either bipolar or major depressive disorders. This relationship should be considered in treatment planning and suicide prevention, including consideration of mood-stabilizing and antimanic agents in the treatment of severe adolescent depression.

Disclosure of interest The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.01.2219>

EW0350

Pediatric mania: The controversy between euphoria and irritability

G. Serra^{1,*}, M. Uchida², C. Battaglia¹, M.P. Casini¹, L. De Chiara³, J. Biederman², S. Vicari¹, J. Wozniak²

¹ Pediatric Hospital Bambino Gesù, Department of Neuroscience, Roma, Italy

² Harvard Medical School, Department of Pediatric Psychopharmacology, Boston, USA

³ Harvard Medical School, Department of Psychiatry, Boston, USA

* Corresponding author.

Introduction Pediatric bipolar disorder (BD) is a highly morbid pediatric psychiatric disease, consistently associated with family psychiatric history of mood disorders, with high levels of morbidity and disability and with a great risk of suicide.

Objectives While there is a general consensus on the symptomatology of depression in childhood, the phenomenology of pediatric mania is still highly debated and the course and long-term outcome of pediatric BD still need to be clarified.