



A reproducing kernel approach to Lebesgue decomposition*

Jashan Bal, Robert T.W. Martin, Fouad Naderi

Abstract. We show that properties of pairs of finite, positive and regular Borel measures on the complex unit circle such as domination, absolute continuity and singularity can be completely described in terms of containment and intersection of their reproducing kernel Hilbert spaces of 'Cauchy transforms' in the complex unit disk. This leads to a new construction of the classical Lebesgue decomposition and proof of the Radon–Nikodym theorem using reproducing kernel theory and functional analysis.

Given a finite, positive and regular Borel measure, μ , on the complex unit circle, $\partial\mathbb{D}$, it is natural to consider its L^2 -space, $L^2(\mu) := L^2(\mu, \partial\mathbb{D})$, as well as $H^2(\mu) := \overline{\mathbb{C}[\zeta]}^{\|\cdot\|_{L^2(\mu)}}$, the closure of the analytic polynomials, $\mathbb{C}[\zeta]$, in $L^2(\mu)$. The linear operator of multiplication by the independent variable, M_ζ^μ , is unitary on $L^2(\mu)$ and has $H^2(\mu)$ as a closed invariant subspace so that $Z^\mu := M_\zeta^\mu|_{H^2(\mu)}$ is an isometry that will play a central role in our analysis. The μ -Cauchy transform of any $h \in H^2(\mu)$ is the analytic function,

$$(\mathcal{E}_\mu h)(z) := \int_{\partial\mathbb{D}} \frac{h(\zeta)}{1 - z\bar{\zeta}} \mu(d\zeta) \in \mathcal{O}(\mathbb{D}),$$

in the complex unit disk, $\mathbb{D} := (\mathbb{C})_1$. Here, given a Banach space, X , $(X)_1$ and $[X]_1$ denote its open and closed unit balls in the norm topology.

Recall that a reproducing kernel Hilbert space (RKHS), \mathcal{H} , is a Hilbert space of functions on a set, X , so that point evaluation at any point $x \in X$ yields a bounded linear functional on the space. The Riesz representation lemma then implies the existence of *kernel vectors*, $k_x, x \in X$, so that the bounded linear functional of point evaluation at x is implemented by inner products against k_x . The function of two variables, $k : X \times X \rightarrow \mathbb{C}$,

$$k(x, y) := \langle k_x, k_y \rangle_{\mathcal{H}},$$

is then called the *reproducing kernel* of \mathcal{H} . In this paper, all inner products and sesquilinear forms are conjugate linear in their first argument and linear in their second argument. Any reproducing kernel function is a *positive kernel* function on $X \times X$, i.e. for any finite set $\{x_1, \dots, x_n\} \subseteq X$, the $n \times n$ matrix,

$$[k(x_i, x_j)]_{1 \leq i, j \leq n} \geq 0, \tag{0.1}$$

is positive semi-definite. Conversely, by a theorem of Aronszajn and Moore, given any positive kernel function, k , on $X \times X$, one can construct a RKHS of functions on X with reproducing kernel k [3], see Subsection 1.2. Given this bijective correspondence between positive kernel functions on X and RKHS of functions on X , one writes $\mathcal{H} = \mathcal{H}(k)$ if \mathcal{H} is a RKHS with reproducing kernel k .

Equipping the vector space of μ -Cauchy transforms with the $H^2(\mu)$ -inner product,

$$\langle \mathcal{E}_\mu g, \mathcal{E}_\mu h \rangle_\mu := \langle g, h \rangle_{L^2(\mu)}; \quad g, h \in H^2(\mu),$$

yields a reproducing kernel Hilbert space (RKHS) of analytic functions in \mathbb{D} , $\mathcal{H}^+(\mu)$, with reproducing kernel,

$$k^\mu(z, w) = \int_{\partial\mathbb{D}} \frac{1}{1 - z\bar{\zeta}} \frac{1}{1 - w\bar{\zeta}} \mu(d\zeta); \quad z, w \in \mathbb{D}. \tag{0.2}$$

Using the above formula (0.2), it is easy to check that domination of measures implies domination of the reproducing kernels for their spaces of Cauchy transforms,

$$\mu \leq t^2 \lambda, \quad t > 0 \quad \Rightarrow \quad k^\mu \leq t^2 k^\lambda,$$

see Theorem 3.1, where we write $k \leq K$ for positive kernel functions k, K on X , if $K - k$ is a positive kernel function on X . We will say that λ *dominates* μ in the *reproducing kernel sense* (by $t^2 > 0$) and write $\mu \leq_{RK} t^2 \lambda$ to denote that $k^\mu \leq t^2 k^\lambda$. By results of Aronszajn, domination of kernels, $k \leq t^2 K$, is equivalent to bounded containment of their RKHS, i.e. $k \leq t^2 K$ if and only if $\mathcal{H}(k) \subseteq \mathcal{H}(K)$ and the norm of the linear embedding $e : \mathcal{H}(k) \hookrightarrow \mathcal{H}(K)$ is at most $t > 0$ [3]. See Subsection

AMS subject classification: 47B32, 15A63, 28A99.

Keywords: reproducing kernels, Hilbert spaces of analytic functions, Lebesgue decomposition, sesquilinear forms in Hilbert space.

*R.T.W.M. partially supported by NSERC grant 2020-05683

1.2 for a review of RKHS theory and these results. In summary, domination of measures implies bounded containment of their spaces of Cauchy transforms:

$$\mu \leq t^2 \lambda \quad \Rightarrow \quad \mathcal{H}^+(\mu) \subseteq \mathcal{H}^+(\lambda), \quad e_{\mu,\lambda} : \mathcal{H}^+(\mu) \hookrightarrow \mathcal{H}^+(\lambda), \quad \|e_{\mu,\lambda}\| \leq t,$$

i.e. $\mu \leq t^2 \lambda \Rightarrow \mu \leq_{RK} t^2 \lambda$.

Building on this observation, we show that domination and, more generally, absolute continuity, as well as mutual singularity of measures can be completely characterized in terms of their spaces of Cauchy transforms. Moreover, we develop an independent construction of the Lebesgue decomposition and new proof of the Radon–Nikodym theorem using reproducing kernel methods and operator theory.

Outline

The following Background section, Section 1, provides an introduction to (i) the bijective correspondence between positive, finite and regular Borel measures on the circle and contractive analytic functions in the disk, (ii) reproducing kernel theory and (iii) the theory of densely–defined and positive semi-definite quadratic forms in a separable, complex Hilbert space.

Section 2 introduces the reproducing kernel Hilbert spaces, $\mathcal{H}^+(\mu)$, of μ –Cauchy transforms associated to any positive, finite and regular Borel measure, μ , on the complex unit circle. These are Hilbert spaces of holomorphic functions in the complex unit disk.

Our first main results appear in Section 3. Theorem 3.1 proves that domination of positive measures in the reproducing kernel sense is equivalent to domination in the classical sense:

Theorem 3.1 *Given positive, finite and regular Borel measures, μ and λ on the unit circle, $\mu \leq_{RK} t^2 \lambda$ for some $t > 0$ if and only if $\mu \leq t^2 \lambda$.*

This result is extended to general absolute continuity, written $\mu \ll \lambda$, in Theorem 3.12. Namely, we say that μ is absolutely continuous in the reproducing kernel sense with respect to λ , written $\mu \ll_{RK} \lambda$, if the intersection of the space of μ –Cauchy transforms with the space of λ –Cauchy transforms, $\text{int}(\mu, \lambda)$, is norm-dense in $\mathcal{H}^+(\mu)$.

Theorem 3.12 *Let μ, λ be positive, finite and regular Borel measures on $\partial\mathbb{D}$. Then $\mu \ll \lambda$ if and only if $\mu \ll_{RK} \lambda$.*

Moreover, Theorem 3.12 gives a formula for the Radon–Nikodym derivative of μ with respect to λ in terms of the closed, densely–defined embedding, $e_{\mu,\lambda} : \text{int}(\mu, \lambda) := \mathcal{H}^+(\mu) \cap \mathcal{H}^+(\lambda) \subseteq \mathcal{H}^+(\mu) \hookrightarrow \mathcal{H}^+(\lambda)$.

These are satisfying results, however, actual construction of the Lebesgue decomposition of μ with respect to λ using reproducing kernel methods is more subtle and bifurcates into the two cases, where: The intersection space, $\text{int}(\mu, \lambda) = \mathcal{H}^+(\mu) \cap \mathcal{H}^+(\lambda)$, of the spaces of μ and λ –Cauchy transforms is (i) invariant, or, (ii) not invariant, for the image, V^μ , of $Z^\mu = M_\zeta^\mu|_{H^2(\mu)}$ under Cauchy transform. Some necessary and sufficient conditions for this to hold are obtained in Lemma 4.3 and Proposition 4.7. Namely, as described in Subsection 1.1, there is a bijection between contractive analytic functions in the complex unit disk and positive, finite and regular Borel measures on the circle. If a positive measure, μ , corresponds to an extreme point of this compact, convex set of contractive analytic functions, we say that μ is *extreme*, otherwise μ is *non-extreme*. As established in Lemma 4.3 and Proposition 4.7, the intersection space, $\text{int}(\mu, \lambda)$ will be V_μ –reducing if (i) λ is non-extreme or if (ii) $\mu + \lambda$ is extreme, and the intersection space will be non-trivial and not V_μ –invariant if μ, λ are both extreme but $\mu + \lambda$ is non-extreme.

In the positive direction, we obtain:

Theorem 4.5 *Let μ and λ be finite, positive and regular Borel measures on the unit circle. If the intersection space, $\text{int}(\mu, \lambda)$ is V_μ –invariant and $\mu = \mu_{ac} + \mu_s$ is the Lebesgue decomposition of μ with respect to λ , then*

$$\mathcal{H}^+(\mu) = \mathcal{H}^+(\mu_{ac}) \oplus \mathcal{H}^+(\mu_s).$$

In this case,

$$\mathcal{H}^+(\mu_{ac}) = \text{int}(\mu, \lambda)^{-\|\cdot\|_\mu}, \quad \text{and} \quad \mathcal{H}^+(\mu_s) \cap \mathcal{H}^+(\lambda) = \{0\}.$$

Given two positive, finite and regular Borel measures, μ and λ , on the complex unit circle, $\partial\mathbb{D}$, one can associate to μ a densely–defined and positive semi-definite sesquilinear or quadratic form in $H^2(\lambda)$. Namely, we define the *form domain*, $\text{Dom } q_\mu \subseteq H^2(\lambda)$, as the *disk algebra*, $\text{Dom } q_\mu := A(\mathbb{D})$, the unital Banach algebra of all uniformly bounded analytic functions in the unit disk which extend continuously to the boundary, equipped with the supremum norm. The disk algebra embeds isometrically into the continuous functions on the circle, $\mathcal{C}(\partial\mathbb{D})$ and $A(\mathbb{D})$ can be viewed as a dense subspace of

$H^2(\lambda)$. The quadratic form, $q_\mu : \text{Dom } q_\mu \times \text{Dom } q_\mu \rightarrow \mathbb{C}$ is then defined in the obvious way by integration against μ ,

$$q_\mu(g, h) := \int_{\partial\mathbb{D}} \overline{g(\zeta)} h(\zeta) \mu(d\zeta), \quad g, h \in A(\mathbb{D}) = \text{Dom } q_\mu. \quad (0.3)$$

As described in Section 3 and Theorem 3.8, there is a theory of Lebesgue decomposition of densely-defined and positive semi-definite quadratic forms in a Hilbert space, \mathcal{H} . Namely, given any such form, there is a unique *Simon–Lebesgue form decomposition*,

$$q = q_{ac} + q_s,$$

where $0 \leq q_{ac}, q_s \leq q$, q_{ac} is *absolutely continuous* in the sense that it is *closeable* and it is maximal in the sense that q_{ac} is the largest closeable quadratic form bounded above by q . The form q_s is singular in the sense that the only closeable positive semi-definite form it dominates is the identically 0 form. Here, a positive semi-definite quadratic form, q , with dense form domain $\text{Dom } q$ in \mathcal{H} , is *closed*, if $\text{Dom } q$ is a Hilbert space, *i.e.* complete, with respect to the norm induced by the inner product $q(\cdot, \cdot) + \langle \cdot, \cdot \rangle_{\mathcal{H}}$. A form is then *closeable* if it has a closed extension. See Subsection 1.3 for an introduction to the theory of densely-defined and positive semi-definite quadratic forms.

An immediate question is whether the Simon–Lebesgue decomposition of the form, q_μ , in $H^2(\lambda)$ coincides with the Lebesgue decomposition of μ with respect to λ . Namely, if $\mu = \mu_{ac} + \mu_s$ and $q_\mu = q_{ac} + q_s$, then is it true that $q_{ac} = q_{\mu_{ac}}$ and $q_s = q_{\mu_s}$? A complete answer, summarized in the theorem below, is provided in Theorem 4.12, Theorem 4.18, Corollary 4.14 and Corollary 4.15.

Theorem If $q_\mu = q_{ac} + q_s$ is the Simon–Lebesgue form decomposition of q_μ in $H^2(\lambda)$, then

$$\mathcal{H}^+(\mu) = \mathcal{H}^+(q_{ac}) \oplus \mathcal{H}^+(q_s),$$

where

$$\mathcal{H}^+(q_{ac}) = \text{int}(\mu, \lambda)^{-\|\cdot\|_\mu}.$$

If $\mu = \mu_{ac} + \mu_s$ is the Lebesgue decomposition of μ with respect to λ , then

$$\mathcal{H}^+(\mu) = \mathcal{H}^+(\mu_{ac}) + \mathcal{H}^+(\mu_s),$$

is a complementary space decomposition in the sense of de Branges and Rovnyak, with $\mathcal{H}^+(\mu_{ac}), \mathcal{H}^+(\mu_s)$ contractively contained in $\mathcal{H}^+(\mu)$. Moreover, $\mathcal{H}^+(\mu_{ac})$ is the largest RKHS, $\mathcal{H}(k)$, contractively contained in $\mathcal{H}^+(q_{ac}) \subseteq \mathcal{H}^+(\mu)$ so that the closed embedding, $e : \mathcal{H}(k) \cap \mathcal{H}^+(\lambda) \subseteq \mathcal{H}(k) \hookrightarrow \mathcal{H}^+(\lambda)$, is such that $\tau := ee^*$ is Toeplitz for the image, V^λ , of Z^λ under Cauchy transform, *i.e.* $V^{\lambda*} \tau V^\lambda = \tau$. In particular, the Simon–Lebesgue decomposition of the quadratic form, q_μ , in $H^2(\lambda)$ coincides with the Lebesgue decomposition of μ with respect to λ if and only if $\text{int}(\mu, \lambda)$ is V^μ -invariant.

In the above, the spaces of q_{ac} and q_s -Cauchy transforms are defined in an analogous way to the space of μ -Cauchy transforms, see Subsection 4.1. By Proposition 4.7, the intersection space, $\text{int}(\mu, \lambda)$, is not always V^μ -invariant. Example 4.9 (continued in Example 4.17) provides a concrete example, where $\mu = m_+$ and $\lambda = m_-$ are the mutually singular restrictions of normalized Lebesgue measure, m , to the upper and lower half-circles, so that the Lebesgue decomposition of m_+ with respect to m_- has $m_{+,ac} = 0$ but $\text{int}(m_+, m_-) \neq \{0\}$, so that $q_{m_+,ac} \neq 0$.

Remark This ‘reproducing kernel approach’ to measure theory on the circle and Lebesgue decomposition of a positive measure with respect to Lebesgue measure was first considered and studied in [14, 15], in a more general and non-commutative context.

1 Background

1.1 Function theory in the disk, measure theory on the circle

Classical analytic function theory in the complex unit disk and measure theory on the complex unit circle are fundamentally intertwined. There are bijective correspondences between (i) contractive analytic functions in the disk, (ii) analytic functions in the disk with positive semi-definite real part, *i.e.* *Herglotz functions* and (iii) positive, finite and regular Borel measures on the complex unit circle. Namely, starting with such a positive measure, μ , its *Herglotz–Riesz transform* is the Herglotz function,

$$H_\mu(z) := \int_{\partial\mathbb{D}} \frac{1+z\bar{\zeta}}{1-z\bar{\zeta}} \mu(d\zeta) \in \mathcal{O}(\mathbb{D}).$$

It is easily verified that $\operatorname{Re} H_\mu(z) \geq 0$, is a positive harmonic function. Applying the inverse *Cayley transform* to any Herglotz function, *i.e.* the Möbius transformation sending the open right half-plane onto the open unit disk, \mathbb{D} , which interchanges the points 1 and 0, yields a contractive analytic function, b_μ , in the disk,

$$b_\mu(z) := \frac{H_\mu(z) - 1}{H_\mu(z) + 1}, \quad |b_\mu(z)| \leq 1, \quad z \in \mathbb{D}.$$

(By the maximum modulus principle, b_μ is strictly contractive in \mathbb{D} unless it is constant.) Each of these transformations is essentially reversible. Namely, given any contractive analytic function, b , the Cayley transform, $H_b := \frac{1+b}{1-b}$, is a Herglotz function and the Herglotz representation theorem states that if H is any Herglotz function in the disk, then there is a unique finite, positive and regular Borel measure, μ on the circle, so that

$$H(z) = i \operatorname{Im} H(0) + \int_{\partial\mathbb{D}} \frac{1 + z\bar{\zeta}}{1 - z\zeta} \mu(d\zeta) = i \operatorname{Im} H(0) + H_\mu(z),$$

see [13, Boundary Values, Chapter 3]. To be precise, two Herglotz functions correspond to the same positive measure, μ , if and only if they differ by an imaginary constant. If H_1, H_2 are two Herglotz functions so that $H_2 = H_1 + it$ for some $t \in \mathbb{R}$, then their corresponding inverse Cayley transforms obey

$$b_2 = \frac{\bar{z(t)}}{z(t)} \cdot \mathfrak{m}_{z(t)} \circ b_1, \quad \mathfrak{m}_{z(t)}(z) = \frac{z - z(t)}{1 - \bar{z(t)}z}, \quad z(t) := \frac{t}{2i + t} \in \mathbb{D},$$

so that b_2 is, up to multiplication by the unimodular constant $\frac{\bar{z(t)}}{z(t)}$, a Möbius transformation, $\mathfrak{m}_{z(t)}$, of b_1 , where $\mathfrak{m}_{z(t)}$ defines an automorphism of the disk interchanging 0 with $z(t)$.

If a contractive analytic function, b , corresponds, essentially uniquely, to a positive measure, μ , in this way, we write $\mu := \mu_b$, and μ_b is called the *Clark* or *Aleksandrov–Clark measure* of b [5]. Many properties of contractive analytic functions in the disk can be described in terms of corresponding properties of their Clark measures and vice versa [1, 2]. For example, by Fatou's theorem, the Radon–Nikodym derivative of any Clark measure, μ_b , with respect to normalized Lebesgue measure, m , on the circle is given by the radial, or more generally non-tangential, limits of the real part of its Herglotz function,

$$\begin{aligned} \frac{\mu_b(d\zeta)}{m(d\zeta)} &= \lim_{r \uparrow 1} \operatorname{Re} H_b(r\zeta); \quad m - a.e., \quad \zeta \in \partial\mathbb{D} \\ &= \lim_{r \uparrow 1} \frac{1 - |b(r\zeta)|^2}{|1 - b(r\zeta)|^2} \geq 0, \end{aligned}$$

[7], [13, Fatou's Theorem, Chapter 3]. As a corollary of this formula, we see that b is *inner*, *i.e.* it has unimodular radial boundary limits m -a.e. on the circle, if and only if its Radon–Nikodym derivative vanishes almost everywhere, *i.e.* if and only if its Clark measure is singular with respect to Lebesgue measure.

As a second example which will be relevant for our investigations here, b is an extreme point of the closed convex set of contractive analytic functions in the disk if and only if its Radon–Nikodym derivative with respect to Lebesgue measure is not log-integrable. That is, b is an extreme point if and only if

$$\log \frac{\mu_b(d\zeta)}{m(d\zeta)} \notin L^1 = L^1(m).$$

This follows from the characterization of extreme points in the set of contractive analytic functions given in [13, Extreme Points, Chapter 9] and Fatou's Radon–Nikodym formula as described above. Here, equipping the set of all bounded analytic functions in the disk with the supremum norm, we obtain the unital Banach algebra, H^∞ , the Hardy algebra, whose closed unit ball, $[H^\infty]_1$, is the compact and convex set of contractive analytic functions in the disk. It further follows from a well-known theorem of Szegő (later strengthened by Kolmogoroff and Krein), that $H^2(\mu) = L^2(\mu)$ if and only if $\mu = \mu_b$ for an extreme point $b \in [H^\infty]_1$ [13, Szegő's Theorem, Chapter 4], [23]. Namely, Szegő's theorem gives a formula for the distance from the constant function 1 to the closure of the analytic polynomials with zero constant term in $L^2(\mu)$:

$$\inf_{\substack{p \in \mathbb{C}[\zeta] \\ p(0)=0}} \|1 - p\|_{L^2(\mu)}^2 = \exp \int_{\partial\mathbb{D}} \log \frac{\mu(d\zeta)}{m(d\zeta)} m(d\zeta).$$

It follows, in particular, that b is an extreme point so that $\frac{d\mu}{dm}$ is not log-integrable if and only if 1 belongs to the closure, $H_0^2(\mu)$, in $L^2(\mu)$ of the analytic polynomials obeying $p(0) = 0$. That is, if and only if $H_0^2(\mu) = H^2(\mu)$. An inductive argument then shows that this is equivalent to $H^2(\mu) = L^2(\mu)$, so that $Z^\mu = M_\zeta^\mu|_{H^2(\mu)} = M_\zeta^\mu$ is unitary. If $\mu = \mu_b$ is the Clark measure of an extreme point, b , we will say that μ is *extreme*, and that μ is *non-extreme* if b is not an extreme point.

The results of this paper reinforce the close relationship between function theory in the disk and measure theory on the circle by establishing the Lebesgue decomposition and Radon–Nikodym theorem for positive measures using functional

analysis and reproducing kernel theory applied to spaces of Cauchy transforms of positive measures. We will see that the reproducing kernel construction of the Lebesgue decomposition of a positive measure μ , with respect to another, λ , bifurcates into the two cases, where: the intersection of the spaces of μ and λ -Cauchy transforms, is (i) invariant, or (ii), not invariant for the image of Z^μ under Cauchy transform. Moreover, whether or not this intersection space is invariant is largely dependent on whether λ , or $\mu + \lambda$ are non-extreme or extreme.

1.2 Reproducing kernel Hilbert spaces

As described in the introduction, a reproducing kernel Hilbert space (RKHS) is any complex, separable Hilbert space of functions, \mathcal{H} , on a set X , with the property that the linear functional of point evaluation at any $x \in X$ is bounded on \mathcal{H} . Further recall, as described above, that for any $x \in X$, there is then a unique *kernel vector* or *point evaluation vector*, $k_x \in \mathcal{H}$ so that $\langle k_x, h \rangle_{\mathcal{H}} = h(x)$ for any $h \in \mathcal{H}$ and we write $\mathcal{H} = \mathcal{H}(k)$, where $k : X \times X \rightarrow \mathbb{C}$ is a *positive kernel function* on X in the sense of Equation (0.1). Much of elementary reproducing kernel Hilbert space theory was developed by N. Aronszajn in his seminal paper, [3]. In particular, there is a bijective correspondence between RKHS on a set X and positive kernel functions on X given by the Aronszajn–Moore theorem, [3, Part I], [20, Proposition 2.13, Theorem 2.14] and this motivates the notation $\mathcal{H} = \mathcal{H}(k)$.

Theorem (Aronszajn–Moore) *If $\mathcal{H} = \mathcal{H}(k)$ is a RKHS of functions on a set, X , then k is a positive kernel function on X . Conversely, if k is a positive kernel function on X , then there is a (necessarily unique) RKHS of functions on X with reproducing kernel, k .*

Any RKHS, $\mathcal{H}(k)$, of functions on a set X , is naturally equipped with a *multiplier algebra*, $\text{Mult}(k)$, the unital algebra of all functions on X which ‘multiply’ $\mathcal{H}(k)$ into itself. That is, $g \in \text{Mult}(k)$ if and only if $g \cdot h \in \mathcal{H}(k)$ for any $h \in \mathcal{H}(k)$. Any $h \in \text{Mult}(k)$ can be identified with the linear multiplication operator $M_h : \mathcal{H}(k) \rightarrow \mathcal{H}(k)$. More generally, one can consider the set of multipliers, $\text{Mult}(k, K)$, between two RKHS on the same set. If $h \in \text{Mult}(k, K)$, then M_h is always bounded, by the closed graph theorem. Adjoints of multiplication operators have a natural action on kernel vectors: If $h \in \text{Mult}(k, K)$, then

$$M_h^* K_z = k_z \overline{h(z)}.$$

All RKHS in this paper will be RKHS, $\mathcal{H}(k)$, of analytic functions in the complex unit disk, $\mathbb{D} = (\mathbb{C})_1$, with the additional property that evaluation of the Taylor coefficients of any $h \in \mathcal{H}(k)$ (at 0) defines a bounded linear functional on $\mathcal{H}(k)$. Again, by the Riesz representation lemma, for any $j \in \mathbb{N} \cup \{0\}$, there is then a unique *Taylor coefficient kernel vector*, $k_j \in \mathcal{H}(k)$, so that if $h \in \mathcal{H}(k)$ has Taylor series at 0,

$$h(z) = \sum_{j=0}^{\infty} \hat{h}_j z^j,$$

then $\langle k_j, h \rangle_{\mathcal{H}(k)} = \hat{h}_j$. It follows that

$$\hat{k}(i, j) := \langle k_i, k_j \rangle_{\mathcal{H}(k)},$$

defines a positive kernel function, the *coefficient reproducing kernel* of $\mathcal{H}(k)$, on the set $\mathbb{N} \cup \{0\}$. It is easily checked that for any such Taylor coefficient reproducing kernel Hilbert spaces, $\mathcal{H}(k)$ and $\mathcal{H}(K)$, of analytic functions in \mathbb{D} ,

$$k \leq K \quad \Leftrightarrow \quad \hat{k} \leq \hat{K}.$$

The reproducing and coefficient reproducing kernels of a Taylor coefficient RKHS in \mathbb{D} are related by the formulas:

$$k(z, w) = \sum_{j, \ell=0}^{\infty} \hat{k}(j, \ell) z^j \overline{w}^\ell, \quad \text{and} \quad k_z = \sum_{j=0}^{\infty} \overline{z}^j k_j.$$

Adjoints of multipliers also have a natural convolution action on coefficient kernels, if $h \in \text{Mult}(K, k)$, then,

$$M_h^* K_\ell = \sum_{i+j=\ell} k_i \overline{\hat{h}_j}. \tag{1.1}$$

We will say that a RKHS, $\mathcal{H}(k)$, of analytic functions in $X = \mathbb{D}$ is a *coefficient RKHS in \mathbb{D}* , if Taylor coefficient evaluations define bounded linear functionals on $\mathcal{H}(k)$. In this case the positive coefficient kernel function \hat{k} on $\mathbb{N} \cup \{0\}$ is an example of a discrete or formal reproducing kernel in the sense of [4].

In this paper it will be useful to consider densely-defined multipliers between RKHS $\mathcal{H}(K)$, $\mathcal{H}(k)$ on X , which are not necessarily bounded.

Proposition 1.1 (Multipliers are closeable) Let k, K be positive kernel functions on X , and let h be a function on X so that the linear operator $M_h : \text{Dom } M_h \subseteq \mathcal{H}(k) \rightarrow \mathcal{H}(K)$ has dense domain, $\text{Dom } M_h$. Then M_h is closeable, and closed on its maximal domain, $\text{Dom}_{\max} M_h := \{g \in \mathcal{H}(k) \mid h \cdot g \in \mathcal{H}(K)\}$, $M_h^* K_x = k_x \overline{h(x)}$, and $\bigvee_{x \in X} K_x$ is a core for M_h^* , if M_h is defined on its maximal domain.

Recall that a linear operator with dense domain in a Hilbert space, \mathcal{H} , is said to be *closed* if its graph is a closed subspace of $\mathcal{H} \oplus \mathcal{H}$. Further recall that a dense set, $\mathcal{D} \subseteq \text{Dom } A$, contained in the domain of closed operator, A , is called a *core* for A if A is equal to the *closure* (minimal closed extension) of its restriction to \mathcal{D} . In general, given any two linear transformations A, B , we say that B is an *extension* of A or that A is a restriction of B , written $A \subseteq B$, if $\text{Dom } A \subseteq \text{Dom } B$ and $B|_{\text{Dom } A} = A$. Equivalently, the set of all pairs (x, Ax) , for $x \in \mathcal{D}$, is dense in the graph of A . Finally, A is *closeable* if it has a closed extension.

Proof Define $\text{Dom}_{\max} M_h$ to be the linear space of all $g \in \mathcal{H}(k)$ so that $h \cdot g \in \mathcal{H}(K)$. This is the largest domain on which M_h makes sense. If $g_n \in \text{Dom}_{\max} M_h$ is such that $g_n \rightarrow g$ and $M_h g_n \rightarrow f$, then since $\mathcal{H}(k), \mathcal{H}(K)$ are RKHS, it necessarily follows that

$$g_n(x) \rightarrow g(x), \quad \text{and} \quad h(x)g_n(x) \rightarrow h(x)g(x) = f(x), \quad x \in X.$$

This proves that $f = h \cdot g$, so that $g \in \text{Dom}_{\max} M_h$ and M_h is closed on $\text{Dom}_{\max} M_h$. If M_h is densely-defined on some other domain, $\text{Dom } M_h$, then $\text{Dom } M_h \subseteq \text{Dom}_{\max} M_h$ by maximality, so that M_h has a closed extension, and is hence closeable.

The fact that $\bigvee K_x$ is a core for M_h^* follows from the assumption that M_h is defined (and closed) on its maximal domain. By maximality, M_h , with domain $\text{Dom}_{\max} M_h$, has no non-trivial closed extensions which act as multiplication by h . Let T_* be the closure of the restriction of M_h^* to $\bigvee K_x$. Then $T_* \subseteq M_h^*$ is densely-defined and closed so that $M_h \subseteq T := T_*^*$, where T_*^* , the adjoint of T_* is necessarily closed so that $T^* = T_*$. However,

$$T^* K_x = M_h^* K_x = K_x \overline{h(x)},$$

so that T necessarily acts as multiplication by h on its domain. By maximality, $\text{Dom } T = \text{Dom}_{\max} M_h$ and $M_h = T$. ■

Remark 1.2 If $\mathcal{H}(k)$ and $\mathcal{H}(K)$ are Taylor coefficient RKHS in \mathbb{D} , then one can further show that the adjoint of any closed multiplication operator, $M_h : \mathcal{H}(k) \rightarrow \mathcal{H}(K)$ acts as a convolution operator on coefficient kernels, as in Equation (1.1), and the linear span of all Taylor coefficient kernels is also a core for M_h^* .

One can define a natural partial order on positive kernel functions on a fixed set, X . Namely, if k and K are two positive kernel functions on the same set, X , we write $k \leq K$, if $K - k$ is a positive kernel function on X . Notice that the identically zero kernel function is a positive kernel on X , so that $k \leq K$ can be equivalently written as $K - k \geq 0$. The following theorem of Aronszajn describes when one RKHS of functions on X is boundedly contained in another in terms of this partial order, [3, Section 7] [20, Theorem 5.1].

Theorem (Aronszajn's inclusion theorem) Let k, K be positive kernel functions on a set, X . Then $\mathcal{H}(k) \subseteq \mathcal{H}(K)$ and the norm of the embedding $e : \mathcal{H}(k) \hookrightarrow \mathcal{H}(K)$ is at most $t^2 > 0$ if and only if $t^2 K \geq k$.

If k and K are both positive kernel functions on a set, X , it is immediate that $k + K$ is also a positive kernel function on X . The following 'sums of kernels' theorem of Aronszajn describes the norm of $\mathcal{H}(k + K)$ and the decomposition of this space in terms of $\mathcal{H}(k)$ and $\mathcal{H}(K)$ [3], [20, Theorem 5.4, Corollary 5.5]. Notice, in particular that $k, K \leq k + K$ as kernel functions so that $\mathcal{H}(k)$ and $\mathcal{H}(K)$ are contractively contained in $\mathcal{H}(k + K)$, by the inclusion theorem.

Theorem (Aronszajn's sums of kernels theorem) Let k, K be positive kernel functions on a set, X . Then, $\mathcal{H}(k + K) = \mathcal{H}(k) + \mathcal{H}(K)$ and

$$\|h\|_{\mathcal{H}(k+K)}^2 = \min \{ \|f\|_{\mathcal{H}(k)}^2 + \|g\|_{\mathcal{H}(K)}^2 \mid f + g = h \}.$$

In particular, $\mathcal{H}(k + K) = \mathcal{H}(k) \oplus \mathcal{H}(K)$ if and only if $\mathcal{H}(k) \cap \mathcal{H}(K) = \{0\}$.

Observe that the sums of kernels theorem asserts that the algebraic sum $\mathcal{H}(k + K) = \mathcal{H}(k) + \mathcal{H}(K)$ is a direct sum if and only if it is an orthogonal direct sum. More can be said about this decomposition and the structure of $\mathcal{H}(k + K)$ using the theory of operator-range spaces of contractions and their complementary spaces in the sense of de Branges and Rovnyak [6], [8, Chapter 16]. Let $A \in \mathcal{L}(\mathcal{H}, \mathcal{J})$ be a bounded linear operator. The *operator-range space* of A , $\mathcal{R}(A)$, is the Hilbert space obtained by equipping the range of A with the inner product that makes A a co-isometry onto its range. That is, $\mathcal{R}(A) = \text{Ran } A \subseteq \mathcal{J}$, with inner product,

$$\langle Ax, Ay \rangle_A := \langle x, P_{\text{Ker } A}^\perp y \rangle_{\mathcal{H}}.$$

One can generally show that $\mathcal{R}(A) = \mathcal{R}(\sqrt{AA^*})$, [8, Corollary 16.8]. If A is a contraction, $\|A\| \leq 1$, then $\mathcal{R}(A) \subseteq \mathcal{J}$ is contractively contained in \mathcal{J} in the sense that the embedding, $e : \mathcal{R}(A) \hookrightarrow \mathcal{J}$ is a linear contraction. In this case, one can define the *complementary space of A* , $\mathcal{R}^c(A) := \mathcal{R}(\sqrt{I - AA^*})$. The notion of complementary space was originally introduced in a more geometric way by de Branges and Rovnyak [6]. Namely, if \mathcal{H} is any Hilbert space and $\mathcal{R} \subseteq \mathcal{H}$ is a Hilbert space which is contractively contained in \mathcal{H} , then $\mathcal{R} = \mathcal{R}(j)$, where $j : \mathcal{R} \hookrightarrow \mathcal{H}$ is the contractive embedding. L. de Branges and J. Rovnyak defined the complementary space, \mathcal{R}^c of \mathcal{R} as the set of all $y \in \mathcal{H}$ so that

$$\sup_{x \in \mathcal{R}} (\|y + x\|_{\mathcal{H}}^2 - \|x\|_{\mathcal{R}}^2) < +\infty.$$

One can prove that $\mathcal{R}^c = \mathcal{R}^c(j)$ and that the above formula is equal to the norm of y in $\mathcal{R}^c(j)$, so that these two definitions coincide [8, Chapter 16]. The following theorem summarizes several results in the theory of operator–range spaces, see [8, Chapter 16].

Theorem 1.3 (Operator–range spaces of contractions) Let $A \in \mathcal{L}(\mathcal{H}, \mathcal{J})$ be a contraction. If $e : \mathcal{R}(A) \hookrightarrow \mathcal{J}$ and $j : \mathcal{R}^c(A) \hookrightarrow \mathcal{J}$ are the contractive embeddings, then

$$\mathcal{J} = \mathcal{R}(A) + \mathcal{R}^c(A).$$

For any $x = y + z \in \mathcal{J}$ so that $y \in \mathcal{R}(A)$ and $z \in \mathcal{R}^c(A)$, the Pythagorean equality,

$$\|x\|_{\mathcal{J}}^2 = \|y\|_{\mathcal{R}(A)}^2 + \|z\|_{\mathcal{R}^c(A)}^2, \tag{1.2}$$

holds if and only if $y = ee^*x$ and $z = jj^*x$, so that, in particular, $I_{\mathcal{J}} = ee^* + jj^*$. As a vector space, the overlapping space is

$$\mathcal{R}(A) \cap \mathcal{R}^c(A) = A\mathcal{R}^c(A^*),$$

and $A : \mathcal{R}^c(A^*) \rightarrow \mathcal{R}^c(A)$ acts as a linear contraction.

Moreover, the following are equivalent:

- (i) A is a partial isometry,
- (ii) $\mathcal{R}(A)$ and $\mathcal{R}^c(A)$ are isometrically contained in \mathcal{J} as orthogonal complements, $\mathcal{J} = \mathcal{R}(A) \oplus \mathcal{R}^c(A)$,
- (iii) $\mathcal{R}(A) \cap \mathcal{R}^c(A) = \{0\}$.

Observe that, as in Aronszajn’s sums of kernels theorem, the algebraic sum $\mathcal{J} = \mathcal{R}(A) + \mathcal{R}^c(A)$ is a direct sum if and only if it is an orthogonal direct sum.

Theorem 1.4 Let $\mathcal{H}(K)$ be a RKHS on a set, X . If $\mathcal{H}(k)$ is another RKHS on X which embeds, contractively, in $\mathcal{H}(K)$, and $e : \mathcal{H}(k) \hookrightarrow \mathcal{H}(K)$ is the contractive embedding, then $\mathcal{H}(k) = \mathcal{R}(e)$ and the complementary space, $\mathcal{R}^c(e)$, is the RKHS on X with reproducing kernel $K - k$.

An embedding of RKHS, $e : \mathcal{H}(k) \hookrightarrow \mathcal{H}(K)$, is necessarily injective.

Proof Let $e : \mathcal{H}(k) \hookrightarrow \mathcal{H}(K)$ be the contractive embedding and consider the operator–range space of e . Given any $g, h \in \mathcal{H}(k)$, we have that

$$\langle eg, eh \rangle_e = \langle g, h \rangle_{\mathcal{H}(k)},$$

since e is injective. Hence, for any $x \in X$,

$$\langle ek_x, eh \rangle_e = \langle k_x, h \rangle_{\mathcal{H}(k)} = h(x) = (eh)(x), \tag{1.3}$$

and it follows that $\mathcal{R}(e) = \mathcal{H}(k)$. Indeed, equation (1.3) shows that $\mathcal{R}(e)$ is a reproducing kernel Hilbert space on X with point evaluation vectors $\tilde{k}_x := ek_x$ and that for any $x, y \in X$,

$$\tilde{k}(x, y) = \left\langle \tilde{k}_x, \tilde{k}_y \right\rangle_e = \langle k_x, k_y \rangle = k(x, y),$$

so that $\mathcal{R}(e) = \mathcal{H}(\tilde{k}) = \mathcal{H}(k)$. Now consider the complementary space, $\mathcal{R}^c(e)$, of $\mathcal{H}(k) = \mathcal{R}(e)$. Since this complementary space is contractively contained in $\mathcal{H}(K)$, for any $\sqrt{I - ee^*}g \in \mathcal{R}^c(e)$,

$$\begin{aligned} \left\langle (I - ee^*)K_x, \sqrt{I - ee^*}g \right\rangle_{\mathcal{R}^c} &= \left\langle \sqrt{I - ee^*}K_x, g \right\rangle_{\mathcal{H}(K)} \\ &= (\sqrt{I - ee^*}g)(x), \end{aligned}$$

proving that $\mathcal{R}^c(e)$ is also a RKHS on X with point evaluation vectors $k'_x := (I - ee^*)K_x$. Hence,

$$\begin{aligned} k'(x, y) &= \langle k'_x, k'_y \rangle_{\mathcal{H}} = \langle (I - ee^*)K_x, (I - ee^*)K_y \rangle_{\mathcal{H}} \\ &= \langle K_x, (I - ee^*)K_y \rangle_{\mathcal{H}(K)} \\ &= K(x, y) - k(x, y). \end{aligned}$$

If $j : \mathcal{R}^c(e) \hookrightarrow \mathcal{H}^+(\mu)$ is the contractive embedding, then observe that $jj^* + ee^* = I_{\mathcal{H}(K)}$, so that

$$\mathcal{R}(j) = \mathcal{R}(\sqrt{jj^*}) = \mathcal{R}^c(e). \quad \blacksquare$$

The previous theorem and Theorem 1.3 provide additional information on the structure and decomposition of $\mathcal{H}(k+K)$ in Aronszajn's sums of kernels theorem.

Corollary 1.5 *Let k, K be positive kernel functions on a set, X and let $e : \mathcal{H}(k) \hookrightarrow \mathcal{H}(k+K)$ and $j : \mathcal{H}(K) \hookrightarrow \mathcal{H}(k+K)$ be the contractive embeddings. Then we can identify $\mathcal{H}(k)$ and $\mathcal{H}(K)$ with the operator range spaces $\mathcal{R}(e)$ and $\mathcal{R}(j)$, respectively. Moreover, $I_{\mathcal{H}(k+K)} = ee^* + jj^*$ so that $\mathcal{H}(K) = \mathcal{R}^c(e)$ is the complementary space of $\mathcal{R}(e) = \mathcal{H}(k)$, and given any $h \in \mathcal{H}(k+K)$,*

$$\|h\|_{k+K}^2 = \|e^*h\|_k^2 + \|j^*h\|_K^2.$$

The intersection space, $\mathcal{H}(k) \cap \mathcal{H}(K)$ is equal to $e\mathcal{R}^c(e^)$ and $j\mathcal{R}^c(j^*)$, and $e : \mathcal{R}^c(e^*) \rightarrow \mathcal{R}^c(e) = \mathcal{H}(K)$, $j : \mathcal{R}^c(j^*) \rightarrow \mathcal{R}^c(j)$ are contractions.*

Finally, as described in [18] and [17, Section 5], we can define a pair of natural 'lattice operations', \vee and \wedge on the set of all positive kernel functions on a fixed set, X . Given two positive kernel functions, k and K , on X , let $k \vee K := k + K$, a positive kernel function on X . We can also construct a second RKHS on X by defining

$$\text{int}(k, K) := \mathcal{H}(k) \cap \mathcal{H}(K),$$

equipped with the inner product

$$\langle g, h \rangle_{k \wedge K} := \langle g, h \rangle_k + \langle g, h \rangle_K.$$

It is not difficult to verify that $\text{int}(k, K)$, equipped with this inner product is complete, and that point evaluation at any $x \in X$ defines a bounded linear functional on $\text{int}(k, K)$, so that this is a RKHS, $\mathcal{H}(k \wedge K)$, of functions on X . The following theorem describes a useful relationship between $\mathcal{H}(k+K)$ and $\mathcal{H}(k \wedge K)$ [17, Theorem 5.2], [18].

Theorem (Sums and intersections of RKHS) *Let $k, K \geq 0$ be positive kernel functions on a set, X . Define two linear maps, U_\vee and U_\wedge of $\mathcal{H}(k+K)$ and $\mathcal{H}(k \wedge K)$, respectively, into $\mathcal{H}(k) \oplus \mathcal{H}(K)$ by*

$$U_\vee(k+K)_x := k_x \oplus K_x, \quad x \in X, \quad \text{and} \quad U_\wedge f := f \oplus -f, \quad f \in \mathcal{H}(k \wedge K).$$

Then, U_\vee, U_\wedge both define isometries into $\mathcal{H}(k) \oplus \mathcal{H}(K)$ with $\text{Ran } U_\vee = \text{Ran } U_\wedge^\perp$, so that

$$\mathcal{H}(k) \oplus \mathcal{H}(K) = U_\vee \mathcal{H}(k+K) \oplus U_\wedge \mathcal{H}(k \wedge K).$$

The point evaluation vectors for $\mathcal{H}(k \wedge K) = \mathcal{H}(k) \cap \mathcal{H}(K)$ are given by the formulas

$$(k \wedge K)_x = \frac{1}{2} U_\wedge^*(K_x \oplus -k_x) = U_\wedge^*(K_x \oplus 0) = U_\wedge^*(0 \oplus -k_x).$$

1.3 Positive quadratic forms

A quadratic or sesquilinear form, $q : \text{Dom } q \times \text{Dom } q \rightarrow \mathbb{C}$, with dense *form domain*, $\text{Dom } q$ in a separable Hilbert space, \mathcal{H} , is said to be positive semi-definite if $q(x, x) \geq 0$ for all $x \in \text{Dom } q$. Such a quadratic form is said to be *closed*, if $\text{Dom } q$ is complete with respect to the norm induced by the inner product

$$\langle \cdot, \cdot \rangle_{q+\text{id}} := \langle \cdot, \cdot \rangle_{\mathcal{H}} + q(\cdot, \cdot)$$

and q is *closeable* if it has a closed extension. We will let $\hat{\mathcal{H}}(q)$ denote the Hilbert space completion of $\text{Dom } q$ with respect to this $q + \text{id}$ -inner product. Hence, q is closed if and only if $\hat{\mathcal{H}}(q) = \text{Dom } q$. If $q \geq 0$ is closeable, then its *closure*, \bar{q} , is the minimal closed extension of q . By Kato, closed positive semi-definite forms obey an 'unbounded version' of the Riesz

Lemma [16, Chapter VI, Theorem 2.1 and Theorem 2.23]. Namely, $q \geq 0$ is closed if and only if there is a *unique* self-adjoint, densely-defined and positive semi-definite operator, A , so that $\text{Dom } q = \text{Dom } \sqrt{A}$ and

$$q(x, x) = q_A(x, x) = \left\langle \sqrt{A}x, \sqrt{A}x \right\rangle_{\mathcal{H}}.$$

Any self-adjoint operator is necessarily closed. Following Kato and Simon, we can define a partial order on densely-defined and positive semi-definite forms by $q_1 \leq q_2$ if

- (i) $\text{Dom } q_2 \subseteq \text{Dom } q_1$, and
- (ii) $q_1(x, x) \leq q_2(x, x)$ for all $x \in \text{Dom } q_2$.

In particular, if q_A and q_B are the closed forms of the self-adjoint and positive semi-definite operators A and B , we say that $A \leq B$ in the *form sense* if $q_A \leq q_B$ as forms. This reduces to the usual Löwner partial order on bounded, self-adjoint operators, if A and B are bounded. At first sight, it may seem strange that the ‘larger’ form in the above definition may have a smaller domain. The following result of Kato provides some justification for this [16, Chapter VI, Theorem 2.21], [22, Proposition 1.1].

Lemma 1.6 (Kato) *Let $A, B \geq 0$ be self-adjoint and densely-defined in \mathcal{H} . Then $A \leq B$ in the form sense if and only if*

$$(tI + A)^{-1} \leq (tI + B)^{-1},$$

for any $t > 0$.

Recall that if A is a closed operator with dense domain, $\text{Dom } A \subseteq \mathcal{H}$, that $\mathcal{D} \subseteq \text{Dom } A$ is called a *core* for A , if A is equal to the closure of its restriction to \mathcal{D} . Similarly, if q is a closed, densely-defined and positive semi-definite form, a (necessarily dense) set $\mathcal{D} \subseteq \text{Dom } q$ is called a *form-core* for q , if \mathcal{D} is dense in $\hat{\mathcal{H}}(q)$. It is not difficult to verify that if $q = q_A$ is closed, then \mathcal{D} is a form-core for q if and only if \mathcal{D} is a core for \sqrt{A} .

Toeplitz forms. The classical Hardy space, $H^2 = H^2(\mathbb{D})$, is the Hilbert space of square-summable Taylor series in the complex unit disk, equipped with the ℓ^2 -inner product of these coefficients. By results of Fatou, any element of H^2 has non-tangential boundary limits almost everywhere on the unit circle, $\partial\mathbb{D}$, with respect to normalized Lebesgue measure, m [13]. Identifying any $h \in H^2$ with its boundary limits defines an isometric inclusion of H^2 into $L^2 = L^2(m)$. Classically, Toeplitz operators, T , on H^2 , are defined as the compression of bounded multiplication operators on L^2 to H^2 . Namely, $T = T_g := P_{H^2} M_g|_{H^2}$, and $\|T_g\| = \|M_g\| = \|g\|_\infty$ so that $g \in L^\infty$. A theorem of Brown and Halmos, [10, Theorem 6], characterizes the Toeplitz operators as the set of all bounded operators, T , on H^2 which obey the simple algebraic condition,

$$S^*TS = T,$$

where $S = M_z$, is the *shift* on H^2 , the isometry of multiplication by z . Under the boundary value identification of H^2 with the subspace $H^2(m) \subseteq L^2(m, \partial\mathbb{D})$, the shift is identified with the isometry $S = Z^m = M_z^m|_{H^2(m)}$.

Recall, as described in the Outline, given a positive, finite and regular Borel measure, μ , on $\partial\mathbb{D}$, we can associate to μ the densely-defined and positive semi-definite quadratic form, q_μ , with dense form domain, $\text{Dom } q_\mu := A(\mathbb{D})$ in $H^2 = H^2(m)$, where m is normalized Lebesgue measure. This positive form, q_μ , is an example of a *Toeplitz form*, as studied by Grenander and Szegő in [9]. Namely, $\text{Dom } q_\mu = A(\mathbb{D})$ obeys $S\text{Dom } q_\mu \subseteq \text{Dom } q_\mu$, and

$$q_\mu(Sx, Sy) = q_\mu(x, y); \quad x, y \in A(\mathbb{D}) = \text{Dom } q_\mu.$$

If q_μ is closeable so that $\overline{q_\mu} = q_T$ for a closed, self-adjoint $T \geq 0$, then by Kato’s unbounded Riesz lemma we have that $S^*TS = T$, and our results will show that

$$q_\mu(x, y) = \left\langle M_{\sqrt{f}}x, M_{\sqrt{f}}y \right\rangle_{L^2}; \quad f = \frac{d\mu}{dm} \in L^1, \quad x, y \in A(\mathbb{D}),$$

see Theorem 3.12. Hence, $T = T_f = P_{H^2} M_f|_{H^2}$ is a closed, potentially unbounded Toeplitz operator with symbol f , in this ‘quadratic form sense’. In particular, if $T \geq 0$ is bounded, which happens if and only if $\mu \leq t^2m$ for some $t > 0$, then by the Riesz representation lemma, $S^*TS = T$, so that T is a bounded Toeplitz operator by Brown–Halmos, in which case $T = T_f$ for $f = \frac{d\mu}{dm} \in L^\infty$ and $\|f\|_\infty \leq t$, by Theorem 3.1 and Corollary 3.2.

2 Spaces of Cauchy transforms

Let μ be a positive, finite and regular Borel measure on the complex unit circle. Recall that given any $h \in H^2(\mu) = \mathbb{C}[\zeta]^{-\|\cdot\|_{L^2(\mu)}}$, its μ -Cauchy transform is the function

$$(\mathcal{C}_\mu h)(z) := \int_{\partial\mathbb{D}} h(\zeta) \frac{1}{1 - z\bar{\zeta}} \mu(d\zeta) = \langle k_z, h \rangle_{L^2(\mu)}; \quad k_z(\zeta) := (1 - \bar{z}\zeta)^{-1}. \quad (2.1)$$

We will call the functions $k_z, z \in \mathbb{D}$, Szegő kernel vectors.

Recall that $A(\mathbb{D})$ denotes the disk algebra, the unital Banach algebra of analytic functions in \mathbb{D} which extend continuously to the boundary, $\partial\mathbb{D}$. Since the analytic polynomials are supremum-norm dense in $A(\mathbb{D})$, viewed as a subspace of the continuous functions, $\mathcal{C}(\partial\mathbb{D})$, on the circle and $H^2(\mu) = A(\mathbb{D})^{-\|\cdot\|_{L^2(\mu)}}$, it follows that $H^2(\mu) = \mathbb{C}[\zeta]^{-\|\cdot\|_{L^2(\mu)}} = A(\mathbb{D})^{-\|\cdot\|_{L^2(\mu)}}$. Similarly, $\mathcal{K}_\mathbb{D} := \bigvee_{z \in \mathbb{D}} k_z$ is also supremum-norm dense in $A(\mathbb{D}) \subseteq \mathcal{C}(\partial\mathbb{D})$, so that this subspace is also dense in $H^2(\mu)$.

Lemma 2.1 *The μ -Cauchy transform of any $h \in H^2(\mu)$ is holomorphic in \mathbb{D} .*

Proof Given any $z \in \mathbb{D}$, since $(\mathcal{C}_\mu h)(z) := \langle k_z, h \rangle_{L^2(\mu)}$, consider the Leibniz difference quotient

$$\lim_{\epsilon \rightarrow 0} \frac{(\mathcal{C}_\mu h)(z + \epsilon) - (\mathcal{C}_\mu h)(z)}{\epsilon} = \lim_{\epsilon \rightarrow 0} \epsilon^{-1} \langle k_{z+\epsilon} - k_z, h \rangle_{L^2(\mu)}.$$

(Here, recall that our inner products are conjugate linear in their first argument.) This limit will exist and $\mathcal{C}_\mu h$ will be holomorphic, if and only if the limit of $\bar{\epsilon}^{-1}(k_{z+\epsilon} - k_z)$ exists in $H^2(\mu)$. This limit exists in supremum norm on the circle (and so belongs to $A(\mathbb{D})$), and so it certainly exists in the $L^2(\mu)$ -norm by Cauchy-Schwarz. Indeed,

$$\begin{aligned} \lim_{\epsilon \rightarrow 0} \frac{\overline{k_{z+\epsilon}(\zeta) - k_z(\zeta)}}{\bar{\epsilon}} &= \lim_{\epsilon \rightarrow 0} \frac{1 - \bar{\zeta}z - (1 - \bar{\zeta}z - \bar{\zeta}\epsilon)}{\epsilon(1 - \bar{\zeta}z)(1 - \bar{\zeta}(z + \epsilon))} \\ &= \frac{\bar{\zeta}}{(1 - \bar{\zeta}z)^2}, \end{aligned}$$

and this limit is continuous on $\partial\mathbb{D}$ for any fixed $z \in \mathbb{D}$. Hence $\mathcal{C}_\mu h \in \mathcal{O}(\mathbb{D})$ for any $h \in H^2(\mu)$. ■

Let $\mathcal{H}^+(\mu) := \mathcal{C}_\mu H^2(\mu) \subseteq \mathcal{O}(\mathbb{D})$ be the complex vector space of μ -Cauchy transforms equipped with the inner product,

$$\langle \mathcal{C}_\mu g, \mathcal{C}_\mu h \rangle_\mu := \langle g, h \rangle_{L^2(\mu)}.$$

Lemma 2.2 *The space of μ -Cauchy transforms, $\mathcal{H}^+(\mu)$, is a RKHS of analytic functions in \mathbb{D} with reproducing kernel*

$$\begin{aligned} k^\mu(z, w) &:= \int_{\partial\mathbb{D}} \frac{1}{1 - z\bar{\zeta}} \frac{1}{1 - \bar{w}\zeta} \mu(d\zeta) = \langle k_z, k_w \rangle_{L^2(\mu)} \\ &= \frac{1}{2} \frac{H_\mu(z) + \overline{H_\mu(w)}}{1 - z\bar{w}}, \end{aligned}$$

where $H_\mu(z) = \int_{\partial\mathbb{D}} \frac{1+z\bar{\zeta}}{1-z\bar{\zeta}} \mu(d\zeta)$ is the Herglotz-Riesz transform of μ .

By construction, \mathcal{C}_μ is an isometry of $H^2(\mu)$ onto $\mathcal{H}^+(\mu)$.

Proof To show that this inner product is well-defined, we need to check that $\mathcal{C}_\mu h \equiv 0$ in the disk implies that $h = 0$ in $H^2(\mu)$. Indeed,

$$(\mathcal{C}_\mu h)(z) = \langle k_z, h \rangle_{H^2(\mu)},$$

and since $\bigvee k_z$ is dense in $A(\mathbb{D})$, the linear span of the Szegő kernels is also dense in $H^2(\mu)$ as described above. Hence this vanishes for all $z \in \mathbb{D}$ if and only if $h = 0$.

By definition, for any $z \in \mathbb{D}$,

$$(\mathcal{C}_\mu h)(z) = \langle k_z, h \rangle_{H^2(\mu)} = \langle \mathcal{C}_\mu k_z, \mathcal{C}_\mu h \rangle_\mu,$$

so that $\mathcal{H}^+(\mu)$ is a RKHS in \mathbb{D} with kernel vectors $k_z^\mu := \mathcal{C}_\mu k_z$ and reproducing kernel

$$k^\mu(z, w) := \langle \mathcal{C}_\mu k_z, \mathcal{C}_\mu k_w \rangle_{L^2(\mu)} = \int_{\partial\mathbb{D}} \frac{1}{1-z\bar{\zeta}} \frac{1}{1-\bar{w}\zeta} \mu(d\zeta).$$

Finally,

$$\begin{aligned} H_\mu(z) + \overline{H_\mu(w)} &= \int_{\partial\mathbb{D}} \frac{1+z\bar{\zeta}}{1-z\bar{\zeta}} \mu(d\zeta) + \int_{\partial\mathbb{D}} \frac{1+\bar{w}\zeta}{1-\bar{w}\zeta} \mu(d\zeta) \\ &= 2(1-z\bar{w}) \int_{\partial\mathbb{D}} \frac{1}{1-z\bar{\zeta}} \frac{1}{1-\bar{w}\zeta} \mu(d\zeta), \end{aligned}$$

establishing the second formula. ■

Example 2.3 (Hardy space.) If $\mu = m$ is normalized Lebesgue measure, then,

$$\begin{aligned} H_m(z) &:= \int_{\partial\mathbb{D}} \frac{1+z\bar{\zeta}}{1-z\bar{\zeta}} m(d\zeta) \\ &= \int_{\partial\mathbb{D}} \frac{2}{1-z\bar{\zeta}} m(d\zeta) - m(\partial\mathbb{D}) \\ &= \sum_{j=0}^{\infty} z^j \int_{\partial\mathbb{D}} \bar{\zeta}^j m(d\zeta) - \int_{\partial\mathbb{D}} m(d\zeta) \\ &= \sum_{j=0}^{\infty} z^j \frac{1}{2\pi} \int_0^{2\pi} e^{-ij\theta} d\theta - \frac{1}{2\pi} \int_0^{2\pi} d\theta = 2\delta_{j,0} - 1 = 1. \end{aligned}$$

It follows that $b_m := \frac{H_m^{-1}}{H_m+1} \equiv 0$, so that $m = \mu_0$ is the Clark measure of the identically 0 function. Moreover,

$$k^m(z, w) = \frac{1}{2} \frac{H_m(z) + \overline{H_m(w)}}{1-z\bar{w}} = \frac{1}{1-z\bar{w}} = k(z, w),$$

is the *Szegő kernel*. This is the reproducing kernel for the classical Hardy space $H^2 = H^2(\mathbb{D})$, of square-summable Taylor series in the complex unit disk, equipped with the ℓ^2 -inner product of the Taylor coefficients. That is, $\mathcal{H}^+(m) = H^2$.

Since any $h := \mathcal{C}_\mu g \in \mathcal{H}^+(\mu)$ is holomorphic in the open unit disk, its Taylor series at 0 has radius of convergence at least one,

$$h(z) = \sum_{j=0}^{\infty} \hat{h}_j z^j.$$

Moreover, expanding $k_z(\zeta)$ in a convergent geometric sum,

$$\begin{aligned} h(z) &= \sum_{j=0}^{\infty} \hat{h}_j z^j = \langle k_z, g \rangle_{L^2(\mu)} = \sum_{j=0}^{\infty} z^j \langle \zeta^j, g \rangle_{L^2(\mu)} \\ &= \sum_{j=0}^{\infty} z^j \langle \mathcal{C}_\mu \zeta^j, h \rangle_\mu, \end{aligned}$$

and it follows that the Taylor coefficients are given by

$$\hat{h}_j = \langle \mathcal{C}_\mu \zeta^j, h \rangle_\mu; \quad j \in \mathbb{N} \cup \{0\}.$$

That is, for any $j \in \mathbb{N} \cup \{0\}$, the linear functionals $\ell_j(h) = \hat{h}_j$ are bounded on $\mathcal{H}^+(\mu)$ and are implemented by inner products against the Taylor coefficient kernel vectors $k_j^\mu := \mathcal{C}_\mu \zeta^j$. Hence $\mathcal{H}^+(\mu)$ is a Taylor coefficient RKHS in \mathbb{D} with coefficient reproducing kernel, $\hat{k}^\mu(i, j)$, on the set $\mathbb{N} \cup \{0\}$,

$$\hat{k}^\mu(i, j) := \langle k_i^\mu, k_j^\mu \rangle_\mu,$$

and \hat{k}^μ is then a positive kernel function on $\mathbb{N} \cup \{0\}$.

Given a positive measure μ , let $V_\mu := \mathcal{C}_\mu Z^\mu \mathcal{C}_\mu^*$. This is an isometry on $\mathcal{H}^+(\mu)$ that will play a central role in our analysis. This operator has a natural action on kernel vectors:

$$V_\mu k_z^\mu \bar{z} = k_z^\mu - k_0^\mu, \quad \text{and} \quad V_\mu^*(k_z^\mu - k_0^\mu) = k_z^\mu \bar{z}. \quad (2.2)$$

In particular,

$$\bigvee_{z \in \mathbb{D}} (k_z^\mu - k_0^\mu) = \text{Ran } V_\mu,$$

where, here, \bigvee denotes closed linear span. It is easy to check that a function, $h \in \mathcal{H}^+(\mu)$, is orthogonal to $\text{Ran } V_\mu$ if and only if $h = c1$, $c \in \mathbb{C}$, is constant in the disk. Hence the following statements are equivalent:

- (i) μ is extreme,
- (ii) $H^2(\mu) = L^2(\mu)$,
- (iii) $Z^\mu = M_\zeta^\mu|_{H^2(\mu)}$ and hence V_μ is unitary,
- (iv) $\mathcal{H}^+(\mu)$ does not contain the constant functions.

Lemma 2.4 Given any finite, positive and regular Borel measure, μ , on $\partial\mathbb{D}$, the co-isometry V_μ^* acts as a backward shift on $\mathcal{H}^+(\mu)$, i.e. if $h \in \mathcal{H}^+(\mu)$, $h(z) = \sum_{j=0}^\infty \hat{h}_j z^j$, then

$$(V_\mu^* h)(z) = \sum_{j=0}^\infty \hat{h}_{j+1} z^j = \frac{h(z) - h(0)}{z}.$$

Given any $h \in \mathcal{H}^+(\mu)$,

$$(V_\mu h)(z) = zh(z) + (V_\mu h)(0)1.$$

Given any h in the classical Hardy space, $H^2 = \mathcal{H}^+(m)$, one can check that $S := V_m = M_z$ is the isometry of multiplication by the independent variable, z , on H^2 , the *shift*. In this case, adjoint of S is called the backward shift and acts as

$$(S^* h)(z) = \frac{h(z) - h(0)}{z}.$$

It is straightforward to verify that if $h \in H^2$ has Taylor series $h(z) = \sum \hat{h}_j z^j$, then $(S^* h)(z) = \sum_{j=0}^\infty \hat{h}_{j+1} z^j$. This motivates the terminology 'backward shift' in the above lemma statement. This lemma is easily verified and we omit the proof.

3 Absolute continuity in the reproducing kernel sense

Recall that given positive measures μ and λ , we say that μ is *dominated* by λ if there is a $t > 0$ so that $\mu \leq t^2 \lambda$, and we say that μ is *reproducing kernel* or *RK-dominated* by λ , if $\mathcal{H}^+(\mu) \subseteq \mathcal{H}^+(\lambda)$ and there is a $t > 0$ so that the norm of the embedding $e_{\mu,\lambda} : \mathcal{H}^+(\mu) \hookrightarrow \mathcal{H}^+(\lambda)$ is at most t , written $\mu \leq_{RK} t^2 \lambda$. We will begin this section by showing that these two definitions of domination are equivalent.

Theorem 3.1 Given positive, finite and regular Borel measures μ, λ on the unit circle, $\mu \leq t^2 \lambda$ for some $t > 0$ if and only if $\mu \leq_{RK} t^2 \lambda$.

Proof (Necessity.) If $\mu \leq t^2 \lambda$, then $\gamma := t^2 \lambda - \mu$ is a positive measure and

$$\begin{aligned} t^2 k^\lambda(z, w) - k^\mu(z, w) &= t^2 \int_{\partial\mathbb{D}} \frac{1}{1 - z\bar{\zeta}} \frac{1}{1 - \bar{w}\zeta} \lambda(d\zeta) - \int_{\partial\mathbb{D}} \frac{1}{1 - z\bar{\zeta}} \frac{1}{1 - \bar{w}\zeta} \mu(d\zeta) \\ &= \int_{\partial\mathbb{D}} \frac{1}{1 - z\bar{\zeta}} \frac{1}{1 - \bar{w}\zeta} \gamma(d\zeta) = k^\gamma(z, w). \end{aligned}$$

It follows that $t^2 k^\lambda - k^\mu = k^\gamma$ is a positive kernel so that $k^\mu \leq t^2 k^\lambda$.

First proof of sufficiency. Conversely, suppose that $K := t^2 k^\lambda - k^\mu \geq 0$ is a positive kernel. View the analytic polynomials, $\mathbb{C}[\zeta]$, as a dense subspace of the disk algebra $A(\mathbb{D})$, embedded isometrically in the Banach space $\mathcal{C}(\partial\mathbb{D})$. For any finite, positive and regular Borel measure on the complex unit circle, μ , we define the positive linear functional, $\hat{\mu}$ on $\mathcal{C}(\partial\mathbb{D})$ by

$$\hat{\mu}(f) := \int_{\partial\mathbb{D}} f d\mu.$$

(The map $\mu \mapsto \hat{\mu}$ is a bijection, by the Riesz–Markov theorem.) We then define a bounded linear functional, ℓ_K on $\mathcal{C}(\partial\mathbb{D})$ by $\ell_K := t^2\lambda - \hat{\mu}$.

By Weierstraß approximation, $\mathbb{C}[\zeta] + \overline{\mathbb{C}[\zeta]}$ is supremum-norm dense in the continuous functions, $\mathcal{C}(\partial\mathbb{D})$. Since the Fejér kernel is positive semi-definite, the partial Cesàro sums of any positive semi-definite $f \in \mathcal{C}(\partial\mathbb{D})$ will be a positive trigonometric polynomial, i.e. a positive semi-definite element of $\mathbb{C}[\zeta] + \overline{\mathbb{C}[\zeta]}$ and, by Fourier theory, it follows that the positive cone of $\mathbb{C}[\zeta] + \overline{\mathbb{C}[\zeta]}$ is supremum norm-dense in the positive cone of $\mathcal{C}(\partial\mathbb{D})$. Moreover, by the Fejér–Riesz theorem, any positive trigonometric polynomial, $p + \bar{q} \geq 0$, on $\partial\mathbb{D}$ factors as $|g|^2$ for an analytic $g \in \mathbb{C}[\zeta]$ (and necessarily, $p = q$, $\deg(p) = \deg(g)$). Hence, to check that ℓ_K is a positive linear functional on $\mathcal{C}(\partial\mathbb{D})$, it suffices to check that $\ell_K(p + \bar{p}) \geq 0$ for any $p + \bar{p} = |g|^2 \geq 0$, $p, g \in \mathbb{C}[\zeta]$. If $p = \sum_{j=0}^n \hat{p}_j \zeta^j$ and $g = \sum_{j=0}^n \hat{g}_j \zeta^j$, then by construction

$$\langle \mathcal{E}_\mu g, \mathcal{E}_\mu g \rangle_\mu = \langle g, g \rangle_{H^2(\mu)} = \langle p, 1 \rangle_{H^2(\mu)} + \langle 1, p \rangle_{H^2(\mu)} = \langle \mathcal{E}_\mu p, \mathcal{E}_\mu 1 \rangle_\mu + \langle \mathcal{E}_\mu 1, \mathcal{E}_\mu p \rangle_\mu.$$

Since

$$\mathcal{E}_\mu p = \sum_{j=0}^n \hat{p}_j k_j^\mu,$$

where the k_j^μ , $j \in \mathbb{N} \cup \{0\}$ are the Taylor coefficient evaluation vectors, and since similar formulas hold for λ , we obtain that

$$\ell_K(|g|^2) = \int_{\partial\mathbb{D}} |g(\zeta)|^2 (t^2\lambda(d\zeta) - \mu(d\zeta)) = \sum_{i,j=0}^n \bar{\hat{g}}_i \hat{g}_j \langle K_i, K_j \rangle_{\mathcal{H}(K)},$$

where the K_i , $i \in \mathbb{N} \cup \{0\}$ are the Taylor coefficient evaluation vectors in $\mathcal{H}(K)$. Namely, $\mathcal{H}(K)$ is also a Taylor coefficient RKHS in \mathbb{D} so that $\hat{K}(i, j) := \langle K_i, K_j \rangle_{\mathcal{H}(K)}$ defines a positive kernel function on the set $\mathbb{N} \cup \{0\}$. It follows that

$$\int_{\partial\mathbb{D}} |g(\zeta)|^2 (t^2\lambda(d\zeta) - \mu(d\zeta)) \geq 0,$$

for any $g \in \mathbb{C}[\zeta]$, or, equivalently,

$$\ell_K(p + \bar{p}) = \int_{\partial\mathbb{D}} (p + \bar{p})(t^2 d\lambda - d\mu) \geq 0,$$

for any positive semi-definite $p + \bar{p} \in \mathbb{C}[\zeta] + \overline{\mathbb{C}[\zeta]}$. By density of the positive cone of $\mathbb{C}[\zeta] + \overline{\mathbb{C}[\zeta]}$ in the positive cone of the continuous functions, it follows that ℓ_K is a bounded, positive linear functional on $\mathcal{C}(\partial\mathbb{D})$, with norm $\|\ell_K\| = \ell_K(1) = t^2\lambda(\partial\mathbb{D}) - \mu(\partial\mathbb{D}) = K(0, 0) \geq 0$. By the Riesz–Markov theorem, there is then a unique, positive, finite and regular Borel measure, γ , on $\partial\mathbb{D}$, so that

$$\ell_K(f) = \int_{\partial\mathbb{D}} f(\zeta)\gamma(d\zeta),$$

for any $f \in \mathcal{C}(\partial\mathbb{D})$, i.e. $\ell_K = \hat{\gamma}$, and we conclude that $\gamma = t^2\lambda - \mu \geq 0$ so that $t^2\lambda \geq \mu$.

Second proof of sufficiency. If $t^2k^\lambda \geq k^\mu$, then by Aronszajn’s inclusion theorem, $\mathcal{H}^+(\mu) \subseteq \mathcal{H}^+(\lambda)$ and the norm of the embedding $e_{\mu,\lambda} : \mathcal{H}^+(\mu) \hookrightarrow \mathcal{H}^+(\lambda)$ is at most $t > 0$. Observe that $e := e_{\mu,\lambda}$ acts trivially as a multiplier by the constant function 1, so that

$$e^* k_z^\lambda = k_z^\mu, \quad \text{and} \quad e^* k_j^\lambda = k_j^\mu,$$

for any $z \in \mathbb{D}$ and $j \in \mathbb{N} \cup \{0\}$. Hence,

$$\begin{aligned} e^* V^\lambda k_z^\lambda \bar{z} &= e^*(k_z^\lambda - k_0^\lambda) \\ &= k_z^\mu - k_0^\mu = V^\mu k_z^\mu \bar{z} \\ &= V^\mu e^* k_z^\lambda \bar{z}, \end{aligned}$$

so that e^* intertwines V^λ with V^μ , $e^* V^\lambda = V^\mu e^*$. Setting $E := \mathcal{E}_\mu^* e^* \mathcal{E}_\lambda$, we see that for any monomial,

$$\begin{aligned} E \zeta^j &= \mathcal{E}_\mu^* e^* k_j^\lambda \\ &= \mathcal{E}_\mu^* k_j^\mu = \zeta^j \in H^2(\mu), \end{aligned}$$

so that $E : H^2(\lambda) \hookrightarrow H^2(\mu)$ obeys $Ep = p$ for any $p \in \mathbb{C}[\zeta] \subseteq H^2(\lambda)$. In particular, $EZ^\lambda = Z^\mu E$. At this point one could argue using the Riesz–Markov theorem as above, however, here is an alternative argument. Since Z^λ and Z^μ are contractions (they are isometries), we can apply the intertwining version of the commutant lifting theorem [19, Corollary 5.9] to conclude that E can be ‘lifted’ to a bounded operator $\hat{E} : L^2(\lambda) \rightarrow L^2(\mu)$, with norm $\|\hat{E}\| = \|E\|$, so that

$\hat{E}M_\zeta^\lambda = M_\zeta^\mu \hat{E}$, and $P_{H^2(\mu)} \hat{E}|_{H^2(\lambda)} = E$. Setting $\hat{T} := \hat{E}^* \hat{E} \in \mathcal{L}(L^2(\lambda))$, $\hat{T} \geq 0$ is a positive semi-definite M_ζ^λ -Toeplitz operator in the sense that

$$M_\zeta^{\lambda*} \hat{T} M_\zeta^\lambda = \hat{E}^* M_\zeta^{\mu*} M_\zeta^\mu \hat{E} = \hat{T},$$

since $M_\zeta^\lambda, M_\zeta^\mu$ are isometries. Since, M_ζ^λ (and M_ζ^μ) is in fact unitary, it follows that \hat{T} commutes with M_ζ^λ . Since $\mathbb{C}[\zeta] + \overline{\mathbb{C}[\zeta]} \subseteq \text{Ran } M_\zeta^\lambda$ is a dense set in $L^2(\lambda)$, a simple argument shows that \hat{T} acts as multiplication by $f := T1 \in L^2(\lambda)$. However, since $T = M_f$ is a bounded and positive semi-definite operator, it is easy to check that $t^2 \geq \|\hat{T}\| = \|f\|_\infty$, $\|\hat{T}\| = \|\hat{E}\|^2 = \|E\|^2 = \|T\|$, and $f \geq 0$ λ -a.e. Finally, one can also check that $T = P_{H^2(\lambda)} \hat{T}|_{H^2(\lambda)}$ is Z^λ -Toeplitz, i.e. $Z^{\lambda*} T Z^\lambda = T$. In conclusion, for any $p, q \in \mathbb{C}[\zeta]$,

$$\begin{aligned} \int_{\partial\mathbb{D}} \overline{p(\zeta)} q(\zeta) \mu(d\zeta) &= \langle p, q \rangle_{H^2(\mu)} = \langle Ep, Eq \rangle_{H^2(\mu)} \\ &= \langle \sqrt{T}p, \sqrt{T}q \rangle_{H^2(\lambda)} = \langle p, \hat{T}q \rangle_{L^2(\lambda)} \\ &= \langle p, M_f q \rangle_{L^2(\lambda)} = \int_{\partial\mathbb{D}} \overline{p(\zeta)} q(\zeta) f(\zeta) \lambda(d\zeta). \end{aligned}$$

This formula extends to elements of the form $g = \overline{p} + q \in \mathbb{C}[\zeta] + \overline{\mathbb{C}[\zeta]}$ since

$$\langle \overline{p_1} + q_1, \overline{p_2} + q_2 \rangle_{L^2(\mu)} = \langle q_1, q_2 \rangle_{H^2(\mu)} + \langle q_1 p_2, 1 \rangle_{H^2(\mu)} + \langle p_2, p_1 \rangle_{H^2(\mu)} + \langle 1, p_1 q_2 \rangle_{H^2(\mu)}.$$

Again, by Weierstraß approximation, since $\mathbb{C}[\zeta] + \overline{\mathbb{C}[\zeta]}$ is supremum-norm dense in $\mathcal{C}(\partial\mathbb{D})$, which is in turn dense in $L^2(\lambda)$ and $L^2(\mu)$, it follows that for any $g, h \in L^2(\lambda)$,

$$\int_{\partial\mathbb{D}} \overline{g(\zeta)} h(\zeta) \mu(d\zeta) = \int_{\partial\mathbb{D}} \overline{g(\zeta)} h(\zeta) f(\zeta) \lambda(d\zeta),$$

where $f \geq 0$, λ -a.e. and $\|f\|_\infty \leq t^2$. We conclude that $\mu \leq t^2 \lambda$ and that

$$f = \frac{\mu(d\zeta)}{\lambda(d\zeta)},$$

is the (bounded) Radon–Nikodym derivative of μ with respect to λ . ■

Definition 1. Let $T \in \mathcal{L}(\mathcal{H})$ be a bounded operator and let V be an isometry on \mathcal{H} . We say that T is V -Toeplitz if

$$V^* T V = T.$$

If $q \geq 0$ is a positive semi-definite quadratic form with dense form domain, $\text{Dom } q \subseteq \mathcal{H}$, we say that q is V -Toeplitz if $\text{Dom } q$ is V -invariant and

$$q(Vg, Vh) = q(g, h); \quad g, h \in \text{Dom } q.$$

In particular, if $T \geq 0$ is a positive semi-definite, self-adjoint and densely-defined operator in \mathcal{H} , we say that T is V -Toeplitz if the closed, positive semi-definite form it generates,

$$q_T(x, y) := \langle \sqrt{T}x, \sqrt{T}y \rangle_{\mathcal{H}}; \quad x, y \in \text{Dom } q_T = \text{Dom } \sqrt{T},$$

is V -Toeplitz. If $T \geq 0$ is bounded, this latter definition reduces to the definition of a bounded, positive semi-definite V -Toeplitz operator.

Corollary 3.2 Let μ, λ be positive, finite and regular Borel measures on $\partial\mathbb{D}$ so that $\mu \leq t^2 \lambda$. In this case,

$$E_{\mu, \lambda} := \mathcal{C}_\mu^* e_{\mu, \lambda}^* \mathcal{C}_\lambda : H^2(\lambda) \hookrightarrow H^2(\mu),$$

is a co-embedding in the sense that $E_{\mu, \lambda} p = p$ for any $p \in \mathbb{C}[\zeta]$. Moreover,

$$e_{\mu, \lambda}^* k_z^\lambda = k_z^\mu, \quad e_{\mu, \lambda}^* k_j^\lambda = k_j^\mu, \quad \forall z \in \mathbb{D}, j \in \mathbb{N} \cup \{0\},$$

$e_{\mu, \lambda}^* V \lambda = V^\mu e_{\mu, \lambda}^*$ and equivalently $E_{\mu, \lambda} Z^\lambda = Z^\mu E_{\mu, \lambda}$. If $T_\mu := E_{\mu, \lambda}^* E_{\mu, \lambda}$ then T_μ is Z^λ -Toeplitz and T is the compression of a bounded multiplication operator, $T = P_{H^2(\lambda)} M_f|_{H^2(\lambda)}$, where $f \geq 0$ λ -a.e., $\|f\|_\infty \leq t^2$, and $f = \frac{\mu(d\zeta)}{\lambda(d\zeta)}$ is the Radon–Nikodym derivative of μ with respect to λ .

Remark 3.3 While the co-embedding, $E_{\mu, \lambda} : H^2(\lambda) \hookrightarrow H^2(\mu)$ always has dense range, it may have non-trivial kernel. For example, if λ is the sum of two Dirac point masses at distinct points $\zeta, \xi \in \partial\mathbb{D}$, μ is the point mass at ζ , then $\mu \leq \lambda$

and if p is any polynomial vanishing at ζ , then $E_{\mu,\lambda}p = 0 \in H^2(\mu)$. To be precise, $H^2(\mu)$ is the closure of the disk algebra, $A(\mathbb{D})$, or the polynomials, $\mathbb{C}[\zeta]$, in the $L^2(\mu)$ -norm, so that if $p \in \mathbb{C}[\zeta]$ or $a \in A(\mathbb{D})$ vanishes on the support of μ , then $p = 0 = a$ in $H^2(\mu)$.

More generally, absolute continuity of positive measures can also be described in terms of their spaces of Cauchy transforms. It is a straightforward exercise, using the Radon–Nikodym formula, to show that μ is absolutely continuous with respect to λ , if and only if one can construct a monotonically increasing sequence of positive measures, $\mu_n \geq 0$, so that $\mu_n \leq \mu$ for all n , the $\mu_n \uparrow \mu$ increase monotonically to μ , and there is a sequence of positive constants, $t_n > 0$ so that $\mu_n \leq t_n^2 \lambda$. Indeed, this can be readily established by taking the ‘join’ or point-wise maxima of $\frac{d\mu}{d\lambda}$ and the constant functions $t_n^2 \cdot 1$. Since $\mu_n \leq \mu$ for all n , Aronszajn’s inclusion theorem implies that $\mathcal{H}^+(\mu_n) \subseteq \mathcal{H}^+(\mu)$ and that the embeddings $e_n : \mathcal{H}^+(\mu_n) \hookrightarrow \mathcal{H}^+(\mu)$ are all contractive. Moreover, and again by Aronszajn’s inclusion theorem, each $\mathcal{H}^+(\mu_n) \subseteq \mathcal{H}^+(\lambda)$ is boundedly contained in $\mathcal{H}^+(\lambda)$ so that

$$\mathcal{H}^+(\mu_n) \subseteq \mathcal{H}^+(\mu) \cap \mathcal{H}^+(\lambda) =: \text{int}(\mu, \lambda). \quad (3.1)$$

Proposition 3.4 *If $\mu \ll \lambda$ then the intersection space, $\text{int}(\mu, \lambda) = \mathcal{H}^+(\mu) \cap \mathcal{H}^+(\lambda)$ is dense in $\mathcal{H}^+(\mu)$.*

Proof We have that for all $n, 0 \leq \mu_n \leq \mu$ and $\mu_n \uparrow \mu$. Moreover, $\mathcal{H}^+(\mu_n) \subseteq \text{int}(\mu, \lambda)$ for all n . If $1 \geq g_n \geq 0, \mu$ -a.e. are the Radon–Nikodym derivatives of the μ_n with respect to μ , and $p \in \mathbb{C}[\zeta]$, let $\mathcal{C}_n := \mathcal{C}_{\mu_n}$ and let $e_n : \mathcal{H}^+(\mu_n) \hookrightarrow \mathcal{H}^+(\mu)$. Then,

$$\begin{aligned} \|\mathcal{C}_\mu p - e_n \mathcal{C}_n p\|_\mu^2 &= \|\mathcal{C}_\mu p\|_\mu^2 - 2\text{Re} \langle \mathcal{C}_\mu p, e_n \mathcal{C}_n p \rangle_\mu + \|e_n \mathcal{C}_n p\|_\mu^2 \\ &= \|p\|_{H^2(\mu)}^2 - 2\text{Re} \langle E_n p, p \rangle_{H^2(\mu_n)} + \|E_n^* p\|_{H^2(\mu)}^2 \\ &\leq 2\|p\|_{H^2(\mu)}^2 - 2\|p\|_{H^2(\mu_n)}^2 \\ &= 2 \int_{\partial\mathbb{D}} |p(\zeta)|^2 (1 - g_n(\zeta)) \mu(d\zeta) \rightarrow 0, \end{aligned}$$

by the Lebesgue monotone convergence theorem. In conclusion,

$$\mathcal{H}^+(\mu) = \bigvee_{n=1}^{\infty} \mathcal{H}^+(\mu_n),$$

where, here, \bigvee denotes closed linear span. ■

This motivates the following definitions:

Definition 2. Let μ, λ be finite, positive and regular Borel measures on $\partial\mathbb{D}$. We say that μ is *absolutely continuous with respect to λ in the reproducing kernel sense*, $\mu \ll_{RK} \lambda$, if the intersection space,

$$\text{int}(\mu, \lambda) = \mathcal{H}^+(\mu) \cap \mathcal{H}^+(\lambda)$$

is norm-dense in $\mathcal{H}^+(\mu)$.

We say that μ is *reproducing kernel singular* with respect to λ , written $\mu \perp_{RK} \lambda$, if the intersection space is trivial, $\text{int}(\mu, \lambda) = \{0\}$.

By the previous proposition, $\mu \ll \lambda$ implies that $\mu \ll_{RK} \lambda$. The main result of this section will be to show that this new ‘reproducing kernel’ definition of absolute continuity is equivalent to the classical one.

Lemma 3.5 *If $\mu \ll_{RK} \lambda$ then the embedding, $e_{\mu,\lambda} : \text{int}(\mu, \lambda) \subseteq \mathcal{H}^+(\mu) \hookrightarrow \mathcal{H}^+(\lambda)$, is closed with dense domain $\text{int}(\mu, \lambda)$. In this case, the co-embedding, $E_{\mu,\lambda} : \text{Dom } E_{\mu,\lambda} \subseteq H^2(\lambda) \hookrightarrow H^2(\mu)$, is densely-defined and closed. Both $\mathbb{C}[\zeta]$ and $\mathcal{K}_{\mathbb{D}} = \bigvee_{z \in \mathbb{D}} k_z$ are cores for $E_{\mu,\lambda}$ and $E_{\mu,\lambda} k_z = k_z, E_{\mu,\lambda} p = p$ for all $k_z \in \mathcal{K}_{\mathbb{D}}$ and $p \in \mathbb{C}[\zeta]$. The (closed) self-adjoint and positive semi-definite operator, $T_\mu := E_{\mu,\lambda}^* E_{\mu,\lambda}$ is Z^λ -Toeplitz.*

Proof Let $e := e_{\mu,\lambda}$ and observe that e is, trivially, a multiplier by the constant function 1 from $\mathcal{H}^+(\mu)$ into $\mathcal{H}^+(\lambda)$. By Proposition 1.1 and Remark 1.2, e is closed on its maximal domain, $\text{int}(\mu, \lambda)$, its adjoint acts as

$$e^* k_z^\lambda = k_z^\mu, \quad \text{and} \quad e^* k_j^\lambda = k_j^\mu,$$

on kernels and coefficient kernels and the linear spans of the point evaluation and Taylor coefficient kernels in $\mathcal{H}^+(\lambda)$ are both cores for e^* . Since $E = E_{\mu,\lambda} := \mathcal{C}_\mu^* e^* \mathcal{C}_\lambda$, E is closed, the formulas $Ep = p$ for $p \in \mathbb{C}[\zeta]$ and $Ek_z = k_z$ are easily verified, and it further follows that $\mathcal{K}_\mathbb{D}$ and $\mathbb{C}[\zeta]$ are cores for E .

To check that $T := T_\mu = E_{\mu,\lambda}^* E_{\mu,\lambda} \geq 0$ is λ -Toeplitz, consider, for any $p, q \in \mathbb{C}[\zeta]$,

$$\begin{aligned} \left\langle \sqrt{T} Z^\lambda p, \sqrt{T} Z^\lambda q \right\rangle_{H^2(\lambda)} &= \langle E\zeta \cdot p, E\zeta \cdot q \rangle_{H^2(\mu)} \\ &= \langle \zeta p, \zeta q \rangle_{H^2(\mu)} = \langle Z^\mu p, Z^\mu q \rangle_{H^2(\mu)} \\ &= \langle p, q \rangle_{H^2(\mu)} = \langle Ep, Eq \rangle_{H^2(\mu)} \\ &= \left\langle \sqrt{T} p, \sqrt{T} q \right\rangle_{H^2(\lambda)}. \end{aligned}$$

As the polynomials are a core for $E = E_{\mu,\lambda}$, this calculation holds on $\text{Dom } E = \text{Dom } \sqrt{T}$. Moreover, since $\text{Dom } E = \text{Dom } \sqrt{T}$, by polar decomposition of closed operators, and since $\mathbb{C}[\zeta]$ and $\mathcal{K}_\mathbb{D}$ are Z^λ -invariant cores for E , they are also cores for \sqrt{T} , and it follows that $\text{Dom } \sqrt{T}_\mu = \text{Dom } E$ is also Z^λ -invariant. ■

Proposition 3.6 Given $\mu, \lambda \geq 0$, if λ is extreme, then $\mu \ll_{RK} \lambda$ if and only if $\mu \ll \lambda$.

Proof Recall that λ is extreme if and only if $H^2(\lambda) = L^2(\lambda)$ so that $Z^\lambda = M_\zeta^\lambda$. In this case, the self-adjoint λ -Toeplitz operator $T_\mu \geq 0$ is Toeplitz with respect to the unitary M_ζ^λ . That is,

$$\left\langle \sqrt{T_\mu} M_\zeta^\lambda h, \sqrt{T_\mu} M_\zeta^\lambda h \right\rangle_{H^2(\lambda)} = \left\langle \sqrt{T_\mu} h, \sqrt{T_\mu} h \right\rangle_{H^2(\lambda)},$$

for all $h \in \text{Dom } \sqrt{T_\mu}$. Hence the quadratic forms for $(M_\zeta^\lambda)^* T_\mu M_\zeta^\lambda$ and T_μ are the same. By uniqueness of the unbounded Riesz representation, $(M_\zeta^\lambda)^* T_\mu M_\zeta^\lambda = T_\mu$, so that, by Lemma 1.6,

$$(I + (M_\zeta^\lambda)^* T_\mu M_\zeta^\lambda)^{-1} = (I + T_\mu)^{-1},$$

or, equivalently,

$$M_\zeta^\lambda (I + T_\mu)^{-1} = (I + T_\mu)^{-1} M_\zeta^\lambda.$$

This shows that T_μ , and hence $\sqrt{T_\mu}$ are affiliated to the commutant of the unitary operator M_ζ^λ . Since $\mathbb{C}[\zeta] \subseteq \text{Dom } \sqrt{T_\mu}$, we conclude that $\sqrt{T_\mu} = M_\zeta^\lambda \sqrt{T_\mu} 1$ acts as multiplication by $\sqrt{T_\mu} 1 =: f \in L^2(\lambda)$. Since $\sqrt{T_\mu} \geq 0$, we necessarily have that $f \geq 0$, λ -a.e., and we conclude that for any polynomials p, q ,

$$\langle p, q \rangle_{H^2(\mu)} = \left\langle \sqrt{T_\mu} p, \sqrt{T_\mu} q \right\rangle_{H^2(\lambda)} = \int_{\partial\mathbb{D}} \overline{p(\zeta)} q(\zeta) f(\zeta)^2 \lambda(d\zeta).$$

As in the proof of sufficiency in Theorem 3.1, we conclude that the above formula holds for any $g, h \in \mathbb{C}[\zeta] + \overline{\mathbb{C}[\zeta]}$, which is dense in $\mathcal{C}(\partial\mathbb{D})$ and $L^\infty(\mu)$. In particular, the formula holds for all simple functions and characteristic functions of Borel sets. Since $f \in L^2(\lambda)$, $f^2 \in L^1(\lambda)$ and it follows that

$$f^2 = \frac{\mu(d\zeta)}{\lambda(d\zeta)},$$

is the Radon–Nikodym derivative of μ with respect to λ . ■

To prove that absolute continuity in the reproducing kernel sense is equivalent to absolute continuity in general, we will appeal to B. Simon's Lebesgue decomposition theory for positive quadratic forms in Hilbert space [22], [21, Supplement to VIII.7]. Let $\hat{\mathcal{H}}(\mathfrak{q})$ be the Hilbert space completion of $\text{Dom } \mathfrak{q}$ with respect to the inner product $\langle \cdot, \cdot \rangle_{\mathcal{H}\mathfrak{C}} + \mathfrak{q}(\cdot, \cdot)$, and let $j_\mathfrak{q} : \text{Dom } \mathfrak{q} \hookrightarrow \hat{\mathcal{H}}(\mathfrak{q})$ denote the formal embedding. Further define the co-embedding $E_\mathfrak{q} : \hat{\mathcal{H}}(\mathfrak{q}) \hookrightarrow \mathcal{H}$ by

$$E_\mathfrak{q}(j_\mathfrak{q}(x)) := x, \quad x \in \text{Dom } \mathfrak{q}.$$

By construction, $j_\mathfrak{q}$ is densely-defined, has dense range, and $E_\mathfrak{q}$ is contractive with dense range in \mathcal{H} . Hence $E_\mathfrak{q}$ extends by continuity to a contraction, also denoted by $E_\mathfrak{q}$, $E_\mathfrak{q} : \hat{\mathcal{H}}(\mathfrak{q}) \hookrightarrow \mathcal{H}$.

Lemma 3.7 A densely-defined and positive semi-definite quadratic form, \mathfrak{q} , in \mathcal{H} , is closeable if and only if $j_\mathfrak{q}$ is closeable, or equivalently, if and only if $E_\mathfrak{q}$ is injective.

This lemma is a straightforward consequence of the definitions, see also [22].

Theorem 3.8 (Simon–Lebesgue decomposition of positive forms) Let $q \geq 0$ be a positive semi-definite quadratic form with dense form domain, $\text{Dom } q$, in a separable, complex Hilbert space, \mathcal{H} . Then q has a unique Lebesgue decomposition, $q = q_{ac} + q_s$, where $0 \leq q_{ac}, q_s \leq q$ in the quadratic form sense, q_{ac} is the maximal absolutely continuous form less than or equal to q and q_s is a singular form.

If P_s denotes the projection onto $\text{Ker } E_q$, and $P_{ac} = I - P_s$, then q_{ac} is given by the formula,

$$q_{ac}(x, y) = \langle j_q(x), (P_{ac} - E_q^* E_q) j_q(y) \rangle_{\mathcal{H}(q)} = \langle j_q(x), P_{ac} j_q(y) \rangle_{\mathcal{H}(q)} - \langle x, y \rangle_{\mathcal{H}}. \quad (3.2)$$

In the above theorem statement, recall that we defined the notions of an absolutely continuous or singular positive quadratic form in the introduction. Namely, a positive semi-definite and densely-defined quadratic form, $q : \text{Dom } q \times \text{Dom } q \rightarrow \mathcal{H}$, $\text{Dom } q \subseteq \mathcal{H}$, is called absolutely continuous if it is closeable, and singular if the only absolutely continuous and positive semi-definite form it dominates is the identically zero form.

Remark 3.9 If, now, $\mu, \lambda \geq 0$ are measures on the circle, we can take $\mathcal{H} := L^2(\lambda)$ or $H^2(\lambda)$, and define $q_\mu \geq 0$ on a dense form domain in \mathcal{H} by the formula

$$q_\mu(f, g) = \int_{\partial\mathbb{D}} \overline{f(\zeta)} g(\zeta) \mu(d\zeta).$$

For example, if $\mathcal{H} = L^2(\lambda)$, one can take $\text{Dom } q_\mu = \mathcal{C}(\partial\mathbb{D})$, the continuous functions. In this case, by the remark on [22, p. 381], the quadratic form Lebesgue decomposition of q_μ coincides with the classical Lebesgue decomposition of μ with respect to λ . Namely, in this case, the absolutely continuous part of q_μ , $q_{\mu;ac}$ is equal to $q_{\mu_{ac}}$, the positive form of the absolutely continuous part of μ with respect to λ , μ_{ac} , and $q_{\mu;s} = q_{\mu_s}$. In particular, $q_T := \overline{q_{ac}}$ is the form of the positive semi-definite, self-adjoint operator $T = M_f \geq 0$, where $f \in L^1(\lambda)$ is the Radon–Nikodym derivative of μ with respect to λ . This follows because, as observed by Simon, in this case his construction of the absolutely continuous and singular parts of q_μ essentially reduces to von Neumann’s functional analytic proof of the Lebesgue decomposition and Radon–Nikodym theorem in [24, Lemma 3.2.3]. See also [12, Section 5], which arrives at the same conclusion with the choice of form domain, $\text{Dom } q_\mu \subseteq L^2(\lambda)$, equal to the simple functions, i.e. linear combinations of characteristic functions of Borel sets.

Theorem 3.10 Let $q \geq 0$ be a densely-defined and positive semi-definite quadratic form in a separable complex Hilbert space, \mathcal{H} . If $q_T = \overline{q_{ac}}$ is the closure of q_{ac} , then $(I + T)^{-1} = E_q E_q^*$, where $E_q : \mathcal{H}(q) \hookrightarrow \mathcal{H}$ is the contractive co-embedding.

Lemma 3.11 Let $A : \text{Dom } A \subseteq \mathcal{H} \rightarrow \mathcal{H}$ be a densely-defined linear operator. Then A is closeable if and only if the positive semi-definite quadratic form $q_{A^*A}(x, y) := \langle Ax, Ay \rangle_{\mathcal{H}}$, with form domain $\text{Dom } q_{A^*A} := \text{Dom } A$, is closeable.

In the above statement, note that A^*A is not defined if A is not closeable.

Proof (Proof of Theorem 3.10.) Let $(x_j)_{j=1}^\infty \subseteq \text{Dom } q$ be a sequence with dense linear span. Apply Gram–Schmidt orthogonalization to (x_j) with respect to the $q + i\text{id}$ -inner product of $\mathcal{H}(q)$. This yields a countable basis $(y_j)_{j=1}^\infty \subseteq \text{Dom } q$, so that the sequence $(j_q(y_j))$ is an orthonormal basis of $\mathcal{H}(q)$. Hence,

$$E_q E_q^* = E_q I_{\mathcal{H}(q)} E_q^* = \sum_{j=1}^\infty \langle E_q j_q(y_j), \cdot \rangle_{\mathcal{H}(q)} E_q j_q(y_j) = \sum \langle y_j, \cdot \rangle_{\mathcal{H}} y_j.$$

By [22, Theorem 2.1, Corollary 2.3], see Theorem 3.8 and Equation (3.2) above,

$$q_{ac}(x, y) + \langle x, y \rangle_{\mathcal{H}} = q_{I+T}(x, y) = \left\langle \sqrt{I+T}x, \sqrt{I+T}y \right\rangle_{\mathcal{H}} = \langle j_q(x), P_{ac} j_q(y) \rangle_{\mathcal{H}(q)},$$

for any $x, y \in \text{Dom } \mathfrak{q} \subseteq \text{Dom } \mathfrak{q}_{ac} \subseteq \text{Dom } \sqrt{I+T}$. Hence, for any $x, y \in \text{Dom } T$,

$$\begin{aligned} q_{E_q E_q^*}((I+T)x, (I+T)y) &= \left\langle \sqrt{E_q E_q^*} (I+T)x, \sqrt{E_q E_q^*} (I+T)y \right\rangle_{\mathcal{H}} \\ &= \sum_{j=1}^{\infty} \langle (I+T)x, y_j \rangle_{\mathcal{H}} \langle y_j, (I+T)y \rangle_{\mathcal{H}} \\ &= \sum \left\langle \sqrt{I+T}x, \sqrt{I+T}y_j \right\rangle_{\mathcal{H}} \left\langle \sqrt{I+T}y_j, \sqrt{I+T}y \right\rangle_{\mathcal{H}} \\ &= \sum \left\langle P_{ac} j_q(x), j_q(y_j) \right\rangle_{\mathfrak{q}+i\mathfrak{d}} \left\langle j_q(y_j), P_{ac} j_q(y) \right\rangle_{\mathfrak{q}+i\mathfrak{d}} \\ &= \left\langle P_{ac} j_q(x), P_{ac} j_q(y) \right\rangle_{\mathfrak{q}+i\mathfrak{d}} = q_{I+T}(x, y) \\ &= \langle x, (I+T)y \rangle_{\mathcal{H}}. \end{aligned}$$

That is, the (closeable) quadratic forms of $I+T$ and $(I+T)E_q E_q^*(I+T)$ agree on $\text{Dom } T$, which is a core for $\sqrt{I+T}$, and a form-core for \mathfrak{q}_{I+T} . Moreover, $\mathcal{D} := \text{Ran } \sqrt{I+T} \cap \text{Dom } \sqrt{I+T}$ is a core for $\sqrt{I+T}$, so that for all $x = \sqrt{I+T}y \in \mathcal{D}$,

$$\begin{aligned} \langle x, x \rangle_{\mathcal{H}} &= \left\langle \sqrt{I+T}y, \sqrt{I+T}y \right\rangle_{\mathcal{H}} = \left\langle (I+T)y, E_q E_q^*(I+T)y \right\rangle_{\mathcal{H}} \\ &= \left\langle \sqrt{I+T}x, E_q E_q^* \sqrt{I+T}x \right\rangle_{\mathcal{H}}. \end{aligned}$$

That is, the (bounded) positive quadratic form of the identity, I , agrees with the quadratic form of $\sqrt{I+T}E_q E_q^* \sqrt{I+T}$ on the dense subspace $\text{Dom } \sqrt{I+T}$. Here, if $V := E_q^* \sqrt{I+T}$, this is a closeable operator by Lemma 3.11. In fact, V extends by continuity to an isometry, since $\mathfrak{q}_{V^*V} = \mathfrak{q}_I|_{\text{Dom } \sqrt{I+T}}$. Moreover, E_q , and hence $\sqrt{E_q E_q^*}$ have dense range, so that $\sqrt{E_q E_q^*} \sqrt{I+T}$ extends to an isometry with dense range, i.e. a unitary. For any $x, y \in \text{Dom } \sqrt{I+T}$, we have that

$$\langle x, y \rangle_{\mathcal{H}} = \left\langle \sqrt{I+T}x, E_q E_q^* \sqrt{I+T}y \right\rangle_{\mathcal{H}}.$$

Hence, by definition of the adjoint, for any $y \in \text{Dom } \sqrt{I+T}$, $E_q E_q^* \sqrt{I+T}y \in \text{Dom } \sqrt{I+T}$, and

$$\sqrt{I+T}E_q E_q^* \sqrt{I+T}y = y.$$

Hence for any $x = \sqrt{I+T}^{-1}g$ and $y = \sqrt{I+T}^{-1}h$,

$$\begin{aligned} \langle g, (I+T)^{-1}h \rangle_{\mathcal{H}} &= \langle x, y \rangle_{\mathcal{H}} = \left\langle E_q^* \sqrt{I+T}x, E_q^* \sqrt{I+T}y \right\rangle_{\mathcal{H}} \\ &= \langle g, E_q E_q^* h \rangle_{\mathcal{H}}, \end{aligned}$$

and we conclude, by the Riesz lemma for bounded sesquilinear forms, that

$$(I+T)^{-1} = E_q E_q^*.$$

■

Theorem 3.12 Let μ, λ be positive, finite and regular Borel measures on $\partial\mathbb{D}$. Then $\mu \ll \lambda$ if and only if $\mu \ll_{RK} \lambda$.

In this case, the co-embedding, $E_{\mu, \lambda} : \text{Dom } E_{\mu, \lambda} \subseteq H^2(\lambda) \hookrightarrow H^2(\mu)$ is closed, and its domain is Z^λ -invariant. Both $\mathbb{C}[\zeta]$ and $\mathcal{K}_{\mathbb{D}} = \vee k_z$ are cores for $E_{\mu, \lambda}$, $E_{\mu, \lambda} p = p$ and $E_{\mu, \lambda} k_z = k_z$ for any $p \in \mathbb{C}[\zeta]$ or Szegő kernel k_z . The self-adjoint and positive semi-definite operator, $T_\mu := E_{\mu, \lambda}^* E_{\mu, \lambda}$, is λ -Toeplitz and

$$T_\mu = P_{H^2(\lambda)} M_f|_{H^2(\lambda)}; \quad f := \frac{d\mu}{d\lambda},$$

in the quadratic form sense, i.e.

$$\left\langle \sqrt{T_\mu} p, \sqrt{T_\mu} q \right\rangle_{H^2(\lambda)} = \left\langle M_{\sqrt{f}} p, M_{\sqrt{f}} q \right\rangle_{L^2(\lambda)}; \quad \forall p, q \in \mathbb{C}[\zeta].$$

Proof Necessity was established in Proposition 3.4 and sufficiency, in the case where λ is extreme, was proven in Proposition 3.6. To prove sufficiency in general, assume that $\mu \ll_{RK} \lambda$ so that the intersection space $\text{int}(\mu, \lambda)$ is dense in the space of μ -Cauchy transforms, $\mathcal{H}^+(\mu)$. Note that $\mu \ll_{RK} \lambda$ if and only if $\mu + \lambda \ll_{RK} \lambda$ (and the same is true for \ll). This follows from Aronszajn's 'sums of kernels' theorem as stated in Subsection 1.2. By Lemma 3.5, $e_{\mu+\lambda, \lambda} : \text{int}(\mu + \lambda, \lambda) \subseteq$

$\mathcal{H}^+(\mu + \lambda) \hookrightarrow \mathcal{H}^+(\lambda)$ is a closed embedding, $E_{\mu+\lambda,\lambda} := \mathcal{C}_{\mu+\lambda}^* e_{\mu+\lambda,\lambda}^* \mathcal{C}_\lambda$ is a closed co-embedding with $\mathcal{K}_\mathbb{D} = \bigvee k_z$ and $\mathbb{C}[\zeta]$ as cores, and $T := T_{\mu+\lambda} = E_{\mu+\lambda,\lambda}^* E_{\mu+\lambda,\lambda} \geq 0$ is self-adjoint, positive semi-definite, densely-defined and λ -Toeplitz.

By Remark 3.9 and Theorem 3.10 above if $\hat{E} : L^2(\mu + \lambda) \hookrightarrow L^2(\lambda)$ is the contractive co-embedding, and $\hat{T} := M_f$, where $f \geq 0$ λ -a.e., $f \in L^1(\lambda)$ is the self-adjoint, positive semi-definite multiplication operator by the Radon–Nikodym derivative, $f = \frac{d\mu}{d\lambda}$, then $(I + \hat{T})^{-1} = \hat{E}\hat{E}^*$. On the other hand, $E := E_{\mu+\lambda,\lambda} : \text{Dom } E_{\mu+\lambda,\lambda} \subseteq H^2(\lambda) \hookrightarrow H^2(\mu + \lambda)$ is closed and bounded below by 1, and so has a contractive inverse, namely, for any $h \in \mathcal{K}_\mathbb{D}$ or in $\mathbb{C}[\zeta]$,

$$\hat{E}Eh = h \in H^2(\lambda), \quad \text{and} \quad E\hat{E}h = h \in H^2(\mu + \lambda).$$

Hence,

$$\hat{E}E|_{\text{Dom } E} = I_{H^2(\lambda)}|_{\text{Dom } E}, \quad \text{and} \quad E\hat{E}|_{H^2(\mu+\lambda)} = I_{\mu+\lambda}.$$

Observe that the contractive co-embedding, \hat{E} is necessarily injective and has dense range, so that it has a closed, potentially unbounded inverse, \hat{E}^{-1} , which is densely-defined. Since $(I + \hat{T})^{-1} = \hat{E}\hat{E}^*$, we conclude that $I + \hat{T} = \hat{E}^{-*}\hat{E}^{-1}$. Hence, for any $p, q \in \mathbb{C}[\zeta] \subseteq H^2(\lambda)$,

$$\begin{aligned} \left\langle \sqrt{I + \hat{T}}p, \sqrt{I + \hat{T}}q \right\rangle_{L^2(\lambda)} &= \left\langle \hat{E}^{-1}p, \hat{E}^{-1}q \right\rangle_{L^2(\mu+\lambda)} \\ &= \langle p, q \rangle_{H^2(\mu+\lambda)} = \langle Ep, Eq \rangle_{H^2(\mu+\lambda)} \\ &= \left\langle \sqrt{T}p, \sqrt{T}q \right\rangle_{H^2(\lambda)}. \end{aligned}$$

This calculation shows that the ‘compression’ of $I + \hat{T}$ to the intersection of its domain with the subspace $H^2(\lambda)$ is equal to T , in this quadratic form sense. In particular $T - I \geq 0$ is the compression of $\hat{T} = M_f$ to $H^2(\lambda)$, where $f \in L^1(\lambda)$ is the Radon–Nikodym derivative of μ with respect to λ . In conclusion, for any polynomials p, q ,

$$\begin{aligned} \int_{\partial\mathbb{D}} \overline{p(\zeta)}q(\zeta)\mu(d\zeta) &= \langle p, q \rangle_{H^2(\mu)} \\ &= \left\langle \sqrt{T}p, \sqrt{T}q \right\rangle_{H^2(\lambda)} - \langle p, q \rangle_{H^2(\lambda)} \\ &= \left\langle \sqrt{\hat{T}}p, \sqrt{\hat{T}}q \right\rangle_{L^2(\lambda)} \\ &= \left\langle M_{\sqrt{f}}p, M_{\sqrt{f}}q \right\rangle_{L^2(\lambda)} \\ &= \int_{\partial\mathbb{D}} \overline{p(\zeta)}q(\zeta)f(\zeta)\lambda(d\zeta). \end{aligned}$$

As in the second proof of sufficiency of Theorem 3.1, this equality can be extended to arbitrary $g, h \in \mathbb{C}[\zeta] + \overline{\mathbb{C}[\zeta]}$, so that by Weierstraß approximation, $\mu \ll \lambda$ with Radon–Nikodym derivative $f \geq 0$, $f \in L^1(\lambda)$. ■

4 Lebesgue decomposition via reproducing kernels

By Theorem 3.12, our definition of reproducing kernel absolute continuity is equivalent to the classical definition of absolute continuity for finite, positive and regular Borel measures on the complex unit circle. In particular, if $\mu \ll \lambda$, it follows that the intersection space of μ and λ -Cauchy transforms is dense in the space of μ -Cauchy transforms. Hence, if $\mu = \mu_{ac} + \mu_s$ is the Lebesgue decomposition of μ with respect to λ , then $\text{int}(\mu_{ac}, \lambda)$ is dense in $\mathcal{H}^+(\mu_{ac})$, and since $\mu \geq \mu_{ac}$, $\text{int}(\mu_{ac}, \lambda) \subseteq \text{int}(\mu, \lambda)$. That is, if $\mu_{ac} \neq 0$, it follows that $\text{int}(\mu, \lambda) \neq \{0\}$ is not trivial. This raises several natural questions: How can we identify the space of μ_{ac} -Cauchy transforms? Is $\text{int}(\mu, \lambda)^{-\mu} := \text{int}(\mu, \lambda)^{-1} \cdot \|\cdot\|_\mu$ equal to the space of μ_{ac} -Cauchy transforms? We will see that the answer to the second question is positive if λ is non-extreme, but that in general, $\text{int}(\mu, \lambda)^{-\mu}$ is not the space of Cauchy transforms of any positive measure, see Corollary 4.15 and Example 4.17.

Theorem 4.1 *If \mathcal{M} is a RKHS in \mathbb{D} that embeds contractively in $\mathcal{H}^+(\mu)$, then $\mathcal{M} = \mathcal{H}^+(\gamma)$ for a positive measure, $\gamma, \gamma \leq \mu$, if and only if $e : \mathcal{M} \hookrightarrow \mathcal{H}^+(\mu)$ is such that the positive semi-definite contraction $\tau := ee^*$ is V_μ -Toeplitz.*

In this case, $\mathcal{M} = \mathcal{H}^+(\gamma) = \mathcal{R}(e)$, and the complementary space of $\mathcal{H}^+(\gamma)$ in $\mathcal{H}^+(\mu)$ is $\mathcal{H}^+(\nu)$, for a positive measure, ν , where $k^\nu = k^\mu - k^\gamma$ so that $\mu = \gamma + \nu$.

Proof First, if $\mathcal{H}^+(\gamma) = \mathcal{M} \subseteq \mathcal{H}^+(\mu)$ is contractively contained, then, $e : \mathcal{H}^+(\gamma) \hookrightarrow \mathcal{H}^+(\mu)$ is trivially a (contractive) multiplier so that, as before, $e^*k_z^\mu = k_z^\gamma$, and

$$e^*V^\mu \bar{=} V^\gamma e^*.$$

In conclusion,

$$V^{\mu*}ee^*V^{\mu} = eV^{\gamma*}V^{\gamma}e^* = ee^* = \tau.$$

Conversely, if $\tau = ee^*$ is V^{μ} -Toeplitz and contractive, then, as in the proof of Theorem 3.1, $T := \mathcal{E}_{\mu}^*ee^*\mathcal{E}_{\mu}$ is a contractive Z^{μ} -Toeplitz operator and we can appeal to the Riesz–Markov theorem to show that there is a $\gamma \geq 0$, so that $T = P_{H^2(\mu)}M_f|_{H^2(\mu)}$, where $f \geq 0$, $\|f\|_{\infty} \leq 1$ is the Radon–Nikodym derivative of γ with respect to μ . Namely, one can define a linear functional, $\dot{\mu}_T$, on $\mathbb{C}[\zeta] + \overline{\mathbb{C}[\zeta]} \subseteq \mathcal{E}(\partial\mathbb{D})$, by

$$\dot{\mu}_T(p + \bar{q}) := \langle 1, Tp \rangle_{H^2(\mu)} + \langle q, T1 \rangle_{H^2(\mu)}.$$

It is easy to check that $\dot{\mu}_T$ is bounded and positive using the Fejér–Riesz theorem, as in the proof of Theorem 3.1. The fact that T is a positive semi-definite Z^{μ} -Toeplitz contraction ensures that $\dot{\mu}_T$ extends to a bounded, positive linear functional on $\mathcal{E}(\partial\mathbb{D})$, and that $\dot{\mu}_T \leq \dot{\mu}$, so that $\dot{\mu}_T = \dot{\gamma}$ for some finite, regular and positive Borel measure, $\gamma \leq \mu$, by the Riesz–Markov theorem.

By Theorem 1.4, the complementary space, $\mathcal{R}^c(e)$, of $\mathcal{R}(e) = \mathcal{H}^+(\gamma)$ is a RKHS in \mathbb{D} with reproducing kernel $k'(z, w) = k^{\mu}(z, w) - k^{\gamma}(z, w)$ and it is contractively contained in $\mathcal{H}^+(\mu)$, by the inclusion theorem. Moreover, if $j : \mathcal{R}^c(e) \hookrightarrow \mathcal{H}^+(\mu)$ is the contractive embedding, then it follows that $jj^* = I - ee^* \geq 0$ is also a positive semi-definite V^{μ} -Toeplitz contraction. Hence, by the first part of the proof, $\mathcal{R} = \mathcal{H}^+(\nu)$ for a positive measure, ν . Finally, since $k^{\mu} = k^{\gamma} + k^{\nu}$, we obtain that $\mu = \gamma + \nu$. ■

Lemma 4.2 Given any μ, λ , the intersection space $\text{int}(\mu, \lambda)$, is both V_{λ} and V_{μ} -co-invariant, and $V_{\lambda}^*|_{\text{int}(\mu, \lambda)} = V_{\mu}^*|_{\text{int}(\mu, \lambda)}$.

Proof This is immediate, by Lemma 2.4, since both V_{μ}^* and V_{λ}^* act as ‘backward shifts’ on power series. ■

Lemma 4.3 If λ is non-extreme, then the intersection space, $\text{int}(\mu, \lambda)$, is V_{μ} -reducing. If λ is extreme, then $\text{int}(\mu, \lambda)$ is V_{μ} -reducing (and V_{λ} -reducing) if and only if $V_{\mu}|_{\text{int}(\mu, \lambda)} = V_{\lambda}|_{\text{int}(\mu, \lambda)}$.

Proof By Lemma 2.4, if $h \in \text{int}(\mu, \lambda)$, then $V_{\mu}h \in \mathcal{H}^+(\mu)$ and

$$(V_{\mu}h)(z) = zh(z) + (V_{\mu}h)(0)1 = (V_{\lambda}h)(z) - (V_{\lambda}h)(0)1 + (V_{\mu}h)(0)1. \quad (4.1)$$

Hence, if $c := (V_{\lambda}h)(0) - (V_{\mu}h)(0) \in \mathbb{C}$ and λ is non-extreme, then both $V_{\lambda}h$ and $c1$ belong to $\mathcal{H}^+(\lambda)$ so that $V_{\mu}h \in \mathcal{H}^+(\lambda) \cap \mathcal{H}^+(\mu) = \text{int}(\mu, \lambda)$. Recall that λ is extreme if and only if $\mathcal{H}^+(\lambda)$ does not contain the constant functions. Hence, if λ is extreme then $\text{int}(\mu, \lambda)$ will be V_{μ} -reducing if and only if $(V_{\mu}h)(0) = (V_{\lambda}h)(0)$ for all $h \in \text{int}(\mu, \lambda)$. By Equation (4.1), this happens if and only if $V_{\mu}h = V_{\lambda}h$. ■

Corollary 4.4 If $\mathcal{M} \subseteq \mathcal{H}^+(\mu)$ is a V_{μ} -reducing subspace, then $\mathcal{M} = \mathcal{H}^+(\gamma)$ for some $\gamma \leq \mu$. Moreover, $\mathcal{M}^{\perp} = \mathcal{H}^+(\gamma')$ for some $\mu \geq \gamma' \geq 0$ so that $\gamma + \gamma' = \mu$.

Proof Let P be the orthogonal projection of $\mathcal{H}^+(\mu)$ onto \mathcal{M} . Then if $e : \mathcal{M} \hookrightarrow \mathcal{H}^+(\mu)$ is the isometric embedding, $P = ee^*$. Hence,

$$V_{\mu}^*ee^*V_{\mu} = V_{\mu}^*PV_{\mu} = V_{\mu}^*V_{\mu}P = P = ee^*,$$

so that $\tau = ee^*$ is V^{μ} -Toeplitz and $\mathcal{M} = \mathcal{H}^+(\gamma)$ for some $0 \leq \gamma \leq \mu$, and $\mathcal{M}^{\perp} = \mathcal{H}^+(\gamma')$, by Theorem 4.1. ■

Theorem 4.5 Let $\mu, \lambda \geq 0$ be finite, positive and regular Borel measures on $\partial\mathbb{D}$. If the intersection space, $\text{int}(\mu, \lambda)$, is V^{μ} -reducing and $\mu = \mu_{ac} + \mu_s$ is the Lebesgue decomposition of μ with respect to λ , then

$$\mathcal{H}^+(\mu) = \mathcal{H}^+(\mu_{ac}) \oplus \mathcal{H}^+(\mu_s).$$

In this case,

$$\mathcal{H}^+(\mu_{ac}) = \text{int}(\mu, \lambda)^{-\mu}, \quad \text{and} \quad \mathcal{H}^+(\mu_s) \cap \mathcal{H}^+(\lambda) = \{0\}.$$

That is, μ_{ac} is the largest positive measure $\leq \mu$ which is RK-ac with respect to λ , and μ_s is RK-singular with respect to λ .

In particular, $\text{int}(\mu, \lambda)$ will be V^{μ} -reducing if λ is non-extreme by Lemma 4.3.

Proof By Theorem 3.12, we have that $\text{int}(\mu_{ac}, \lambda) \subseteq \text{int}(\mu, \lambda)$ is dense in $\mathcal{H}^+(\mu_{ac})$. Since we assume that $\text{int}(\mu, \lambda)$ is V_{μ} -reducing, its closure, $\text{int}(\mu, \lambda)^{-\mu}$, is also V_{μ} -reducing and then $\text{int}(\mu, \lambda)^{-\mu} = \mathcal{H}^+(\gamma)$ for some $0 \leq \gamma \leq \mu$ by Corollary 4.4. By construction $\gamma \ll_{RK} \lambda$ so that $\gamma \ll \lambda$ by Theorem 3.12. By maximality, $\gamma \leq \mu_{ac}$ and by construction $\text{int}(\mu, \lambda) \subseteq \text{int}(\gamma, \lambda)$. However, we also have that $\text{int}(\mu_{ac}, \lambda) \subseteq \text{int}(\mu, \lambda) \subseteq \text{int}(\gamma, \lambda)$. Hence if $e : \mathcal{H}^+(\mu_{ac}) \hookrightarrow \mathcal{H}^+(\mu)$

is the contractive embedding, then e , restricted to the dense subspace $\text{int}(\mu_{ac}, \lambda) \subseteq \mathcal{H}^+(\mu_{ac})$ defines a contraction into $\mathcal{H}^+(\gamma)$, which is isometrically contained in $\mathcal{H}^+(\mu)$. It follows that e extends by continuity to a contractive embedding of $\mathcal{H}^+(\mu_{ac})$ into $\mathcal{H}^+(\gamma)$. Hence, by Theorem 3.1, $\mu_{ac} \leq \gamma$ and we conclude that $\mu_{ac} = \gamma$. ■

Corollary 4.6 *Let $\mu, \lambda \geq 0$ be finite, positive and regular Borel measures on $\partial\mathbb{D}$. If $\text{int}(\mu, \lambda) = \{0\}$ so that $\mu \perp_{RK} \lambda$, then $\mu \perp \lambda$. If $\text{int}(\mu, \lambda)$ is either V_μ or V_λ -reducing then $\mu \perp \lambda$ if and only if $\mu \perp_{RK} \lambda$. In particular, if either μ or λ is non-extreme then $\mu \perp \lambda$ if and only if $\mu \perp_{RK} \lambda$.*

While the previous two results give quite a satisfactory description of the Lebesgue decomposition of μ with respect to λ in terms of reproducing kernel theory in the case where the intersection space, $\text{int}(\mu, \lambda)$ is V^μ -reducing, this is not generally the case, as the next proposition and example show.

In the proposition statement below, recall the definition of the lattice operations \vee, \wedge on positive kernels and the definition of the isometries $U_\vee : \mathcal{H}(K + k) \rightarrow \mathcal{H}(K) \oplus \mathcal{H}(k)$ and $U_\wedge : \mathcal{H}(K \wedge k) = \mathcal{H}(K) \cap \mathcal{H}(k) \rightarrow \mathcal{H}(K) \oplus \mathcal{H}(k)$, see Subsection 1.2.

Proposition 4.7 *If $\mu + \lambda$ is extreme, then $\text{int}(\mu, \lambda)$ is V_μ -reducing. If μ, λ are both extreme but $\mu + \lambda$ is non-extreme, then $\text{int}(\mu, \lambda)$ is non-trivial, but not V_μ -reducing and*

$$\text{int}(\mu, \lambda) \supseteq U_\wedge^* \bigvee_{j=1}^{\infty} (V_\mu^* \oplus V_\lambda^*)^j U_\vee 1.$$

Lemma 4.8 *Let $\mu, \lambda \geq 0$ be positive, finite and regular Borel measures on $\partial\mathbb{D}$. Consider the reproducing kernel Hilbert spaces of μ, λ and $\mu + \lambda$ -Cauchy transforms in $\partial\mathbb{D}$, $\mathcal{H}^+(\mu) = \mathcal{H}(k^\mu)$, $\mathcal{H}^+(\lambda) = \mathcal{H}(k^\lambda)$ and $\mathcal{H}^+(\mu + \lambda) = \mathcal{H}(k^\mu + k^\lambda)$. Then,*

$$U_\vee V_{\mu+\lambda} = V_\mu \oplus V_\lambda U_\vee$$

and $\text{Ran } U_\vee$ is $V_\mu \oplus V_\lambda$ -invariant so that $\text{Ran } U_\wedge$ is $V_\mu \oplus V_\lambda$ co-invariant, and

$$U_\wedge V^{\mu*} |_{\text{int}(\mu, \lambda)} = U_\wedge V^{\lambda*} |_{\text{int}(\mu, \lambda)} = (V^\mu \oplus V^\lambda)^* U_\wedge.$$

Moreover, we have that $V_\mu |_{\text{int}(\mu, \lambda)} = V_\lambda |_{\text{int}(\mu, \lambda)}$ so that $\text{int}(\mu, \lambda) = \mathcal{H}(k^\mu \wedge k^\lambda)$ is both V_μ and V_λ -invariant if and only if $\text{Ran } U_\wedge$ is $V_\mu \oplus V_\lambda$ -invariant.

Proof The intertwining formulas are easily verified. The range of $U_\wedge : \mathcal{H}^+(\mu + \lambda) \rightarrow \mathcal{H}^+(\mu) \oplus \mathcal{H}^+(\lambda)$ is $V_\mu \oplus V_\lambda$ -reducing if and only if, for any $h \oplus -h \in \text{Ran } U_\wedge$, $h \in \text{int}(\mu, \lambda)$,

$$V_\mu h \oplus -V_\lambda h = g \oplus -g,$$

for some $g \in \text{int}(\mu, \lambda)$. Clearly this happens if and only if $V_\mu |_{\text{int}(\mu, \lambda)} = V_\lambda |_{\text{int}(\mu, \lambda)}$. ■

Proof (Proof of Proposition 4.7.) If $\mu + \lambda$ is extreme then $\mathcal{H}^+(\mu + \lambda) = \mathcal{H}^+(\mu) + \mathcal{H}^+(\lambda)$ does not contain the constant functions. Hence both μ and λ must also be extreme. In this case V_μ, V_λ and $V_{\mu+\lambda}$ are all unitary operators. We know that $\text{Ran } U_\vee$ is always $V_\mu \oplus V_\lambda$ -invariant. On the other hand since $\mu + \lambda$ is extreme, $V_{\mu+\lambda}$ is unitary, hence surjective, and

$$\mathcal{H}^+(\mu + \lambda) = \bigvee (k_z^{\mu+\lambda} - k_0^{\mu+\lambda}),$$

so that

$$\text{Ran } U_\vee = \bigvee (k_z^\mu - k_0^\mu) \oplus (k_z^\lambda - k_0^\lambda).$$

Hence,

$$(V_\mu^* \oplus V_\lambda^*) \text{Ran } U_\vee = \bigvee k_z^\mu \bar{z} \oplus k_z^\lambda \bar{z} \subseteq \text{Ran } U_\vee.$$

It follows that $\text{Ran } U_\vee$ is $V_\mu \oplus V_\lambda$ -reducing, so that $\text{Ran } U_\wedge = \text{Ran } U_\vee^\perp$ is also reducing. The previous lemma now implies that $\text{int}(\mu, \lambda)$ is V_μ -reducing.

If, on the other hand, μ, λ are both extreme but $\mu + \lambda$ is not, then V^μ, V^λ are both unitary but $V^{\mu+\lambda}$ is not. Hence, since $1 \perp \text{Ran } V^{\mu+\lambda}$, $1 \in \mathcal{H}^+(\mu + \lambda)$, we have that

$$U_\vee 1 \perp V^\mu \oplus V^\lambda \text{Ran } U_\vee,$$

or, equivalently,

$$(V^\mu \oplus V^\lambda)^* U_\vee 1 \perp \text{Ran } U_\vee.$$

Since $V^\mu \oplus V^\lambda$ is unitary, it follows that

$$0 \neq (V^\mu \oplus V^\lambda)^* U_V 1 \in \text{Ran } U_\wedge,$$

so that $\text{int}(\mu, \lambda) \neq \{0\}$. Since $\text{Ran } U_V$ is not $V^\mu \oplus V^\lambda$ -reducing, neither is $\text{Ran } U_\wedge$, and hence $\text{int}(\mu, \lambda)$ is not V_μ -reducing by the previous lemma. ■

Example 4.9 (Lebesgue measure on the half circles.) Let m_\pm be normalized Lebesgue measure restricted to the upper and lower half-circles. Then $m = m_+ + m_-$, and $m_+ \perp m_-$. Note that both m_\pm are extreme since $\frac{dm_\pm}{dm} = \chi_{\partial\mathbb{D}_\pm}$, where χ_Ω denotes the characteristic function of a Borel set, Ω , is not log-integrable (with respect to m). On the other hand, m is non-extreme. By the previous proposition, $\text{int}(m_+, m_-) \neq \{0\}$ is non-trivial, and yet $m_+ \perp m_-$. If $\text{int}(m_+, m_-)$ contained a non-trivial $V_+ := V^{m_+}$ or $V_- := V^{m_-}$ -reducing subspace, \mathcal{M} , then the closure, \mathcal{M}^+ or \mathcal{M}^- in the norms of $\mathcal{H}^+(m_\pm)$ would be a closed V_+ or V_- -reducing subspace. In the first case, Corollary 4.4 would then imply that $\mathcal{M}^+ = \mathcal{H}^+(\gamma)$ for some $0 \leq \gamma \leq m_+$. On the other hand, $\text{int}(\gamma, m_-) \supseteq \mathcal{M}$ is dense in $\mathcal{M}^+ = \mathcal{H}^+(\gamma)$ so that $\gamma \ll_{RK} m_-$. Since RK-absolute continuity is equivalent to absolute continuity by Theorem 3.12, this contradicts the mutual singularity of m_+ and m_- . A symmetric argument shows that $\text{int}(m_+, m_-)$ cannot contain a non-trivial V_- -reducing subspace either.

Similarly, $m = m_+ + m_-$ can be viewed as the Lebesgue decomposition of m with respect to m_+ . In this case, $\text{int}(m, m_+) = \mathcal{H}^+(m_+) \neq \{0\}$ since $m_+ \leq m$. However, $\text{int}(m, m_+)$ cannot be $S = V_m$ -reducing as then its closure, $\text{int}(m, m_+)^{-m}$ would be a closed, S -reducing subspace of $H^2 = \mathcal{H}^+(m)$ and the shift has no non-trivial reducing subspaces. (Hence this intersection space cannot contain any non-trivial S -reducing subspace.) In fact, $\text{int}(m, m_+)$ cannot (contractively) contain the space of γ -Cauchy transforms of any non-zero positive measure, γ , as then $\gamma \ll_{RK} m_+$ and $\gamma \ll_{RK} m_-$, so that $\gamma \ll m_+, m_-$ by Theorem 3.12 and $\gamma \equiv 0$ since m_+ and m_- are mutually singular. Finally, we cannot have $\text{int}(m, m_+)$ dense in H^2 either as this would imply that $m \ll_{RK} m_+$ which would imply that $m \ll m_+$ by Theorem 3.12.

We can calculate some vectors in $\text{int}(m_+, m_-)$ more explicitly. By the proof of Proposition 4.7, we have that $V_+^* \oplus V_-^* U_V 1 \in \text{Ran } U_\wedge$, and since $\text{Ran } U_\wedge$ is always $V_+ \oplus V_-$ co-invariant,

$$\text{int}(m_+, m_-) \supseteq \bigvee_{j=1}^{\infty} V_+^{*j} k_0^+ = \bigvee V_-^{*j} k_0^-.$$

Here, $1 = k_0^m$, where $m = m_+ + m_-$, so that $U_V 1 = k_0^+ \oplus k_0^-$. Since the unitaries V_\pm^* both act as backward shifts on power series, we can compute these elements of the intersection space explicitly. First, the kernel vectors of $\mathcal{H}^+(m_\pm)$ at 0 are:

$$\begin{aligned} k_0^+(z) &= \frac{1}{2\pi} \int_0^\pi \frac{1}{1 - ze^{-i\theta}} d\theta \\ &= \frac{1}{2\pi i} \log(e^{i\theta} - z) \Big|_{\theta=0}^{\theta=\pi} \\ &= \frac{1}{2\pi i} \log\left(\frac{z+1}{z-1}\right), \end{aligned}$$

where \log is the branch of the logarithm fixed by the choice of the argument function taking values in $[0, 2\pi)$. Here, the branch cut is along the positive real axis, and

$$\text{Re} \frac{z+1}{z-1} = \frac{|z|^2 - 1}{|z-1|^2} < 0,$$

is strictly negative for any $z \in \mathbb{D}$ so that this formula defines a holomorphic function in \mathbb{D} . (We know, of course, that k_0^+ must be holomorphic in \mathbb{D} .) Since

$$1 = k_0(z) = k(z, 0) = k^+(z, 0) + k^-(z, 0) = k_0^+(z) + k_0^-(z),$$

it follows that

$$k_0^-(z) = 1 - k_0^+(z) = 1 - \frac{1}{2\pi i} \log\left(\frac{z+1}{z-1}\right). \quad (4.2)$$

Also note that

$$\frac{1}{2} = \frac{1}{i2\pi} \log(-1),$$

so that $\frac{1}{2} = k_0^+(0) = k_0^-(0)$.

Since V_{\pm}^* act as backward shifts on power series, it follows that $V_+^*k_0^+ = -V_-^*k_0^-$, so that

$$(V_+ \oplus V_-)^*k_0^+ \oplus k_0^- \in \text{Ran } U_{\wedge} = \bigvee_{h \in \text{int}(m_+, m_-)} h \oplus -h,$$

as required.

4.1 Lebesgue decomposition of measures and their forms

As described in Remark 3.9 and Subsection 1.3, if $\mu, \lambda \geq 0$ are positive, finite and regular Borel measures on the unit circle, $\partial\mathbb{D}$, then one can construct the Lebesgue decomposition of μ with respect to λ by considering the densely-defined positive quadratic form, $\mathfrak{q}_{\mu} : \mathcal{E}(\partial\mathbb{D}) \times \mathcal{E}(\partial\mathbb{D}) \rightarrow 0$, with dense form domain $\mathcal{E}(\partial\mathbb{D}) \subseteq L^2(\lambda)$, the continuous functions on the unit circle. Namely, applying the Simon–Lebesgue decomposition to \mathfrak{q}_{μ} , viewed as a positive, densely-defined form in $L^2(\lambda)$, one obtains,

$$\mathfrak{q}_{\mu} = \mathfrak{q}_{\mu;ac} + \mathfrak{q}_{\mu;s},$$

where $\mathfrak{q}_{\mu;ac}$ is an absolutely continuous (closeable) form and \mathfrak{q}_s is a singular form and moreover, $\mathfrak{q}_{\mu;ac} = \mathfrak{q}_{\mu_{ac}}$, $\mathfrak{q}_s = \mathfrak{q}_{\mu_s}$, where $\mu = \mu_{ac} + \mu_s$ is the Lebesgue decomposition.

However, in this paper, since we wish to apply analytic and function theoretic methods, we instead consider the positive quadratic Z^{λ} -Toeplitz form, \mathfrak{q}_{μ} , associated to $\mu \geq 0$, with dense form domain $\text{Dom } \mathfrak{q}_{\mu} = \mathbb{C}[\zeta]$ or $\text{Dom } \mathfrak{q}_{\mu} = A(\mathbb{D})$, in $H^2(\lambda) \subseteq L^2(\lambda)$. As we will show, if $\mathfrak{q}_{\mu} = \mathfrak{q}_{ac} + \mathfrak{q}_s$ is the Simon–Lebesgue form decomposition of \mathfrak{q}_{μ} in $H^2(\lambda)$, then one can define reproducing kernel Hilbert spaces of \mathfrak{q}_{ac} and \mathfrak{q}_s -Cauchy transforms, $\mathcal{H}^+(\mathfrak{q}_{ac})$ and $\mathcal{H}^+(\mathfrak{q}_s)$. The goal of this subsection is to compare the Lebesgue decomposition of μ with respect to λ with the Simon–Lebesgue decomposition of \mathfrak{q}_{μ} in $H^2(\lambda)$.

Let $\mu, \lambda \geq 0$ be finite and regular Borel measures on $\partial\mathbb{D}$. Consider the positive quadratic form, \mathfrak{q}_{μ} , with dense form domain, $A(\mathbb{D}) \subseteq H^2(\lambda)$. Observe that $\hat{\mathcal{H}}(\mathfrak{q}_{\mu}) = H^2(\mu + \lambda)$ so that $\mathbb{C}[\zeta]$ and $\mathcal{K}_{\mathbb{D}}$ are both dense sets in this space. Consider the Simon–Lebesgue decomposition, $\mathfrak{q}_{\mu} = \mathfrak{q}_{ac} + \mathfrak{q}_s$, of \mathfrak{q}_{μ} in $H^2(\lambda)$. By Theorem 3.8, $\mathfrak{q}_{ac} \geq 0$, is the largest closeable quadratic form bounded above by \mathfrak{q}_{μ} . Since $\mathfrak{q}_{ac} \leq \mathfrak{q}_{\mu}$, this implies that $\text{Dom } \mathfrak{q}_{\mu} = A(\mathbb{D}) \subseteq \text{Dom } \mathfrak{q}_{ac}$, and if $\mathfrak{q}_D = \overline{\mathfrak{q}_{ac}}$ denotes the closure of \mathfrak{q}_{ac} , then $A(\mathbb{D})$ must be a form-core for the closed form \mathfrak{q}_D by the maximality statement in Theorem 3.8. We define $H^2(\mathfrak{q}_{ac}), H^2(\mathfrak{q}_s)$ as the Hilbert space completion of the disk algebra, $A(\mathbb{D})$, modulo vectors of zero length, with respect to the pre-inner products, $\mathfrak{q}_{ac}, \mathfrak{q}_s$, respectively. Since $0 \leq \mathfrak{q}_{ac}, \mathfrak{q}_s \leq \mathfrak{q}_{\mu}$, we can define the contractive co-embeddings $E_{ac} : H^2(\mu) \hookrightarrow H^2(\mathfrak{q}_{ac})$ and $E_s : H^2(\mu) \hookrightarrow H^2(\mathfrak{q}_s)$ by $E_{ac}a = a \in H^2(\mathfrak{q}_{ac})$ and $E_s a = a \in H^2(\mu_s)$. (Here, an element $a \in A(\mathbb{D})$ could be equal to 0 as an element of $H^2(\mu)$, or as an element of the spaces $H^2(\mathfrak{q}_{ac}), H^2(\mathfrak{q}_s)$. However, the inequality $0 \leq \mathfrak{q}_{ac}, \mathfrak{q}_s \leq \mathfrak{q}_{\mu}$, ensures that if $a \in A(\mathbb{D})$ is zero as an element of $H^2(\mu)$, i.e. it vanishes μ -a.e., then $a = 0$ as element of both $H^2(\mathfrak{q}_{ac})$ and $H^2(\mathfrak{q}_s)$. A more precise notation would be to let N_{ac} denote the subspace of all elements of $A(\mathbb{D})$ of zero-length with respect to the \mathfrak{q}_{ac} -pre-inner product so that equivalence classes of the form $a + N_{ac}, a \in A(\mathbb{D})$, are dense in $H^2(\mathfrak{q}_{ac})$.)

Observe that if $\mathcal{D} \subseteq A(\mathbb{D})$ is any supremum-norm dense set, such as $\mathcal{K}_{\mathbb{D}} = \bigvee k_z$ or $\mathbb{C}[\zeta]$, then \mathcal{D} is dense in $H^2(\mu)$, and since the co-embedding $E_{ac} : H^2(\mu) \hookrightarrow H^2(\mathfrak{q}_{ac})$ is a contraction with dense range, \mathcal{D} will be dense in $H^2(\mathfrak{q}_{ac})$ and it will be similarly dense in $H^2(\mathfrak{q}_s)$.

Lemma 4.10 *If $\mathfrak{q}_D = \overline{\mathfrak{q}_{ac}}$ is the closure of \mathfrak{q}_{ac} , and $\mathcal{D} \subseteq A(\mathbb{D})$ is supremum-norm dense, then \mathcal{D} is a core for $\sqrt{\mathfrak{q}_D}$.*

Proof Since $\text{Dom } \mathfrak{q}_{\mu} = A(\mathbb{D})$, and $\mathfrak{q}_{ac} \leq \mathfrak{q}_{\mu}$ is the largest closeable and positive semi-definite quadratic form, $A(\mathbb{D})$ is a form-core for \mathfrak{q}_D , and hence a core for $\sqrt{\mathfrak{q}_D}$. Hence, $A(\mathbb{D})$ is dense in $\hat{\mathcal{H}}(\mathfrak{q}_D) = \hat{\mathcal{H}}(\mathfrak{q}_{ac})$. Given any $a \in A(\mathbb{D})$, let $x_n \in \mathcal{D}$ be a sequence which converges to a in supremum-norm. Then

$$\begin{aligned} 0 &\leq \|x_n - a\|_{H^2(\lambda)}^2 + \mathfrak{q}_{ac}(x_n - a, x_n - a) \\ &\leq \|x_n - a\|_{H^2(\lambda)}^2 + \|x_n - a\|_{H^2(\mu)}^2 \\ &\leq \|x_n - a\|_{\infty}^2(\mu(\partial\mathbb{D}) + \lambda(\partial\mathbb{D})) \rightarrow 0. \end{aligned}$$

This proves that \mathcal{D} is dense in the dense subspace $A(\mathbb{D}) \subseteq \hat{\mathcal{H}}(\mathfrak{q}_D)$, and hence \mathcal{D} is a form-core for \mathfrak{q}_D and a core for $\sqrt{\mathfrak{q}_D}$. ■

Given any $h \in H^2(\mathfrak{q}_{ac})$ or in $H^2(\mathfrak{q}_s)$, we can now define the \mathfrak{q}_{ac} or \mathfrak{q}_s -Cauchy transform of h as before:

$$(\mathcal{E}_{ac}h)(z) := \mathfrak{q}_{ac}(k_z, h),$$

and similarly for q_s . As in Lemma 2.1 and Lemma 2.2, Cauchy transforms of elements of $H^2(q_{ac}), H^2(q_s)$ are holomorphic in the unit disk, and if we equip the vector space of q_{ac} -Cauchy transforms with the inner product

$$\langle \mathcal{C}_{ac}x, \mathcal{C}_{ac}y \rangle_{ac} := q_{ac}(x, y),$$

we obtain a reproducing kernel Hilbert space of analytic functions in the disk, $\mathcal{H}^+(q_{ac})$ with reproducing kernel:

$$k^{(ac)}(z, w) := q_{ac}(k_z, k_w).$$

Finally, since $q_\mu = q_{ac} + q_s$, $q_\mu \geq q_{ac}$, $q_s \geq 0$, we obtain the following.

Proposition 4.11 *The RKHS of q_{ac} and q_s -Cauchy transforms are contractively contained in $\mathcal{H}^+(\mu) = \mathcal{H}^+(q_\mu)$ and $k^\mu = k^{(ac)} + k^s$ so that*

$$\mathcal{H}^+(\mu) = \mathcal{H}^+(q_{ac}) + \mathcal{H}^+(q_s).$$

Moreover, if $e_{ac} : \mathcal{H}^+(q_{ac}) \hookrightarrow \mathcal{H}^+(\mu)$ and e_s are the contractive embeddings, then

$$I_{\mathcal{H}^+(\mu)} = e_{ac}e_{ac}^* + e_s e_s^*.$$

Proof To check the decomposition of the identity, it suffices to calculate

$$\begin{aligned} k^\mu(z, w) &= k^{(ac)}(z, w) + k^s(z, w) \\ &= \left\langle k_z^{(ac)}, k_w^{(ac)} \right\rangle_{ac} + \left\langle k_z^s, k_w^s \right\rangle_s \\ &= \left\langle e_{ac}^* k_z^\mu, e_{ac}^* k_w^\mu \right\rangle_{ac} + \left\langle e_s^* k_z^\mu, e_s^* k_w^\mu \right\rangle_s \\ &= \left\langle k_z^\mu, (e_{ac}e_{ac}^* + e_s e_s^*) k_w^\mu \right\rangle_\mu. \end{aligned}$$

■

Theorem 4.12 *Let $q_\mu = q_{ac} + q_s$ be the Simon–Lebesgue decomposition of the form q_μ with dense form domain $A(\mathbb{D})$ in $H^2(\lambda)$. Then,*

$$\mathcal{H}^+(q_{ac}) = \text{int}(\mu, \lambda)^{-\mu} = (\mathcal{H}^+(\mu) \cap \mathcal{H}^+(\lambda))^{-\|\cdot\|_\mu}.$$

If $e : \text{int}(\mu, \lambda) \subseteq \mathcal{H}^+(\mu) \hookrightarrow \mathcal{H}^+(\lambda)$ is the closed embedding and $q_D = \overline{q_{ac}}$, then

$$D = \mathcal{C}_\lambda^* e e^* \mathcal{C}_\lambda.$$

Lemma 4.13 *Let q_1, q_2 be densely-defined, closed and positive semi-definite quadratic forms in a separable, complex Hilbert space, \mathcal{H} . Then $q_1 \leq q_2$ if and only if $q_1(x, x) \leq q_2(x, x)$ for all x in a form-core for q_2 .*

Proof (of Theorem 4.12.) First, since q_{ac} is closeable, $\overline{q_{ac}} = q_B$ for some closed, self-adjoint operator $B \geq 0$. By construction, $A(\mathbb{D}) \subseteq \text{Dom } \sqrt{B}$, and $\mathbb{C}[\zeta], \mathcal{K}_{\mathbb{D}} = \vee k_z$ and $A(\mathbb{D})$ are all cores for \sqrt{B} . Since $B \geq 0$ is closed, $\text{Dom } B$ is also a core for \sqrt{B} . It follows that we can identify $\text{Dom } B$ with a dense subspace of $H^2(q_{ac})$. Namely, if $x \in \text{Dom } B \subseteq H^2(\lambda)$, we can find $a_n \in A(\mathbb{D})$ so that $a_n \rightarrow x$ in $H^2(\lambda)$ and $\sqrt{B}a_n \rightarrow \sqrt{B}x$. Since $\overline{q_{ac}} = q_B$, it follows that (a_n) is a Cauchy sequence in $H^2(q_{ac})$, and we can identify $x \in \text{Dom } B$ with the limit, \hat{x} , of this Cauchy sequence in the Hilbert space $H^2(q_{ac})$. Finally, since $\text{Dom } B$ is a core for \sqrt{B} , for any $a \in A(\mathbb{D}) \subseteq \text{Dom } \sqrt{B}$, we can find $x_n \in \text{Dom } B$ so that $x_n \rightarrow a$ and $\sqrt{B}x_n \rightarrow \sqrt{B}a$ and it follows that $\hat{x}_n \rightarrow a$ in $H^2(q_{ac})$, so that $\text{Dom } B$ can be identified with a dense subspace of $H^2(q_{ac})$.

Furthermore, we can then define the q_{ac} -Cauchy transform of any $x \in \text{Dom } B$,

$$\begin{aligned} (\mathcal{C}_{ac}x)(z) &= \lim_{n \uparrow \infty} q_{ac}(k_z, a_n) = \lim \left\langle \sqrt{B}k_z, \sqrt{B}a_n \right\rangle_{H^2(\lambda)} \\ &= \left\langle \sqrt{B}k_z, \sqrt{B}x \right\rangle_{H^2(\lambda)} \\ &= \langle k_z, Bx \rangle_{H^2(\lambda)} = (\mathcal{C}_\lambda Bx)(z). \end{aligned}$$

This proves that $\mathcal{C}_{ac}x \in \mathcal{H}^+(\lambda)$. Since $\mathcal{C}_{ac}x \in \mathcal{H}^+(q_{ac}) \subseteq \mathcal{H}^+(\mu)$, it follows that $\mathcal{C}_{ac}\text{Dom } B \subseteq \text{int}(\mu, \lambda)$. Moreover, since $\text{Dom } B$ can be identified with a dense subspace of $H^2(q_{ac})$, it follows that $\mathcal{C}_{ac}\text{Dom } B \subseteq \mathcal{H}^+(q_{ac}) \cap \mathcal{H}^+(\lambda) \subseteq \text{int}(\mu, \lambda)$ is dense in $\mathcal{H}^+(q_{ac})$.

Now consider q_D , where $D = \mathcal{C}_\lambda^* e e^* \mathcal{C}_\lambda$ and $e : \text{int}(\mu, \lambda) \subseteq \mathcal{H}^+(\mu) \hookrightarrow \mathcal{H}^+(\lambda)$, as in the theorem statement. By construction, $\mathcal{K}_{\mathbb{D}}$ is a core for \sqrt{D} , and it is also a core for B , so that this set is a form-core for both $q_B = \overline{q_{ac}}$ and q_D . It follows that $q_D|_{\mathcal{K}_{\mathbb{D}}} \leq q_\mu|_{\mathcal{K}_{\mathbb{D}}}$ is a positive closeable form so that by maximality and Lemma 4.13, $q_D \leq q_B$ in the form-sense.

Also, by construction, $e^*k_z^\lambda = k_z^{\mu \cap \lambda}$, where $k^{\mu \cap \lambda}$ is the reproducing kernel for the closed subspace $\text{int}(\mu, \lambda)^{-\mu} \subseteq \mathcal{H}^+(\mu)$. Hence, for any finite subset, $\{z_1, \dots, z_n\} \subseteq \mathbb{D}$, if we consider any finite linear combination of Szegő kernels,

$$h = \sum_{i=1}^n c_i k_{z_i},$$

then

$$\begin{aligned} 0 &\leq \sum \bar{c}_i c_j k^{\mu \cap \lambda}(z_i, z_j) \\ &= \sum \bar{c}_i c_j \left\langle k_{z_i}^{\mu \cap \lambda}, k_{z_j}^{\mu \cap \lambda} \right\rangle_\mu \\ &= \mathfrak{q}_D(h, h) \leq \mathfrak{q}_B(h, h) \\ &= \sum \bar{c}_i c_j \mathfrak{q}_B(k_{z_i}, k_{z_j}) \\ &= \sum \bar{c}_i c_j k^{ac}(z_i, z_j). \end{aligned}$$

That is,

$$0 \leq [k^{\mu \cap \lambda}(z_i, z_j)]_{1 \leq i, j \leq n} = [\mathfrak{q}_D(k_{z_i}, k_{z_j})] \leq [\mathfrak{q}_B(k_{z_i}, k_{z_j})] = [k^{ac}(z_i, z_j)],$$

so that $k^{\mu \cap \lambda} \leq k^{ac}$, and by Aronszajn's inclusion theorem, $\text{int}(\mu, \lambda)^{-\mu}$ is contractively contained in $\mathcal{H}^+(\mathfrak{q}_{ac})$ which is in turn contractively contained in $\mathcal{H}^+(\mu)$. Hence, if e_1 is the first embedding into $\mathcal{H}^+(\mathfrak{q}_{ac})$ and e_2 is the second embedding into $\mathcal{H}^+(\mu)$, the composite embedding, $e = e_2 e_1 : \text{int}(\mu, \lambda)^{-\mu} \hookrightarrow \mathcal{H}^+(\mu)$ is again a contractive embedding and it must be isometric since $\text{int}(\mu, \lambda)^{-\mu}$ is a closed subspace of $\mathcal{H}^+(\mu)$. It follows that e_1 must be an isometric embedding. Indeed, if there is a unit vector x so that $\|e_1 x\| < 1$ then

$$1 = \|x\| = \|ex\| \leq \|e_2\| \|e_1 x\| < 1.$$

Similarly e_2 must be isometric on the range of e_1 . On the other hand, since $\text{int}(\mathfrak{q}_{ac}, \lambda) := \mathcal{H}^+(\mathfrak{q}_{ac}) \cap \mathcal{H}^+(\lambda)$ is dense in $\mathcal{H}^+(\mathfrak{q}_{ac})$ and $\mathcal{H}^+(\mathfrak{q}_{ac})$ is contractively contained in $\mathcal{H}^+(\mu)$, we must have that $\text{int}(\mathfrak{q}_{ac}, \lambda) \subseteq \text{int}(\mu, \lambda) \subseteq \text{Ran } e_1$. Hence, by the previous argument, since $\text{int}(\mathfrak{q}_{ac}, \lambda) \subseteq \text{Ran } e_1$ is dense in $\mathcal{H}^+(\mathfrak{q}_{ac})$ and e_2 is isometric on the range of e_1 , $e_2 : \mathcal{H}^+(\mathfrak{q}_{ac}) \hookrightarrow \mathcal{H}^+(\mu)$ is also an isometric inclusion. In conclusion, $\text{int}(\mu, \lambda)^{-\mu}$ and $\mathcal{H}^+(\mathfrak{q}_{ac})$ are both closed subspaces of $\mathcal{H}^+(\mu)$, $\text{int}(\mu, \lambda)^{-\mu}$ is a closed subspace of $\mathcal{H}^+(\mathfrak{q}_{ac})$ and $\text{int}(\mathfrak{q}_{ac}, \lambda) \subseteq \text{int}(\mu, \lambda)$ is dense in $\mathcal{H}^+(\mathfrak{q}_{ac})$ so that $\text{int}(\mu, \lambda)^{-\mu} = \mathcal{H}^+(\mathfrak{q}_{ac})$. It follows that $\mathfrak{q}_B = \mathfrak{q}_D$ on $\mathcal{H}_\mathbb{D}$ so that by Lemma 4.13 and the uniqueness of representation of closed forms, $D = B$. ■

Corollary 4.14 *If $\mu, \lambda \geq 0$ are finite, positive and regular Borel measures on $\partial\mathbb{D}$ and \mathfrak{q}_μ is the densely-defined positive quadratic form associated to μ with form domain $A(\mathbb{D}) \subseteq H^2(\lambda)$, then the space of μ -Cauchy transforms decomposes as the orthogonal direct sum,*

$$\mathcal{H}^+(\mu) = \mathcal{H}^+(\mathfrak{q}_{ac}) \oplus \mathcal{H}^+(\mathfrak{q}_s).$$

In particular, $\mathcal{H}^+(\mathfrak{q}_s) \cap \text{int}(\mu, \lambda) = \{0\}$.

Proof By Proposition 4.11 and Theorem 3.8, we have that the identity operator on $\mathcal{H}^+(\mu)$ decomposes as

$$I_\mu = e_{ac} e_{ac}^* + e_s e_s^*,$$

and $\mathcal{H}^+(\mathfrak{q}_{ac}) = \text{int}(\mu, \lambda)^{-\mu}$ is a closed subspace of $\mathcal{H}^+(\mu)$ so that the contractive embedding, $e_{ac} : \mathcal{H}^+(\mathfrak{q}_{ac}) \hookrightarrow \mathcal{H}^+(\mu)$ is an isometry. Hence, $P_{ac} := e_{ac} e_{ac}^*$ is an orthogonal projection onto the range of e_{ac} and hence $P_s = I - P_{ac} = e_s e_s^*$ is the projection onto the orthogonal complement of $\text{Ran } e_{ac}$ in $\mathcal{H}^+(\mu)$. It follows that e_s is also an isometric embedding and that we can identify $\mathcal{H}^+(\mathfrak{q}_{ac}), \mathcal{H}^+(\mathfrak{q}_s)$ with the ranges of these isometric embeddings so that

$$\mathcal{H}^+(\mu) = \mathcal{H}^+(\mathfrak{q}_{ac}) \oplus \mathcal{H}^+(\mathfrak{q}_s). \quad \blacksquare$$

Corollary 4.15 *Let μ, λ be positive, finite and regular Borel measures on the unit circle. The Lebesgue decomposition of μ with respect to λ , $\mu = \mu_{ac} + \mu_s$, coincides with the Simon–Lebesgue decomposition of \mathfrak{q}_μ with form domain $\text{Dom } \mathfrak{q}_\mu = A(\mathbb{D})$ in $H^2(\lambda)$, $\mathfrak{q}_\mu = \mathfrak{q}_{ac} + \mathfrak{q}_s$, in the sense that $\mathfrak{q}_{ac} = \mathfrak{q}_{\mu_{ac}}$ and $\mathfrak{q}_s = \mathfrak{q}_{\mu_s}$ if and only if $\text{int}(\mu, \lambda)$ is V^μ -reducing.*

Remark 4.16 More generally, one can apply the methods of this section to construct a Lebesgue decomposition for pairs of positive kernel functions k, K on the same set, X , see Appendix A.

Example 4.17 (Lebesgue measure on the half-circles.) As before, let m_{\pm} denote normalized Lebesgue measure restricted to the upper and lower half-circles. These are mutually singular measures so that $m_+ = m_{+,s}$ is the singular part of m_+ with respect to m_- , and yet by Example 4.9, $\text{int}(m_+, m_-) \neq \{0\}$, so that $q_+ = q_{m_+}$ has a Simon–Lebesgue decomposition $q_+ = q_{ac} + q_s$ in $H^2(m_-)$, where q_{ac} is non-trivial, by Theorem 4.12. Moreover, in this example, m_- is extreme, so that $H^2(m_-) = L^2(m_-)$. This means that while the quadratic form, q_{μ} , associated to μ , with dense form domain, $A(\mathbb{D}) \subseteq L^2(m_-) = H^2(m_-)$ has non-zero absolutely continuous part, if we instead define the form domain of q_{μ} to be $\text{Dom } q_{\mu} = \mathcal{C}(\partial\mathbb{D})$, then, with this form domain, q_{μ} has vanishing absolutely continuous part (since the decompositions of q_{μ} and μ always coincide in this case, see Remark 3.9). This shows, that in dealing with these unbounded positive quadratic Toeplitz forms, the choice of form domain is crucial!

4.2 Lebesgue decomposition for arbitrary measures

The question remains: If $\mu, \lambda \geq 0$ are arbitrary, how can we construct the Lebesgue decomposition of μ with respect of λ using reproducing kernel theory and their spaces of Cauchy transforms? If λ is non-extreme, or more generally if $\text{int}(\mu, \lambda)$ is V_{μ} -reducing, Theorem 4.5 provides a satisfying answer. However, as Proposition 4.7, Example 4.9 and Theorem 4.12 show, the intersection of the spaces of μ and λ Cauchy transforms cannot be reducing in general, and that there are examples of pairs of positive measures μ, λ , for which $\text{int}(\mu, \lambda)$ cannot be equal to, or even contain, the space of Cauchy transforms of any non-zero positive measure.

By Theorem 3.12, we do know that if $\mu = \mu_{ac} + \mu_s$ is the Lebesgue decomposition of μ with respect to λ , that $\mu_{ac} \ll_{RK} \lambda$ so that $\text{int}(\mu_{ac}, \lambda) \subseteq \text{int}(\mu, \lambda) \subseteq \text{int}(\mu, \lambda)^{-\mu} = \mathcal{H}^+(\mu_{ac})$. The final result below provides an abstract characterization of the Lebesgue decomposition for arbitrary pairs of positive measures.

Theorem 4.18 *If $\mu = \mu_{ac} + \mu_s$ is the Lebesgue decomposition of μ with respect to λ and $q_{\mu} = q_{ac} + q_s$ is the Simon–Lebesgue form decomposition of q_{μ} in $H^2(\lambda)$ then $q_{\mu_{ac}} \leq q_{ac}$. Moreover, $\mathcal{H}^+(\mu_{ac})$ is the maximal RKHS, $\mathcal{H}(k)$, in \mathbb{D} with the following property: $\mathcal{H}(k) \cap \mathcal{H}^+(\lambda) \subseteq \text{int}(\mu, \lambda)$ is dense in $\mathcal{H}(k)$, $\mathcal{H}(k) \subseteq \mathcal{H}^+(\mu)$ is contractively contained, and if $e : \mathcal{H}(k) \hookrightarrow \mathcal{H}^+(\mu)$ is the contractive embedding, then ee^* is V_{μ} -Toeplitz. Equivalently, $q_{\mu_{ac}}$ is the largest closeable Z^{λ} -Toeplitz form bounded above by q_{μ} .*

Moreover, if $e_1 := e_{\mu_{ac}}$ and $e_2 = e_{\mu_s}$ then $I_{\mu} = e_1 e_1^* + e_2 e_2^*$. Hence, we can identify $\mathcal{H}^+(\mu_{ac})$ with the operator-range space $\mathcal{R}(e_1)$ and $\mathcal{H}^+(\mu_s)$ with $\mathcal{R}(e_2) = \mathcal{R}^c(e_1)$, the complementary space of $\mathcal{H}^+(\mu_{ac})$ in the sense of deBranges and Rovnyak and

$$\mathcal{H}^+(\mu) = \mathcal{H}^+(\mu_{ac}) + \mathcal{H}^+(\mu_s).$$

Proof This follows from the definition of q_{ac} , Theorem 1.4 and Theorem 4.1. ■

Remark 4.19 In the case where the complementary space decomposition of $\mathcal{H}^+(\mu) = \mathcal{H}^+(\mu_{ac}) + \mathcal{H}^+(\mu_s)$, appearing in the above theorem statement, is not an orthogonal direct sum, this yields a corresponding decomposition of the quadratic form q_{μ} ,

$$q_{\mu} = q_{\mu_{ac}} + q_{\mu_s}, \tag{4.3}$$

where $q_{\mu_{ac}} < q_{ac}$ and $q_{\mu} = q_{ac} + q_s$ is the Simon–Lebesgue decomposition of q_{μ} . In this case, the decomposition of Equation (4.3) is an example of a ‘pseudo-orthogonal’ Lebesgue decomposition of q_{μ} as recently defined and studied in [11].

The previous theorem is, while interesting, admittedly not very practical for construction of the Lebesgue decomposition of μ with respect to λ . A simpler, albeit somewhat ad hoc, approach using our reproducing kernel methods is simply to ‘add Lebesgue measure’. Namely, if $\mu_{ac;\lambda}$ is the absolutely continuous part of μ with respect to λ , then $\mu_{ac;\lambda} = \mu_{ac;\lambda+m} - \mu_{ac;m}$ and both $\lambda + m$ and m are non-extreme so that Theorem 4.5 applies.

A Lebesgue decomposition of positive kernels

Let K be a fixed positive kernel function on a set, X . Given any other positive kernel, k , on X , we can associate to it the densely-defined and positive semi-definite quadratic form, $q_k : \text{Dom } q_k \times \text{Dom } q_k \rightarrow \mathbb{C}$, with dense form domain $\text{Dom } q_k := \bigvee_{x \in X} K_x$ in $\mathcal{H}(K)$,

$$q_k(K_x, K_y) := k(x, y).$$

One can then apply B. Simon’s Lebesgue decomposition of positive quadratic forms to q_k . Such a Lebesgue decomposition of positive kernels was first considered in [12, Section 7, Theorem 7.2]. The theorem below provides some more details about this decomposition.

Theorem A.1 Let k, K be positive kernel functions on a set, X . If \mathfrak{q}_k is the densely-defined positive quadratic form of k in $\mathcal{H}(K)$, as defined above, with Simon–Lebesgue form decomposition $\mathfrak{q}_k = \mathfrak{q}_{ac} + \mathfrak{q}_s$, then there are positive kernels, k^{ac} and k^s on X , so that $\mathfrak{q}_{ac} = \mathfrak{q}_{k^{ac}}$, $\mathfrak{q}_s = \mathfrak{q}_{k^s}$, $k = k^{ac} + k^s$, and

$$\mathcal{H}(k) = \mathcal{H}(k^{ac}) \oplus \mathcal{H}(k^s).$$

Moreover, $\mathcal{H}(k^{ac}) = \text{int}(k, K)^{-k} := (\mathcal{H}(k) \cap \mathcal{H}(K))^{-\|\cdot\|_k}$, and if $e : \text{int}(k, K) \hookrightarrow \mathcal{H}(K)$ is the (closed) embedding, then $\overline{\mathfrak{q}_{ac}} = \mathfrak{q}_{ee^*}$.

In the above, $\text{int}(k, K) := \mathcal{H}(k) \cap \mathcal{H}(K)$.

Proof Let $h := \sum_{i=1}^n c_i K_{x_i}$ be any finite linear combination of the kernels K_{x_i} , $\{x_i\}_{i=1}^n \subseteq X$. Then, since $\mathfrak{q}_{ac} \leq \mathfrak{q}_k$, we obtain that

$$\begin{aligned} \sum \overline{c_i c_j} k(x_i, x_j) &= \sum \overline{c_i c_j} \mathfrak{q}_k(x_i, x_j) \\ &= \mathfrak{q}_k(h, h) \geq \mathfrak{q}_{ac}(h, h) \geq 0, \end{aligned}$$

where

$$0 \leq \mathfrak{q}_{ac}(h, h) = \sum \overline{c_i c_j} \mathfrak{q}_{ac}(K_{x_i}, K_{x_j}).$$

It follows that

$$k^{ac}(x, y) := \mathfrak{q}_{ac}(K_x, K_y),$$

defines a positive kernel function on X so that $0 \leq k^{ac} \leq k$. Similarly, $k^s(x, y) := \mathfrak{q}_s(K_x, K_y)$ defines a positive kernel function on X so that $0 \leq k^s \leq k$, and since $\mathfrak{q}_k = \mathfrak{q}_{ac} + \mathfrak{q}_s$, we obtain that $k^{ac} + k^s = k$.

By definition, \mathfrak{q}_{ac} is the largest closeable quadratic form bounded above by \mathfrak{q}_k . In particular $\overline{\mathfrak{q}_{ac}} = \mathfrak{q}_D$ is the positive form of some densely-defined, self-adjoint and positive semi-definite operator D , so that $\mathcal{K}_X := \bigvee_{x \in X} K_x$ is a core for \sqrt{D} . (Here, \bigvee denotes non-closed linear span.) If $e : \text{int}(k, K) \subseteq \text{int}(k, K)^{-k} \hookrightarrow \mathcal{H}(K)$ is the densely-defined and closed embedding, let $A := ee^*$. We claim that $A = D$. First, $A \geq 0$ is self-adjoint, hence closed, and since e is trivially a multiplier, we obtain that

$$\mathfrak{q}_A(K_x, K_y) = \langle e^* K_x, e^* K_y \rangle_k = \langle k_x^\cap, k_y^\cap \rangle_k = k^\cap(x, y),$$

where k^\cap denotes the reproducing kernel of the subspace $\text{int}(k, K)^{-k} \subseteq \mathcal{H}(k)$, the closure of the intersection space, $\text{int}(k, K)$ in $\mathcal{H}(k)$. In particular, since $k_x^\cap = P_\cap k_x$, where $P_\cap : \mathcal{H}(k) \rightarrow \text{int}(k, K)^{-k}$ is the orthogonal projection, it follows that $k^\cap \leq k$, and hence that $\mathfrak{q}_A \leq \mathfrak{q}_k$. Since $\mathfrak{q}_A|_{\mathcal{K}_X}$ is closeable, it follows, by maximality of the Simon–Lebesgue decomposition, that $\mathfrak{q}_A \leq \mathfrak{q}_D$. This inequality implies that $k^\cap \leq k^{ac}$ as positive kernels on X .

Now suppose that $h \in \text{Dom } D \subseteq \mathcal{H}(K)$ and choose $h_n \in \mathcal{K}_X = \bigvee_{x \in X} K_x$ so that $h_n \rightarrow h$ and $\sqrt{D}h_n \rightarrow \sqrt{D}h$. (This can be done since $\text{Dom } D$ is a core for \sqrt{D} .) If $h_n = \sum_{j=1}^{m_n} c_j(n) K_{x_j(n)}$, a finite linear combination, then note that

$$\begin{aligned} (Dh)(x) &= \lim_{n \uparrow \infty} \sum c_j(n) \langle \sqrt{D}K_x, \sqrt{D}K_{x_j(n)} \rangle_K \\ &= \lim \sum c_j(n) k^{ac}(x, x_j(n)) = \lim g_n(x), \end{aligned}$$

where

$$g_n = \sum c_j(n) k_{x_j(n)}^{ac} \in \mathcal{H}(k^{ac}) \subseteq \mathcal{H}(k).$$

Moreover,

$$\begin{aligned} \|g_n\|_{k^{ac}}^2 &= \sum_{i,j} \overline{c_i(n) c_j(n)} k^{ac}(x_i(n), x_j(n)) \\ &= \langle \sqrt{D}h_n, \sqrt{D}h_n \rangle_K \rightarrow \|\sqrt{D}h\|_K^2, \end{aligned}$$

so that the sequence $(g_n) \subseteq \mathcal{H}(k^{ac})$ is uniformly bounded in norm. Since $g_n(x) \rightarrow (Dh)(x)$ pointwise in X , this and uniform boundedness imply that g_n converges weakly to the function Dh . Since Hilbert spaces are weakly closed, the function $Dh \in \mathcal{H}(k^{ac}) \subseteq \mathcal{H}(k)$, and also $Dh \in \mathcal{H}(K)$ so that $Dh \in \text{int}(k, K) \subseteq \mathcal{H}(k^\cap)$. Hence $Dh \in \mathcal{H}(k^{ac})$ and

$$\|Dh\|_{k^{ac}}^2 = \|\sqrt{D}h\|_K^2.$$

Let $j_1 : \mathcal{H}(k^\cap) \hookrightarrow \mathcal{H}(k^{ac})$ and $j_2 : \mathcal{H}(k^{ac}) \hookrightarrow \mathcal{H}(k)$ be the contractive embeddings. Then $j := j_2 j_1 : \mathcal{H}(k^\cap) \hookrightarrow \mathcal{H}(k)$ is the isometric embedding of the subspace $\mathcal{H}(k^\cap) \subseteq \mathcal{H}(k)$ into $\mathcal{H}(k)$. It follows that j_1 must be isometric and j_2 must be isometric on the range of j_1 in $\mathcal{H}(k^{ac})$.

We claim that $\text{Ran } j_1$ is dense in $\mathcal{H}(k^{ac})$ so that j_1 and j_2 are both isometries. Define a linear map, $V : \mathcal{K}_X^{ac} := \bigvee_{x \in X} k_x^{ac} \rightarrow \mathcal{H}(K)$ by

$$V k_x^{ac} := \sqrt{D} K_x,$$

and extending linearly. Since

$$\langle k_x^{ac}, k_y^{ac} \rangle_{k^{ac}} = k^{ac}(x, y) = \langle \sqrt{D} K_x, \sqrt{D} K_y \rangle_K,$$

it follows that V is an isometry and extends by continuity to an isometry from $\mathcal{H}(k^{ac})$ onto the closure of $\sqrt{D} \mathcal{K}_X$ in $\mathcal{H}(K)$, which we also denote by V . Since \mathcal{K}_X is a core for \sqrt{D} , V is onto $\text{Ran } \sqrt{D}^{-\|\cdot\|_K}$ in $\mathcal{H}(K)$. If there exists a $g \in \mathcal{H}(k^{ac})$ orthogonal to $\text{Ran } D \subseteq \mathcal{H}(k^{ac})$, then choose a sequence $g_n = \sum_{j=1}^{m_n} c_j(n) k_{x_j(n)}^{ac} \in \mathcal{K}_X^{ac}$ so that $g_n \rightarrow g$ and calculate, for any $h \in \text{Dom } D$, that

$$\begin{aligned} 0 &= \langle g, Dh \rangle_{k^{ac}} = \lim_n \langle g_n, Dh \rangle_{k^{ac}} \\ &= \lim_n \sum_j \overline{c_j(n)} \langle k_{x_j(n)}^{ac}, Dh \rangle_{k^{ac}} \\ &= \lim_n \sum_j \overline{c_j(n)} (Dh)(x_j(n)) \\ &= \lim_n \sum_j \overline{c_j(n)} \langle K_{x_j(n)}, Dh \rangle_K \\ &= \lim_n \left\langle \sum_j c_j(n) \sqrt{D} K_{x_j(n)}, \sqrt{D} h \right\rangle_K \\ &= \lim_n \langle V g_n, \sqrt{D} h \rangle_K \\ &= \langle V g, \sqrt{D} h \rangle_K. \end{aligned}$$

This proves that $Vg \in \text{Ran } \sqrt{D}^{-\|\cdot\|_K}$ is orthogonal to $\sqrt{D} \text{Dom } D$. However, $\text{Dom } D$ is a core for \sqrt{D} , so that $\sqrt{D} \text{Dom } D$ is dense in $\text{Ran } \sqrt{D}$. This proves that $Vg = 0$, and hence $g = 0$. In conclusion, $\text{Ran } D \subseteq \text{Ran } j_1 \subseteq \mathcal{H}(k^{ac})$ is dense in $\mathcal{H}(k^{ac})$ so that both j_1 and j_2 are isometric embeddings. That is, $\mathcal{H}(k^\cap)$ embeds, as a closed, dense subspace of $\mathcal{H}(k^{ac})$, which embeds isometrically into $\mathcal{H}(k)$ and we conclude that $\mathcal{H}(k^\cap) = \mathcal{H}(k^{ac})$ so that $k^\cap = k^{ac}$ and $\mathfrak{q}_D = \mathfrak{q}_A$. By the uniqueness of Kato's Riesz representation of closed, positive semi-definite forms, $D = A$ as closed operators.

The fact that $k = k^{ac} + k^s$, implies that if $e_{ac} : \mathcal{H}(k^{ac}) \hookrightarrow \mathcal{H}(k)$ is the isometric embedding and $e_s : \mathcal{H}(k^s) \hookrightarrow \mathcal{H}(k)$ is the contractive (and injective) embedding, that $I = e_{ac} e_{ac}^* + e_s e_s^*$. Hence $e_s e_s^* = I - P_{ac}$, so that $\mathcal{H}(k^s)$ also embeds isometrically in $\mathcal{H}(k)$ as the orthogonal complement of $\mathcal{H}(k^{ac})$. ■

References

- [1] A. B. Aleksandrov. Multiplicity of boundary values of inner functions. (Russian). *Izvestiya Akademii Nauk Armyanskoi SSR*, 22:490–503, 1987.
- [2] A. B. Aleksandrov. On the existence of nontangential boundary values of pseudocontinuable functions. (Russian). *Zapiski Nauchnykh Seminarov POMI*, 222:5–17, 1995.
- [3] N. Aronszajn. Theory of reproducing kernels. *Transactions of the American Mathematical Society*, 68:337–404, 1950.
- [4] J. A. Ball and V. Vinnikov. Formal reproducing kernel Hilbert spaces: the commutative and noncommutative settings. In *Reproducing kernel spaces and applications*, pages 77–134. Springer, 2003.
- [5] D. N. Clark. One dimensional perturbations of restricted shifts. *Journal d'Analyse Mathématique*, 25:169–191, 1972.
- [6] L. de Branges and J. Rovnyak. *Square summable power series*. Holt, Rinehart and Winston, 1966.
- [7] P. Fatou. Séries trigonométriques et séries de Taylor. *Acta Mathematica*, 37:335–400, 1906.
- [8] E. Fricain and J. Mashreghi. *The theory of $\mathcal{H}(b)$ spaces*, volume 2. Cambridge University Press, 2016.
- [9] U. Grenander and G. Szegö. *Toeplitz forms and their applications*. University of California Press, 1958.
- [10] P. R. Halmos and A. Brown. Algebraic properties of Toeplitz operators. *Journal für die reine und angewandte Mathematik*, 213:89–102, 1963.
- [11] S. Hassi and H. de Snoo. Complementation and Lebesgue type decompositions of linear operators and relations. arXiv:2308.09408, 2023.
- [12] S. Hassi, Z. Sebestyén, and H. de Snoo. Lebesgue type decompositions for nonnegative forms. *Journal of Functional Analysis*, 257:3858–3894, 2009.

- [13] K. Hoffman. *Banach spaces of analytic functions*. Prentice-Hall, 1962.
- [14] M. T. Jury and R. T. W. Martin. Fatou's theorem for non-commutative measures. *Advances in Mathematics*, 400:108293, 2022.
- [15] M. T. Jury and R. T. W. Martin. Lebesgue decomposition of non-commutative measures. *International Mathematics Research Notices*, 2022:2968–3030, 2022.
- [16] T. Kato. *Perturbation theory for linear operators*. Springer, 1980.
- [17] R. T. W. Martin and E. Shamovich. A de Branges–Beurling theorem for the full Fock space. *Journal of Mathematical Analysis and Applications*, 496:124765, 2021.
- [18] B. Okutmustur and A. Gheondea. *Reproducing Kernel Hilbert spaces. The Basics, Bergman spaces, and interpolation problems*. Lambert, 2010.
- [19] V. I. Paulsen. *Completely Bounded Maps and Operator Algebras*. Cambridge University Press, New York, NY, 2002.
- [20] V. I. Paulsen and M. Raghupathi. *An introduction to the theory of reproducing kernel Hilbert spaces*, volume 152. Cambridge university press, 2016.
- [21] M. Reed and B. Simon. *Functional Analysis*. Academic Press, 1980.
- [22] B. Simon. A canonical decomposition for quadratic forms with applications to monotone convergence theorems. *Journal of Functional Analysis*, 28:377–385, 1978.
- [23] G. Szegö. Beiträge zur theorie der Toeplitzschens formen. *Mathematische Zeitschrift*, 6:167–202, 1920.
- [24] J. von Neumann. On rings of operators III. *Annals of Mathematics*, 41:94–161, 1940.

Jashan Bal, Department of Pure Mathematics, University of Waterloo
E-mail address: j2bal@uwaterloo.ca

Robert T. W. Martin, Department of Mathematics, University of Manitoba
E-mail address: Robert.Martin@umanitoba.ca

Fouad Naderi, Department of Mathematics, University of Manitoba
E-mail address: naderif@myumanitoba.ca