

Weight concerns among adolescent boys

Colette Kelly^{1,*}, Amanda Fitzgerald², Mariane Sentenac^{3,4}, Jakub Gakewski¹,
Michal Molcho¹ and Saoirse Nic Gabhainn¹

¹Health Promotion Research Centre, School of Health Sciences, National University of Ireland, Galway, Republic of Ireland; ²School of Psychology, University College Dublin, Republic of Ireland; ³Inserm, Toulouse, France; ⁴Université de Toulouse III, Toulouse, France

Submitted 15 August 2014; Final revision received 4 March 2015; Accepted 16 March 2015; First published online 19 June 2015

Abstract

Objective: To investigate weight concerns among adolescent boys and relationships with health indicators and family factors.

Design: Analysis of the 2010 Health Behaviour in School-aged Children survey of 10–17-year-olds.

Setting: Schools in the Republic of Ireland.

Results: Among 6187 boys, 25.1% reported a desire to lose weight (weight ‘loss’ concern) and 7.7% reported a desire to gain weight (weight ‘gain’ concern). Both types of weight concerns were associated with poor self-rated health, life satisfaction and happiness, and with more frequent emotional and physical symptoms. Family factors were associated with boys’ weight concerns. In adjusted analyses, the risk of weight ‘loss’ concerns decreased with daily family breakfasts (OR=0.80; 95% CI 0.66, 0.97). The risk of weight ‘gain’ concerns decreased with frequent family evening meals (OR=0.77; 95% CI 0.60, 0.99). Ease of communication with mother was associated with a decreased risk of weight ‘loss’ and weight ‘gain’ concerns among boys (OR=0.74; 95% CI 0.60, 0.90 and OR=0.61; 95% CI 0.44, 0.82, respectively). An open father–son relationship and having a father present in the home decreased the risk of weight ‘loss’ concerns (OR=0.69; 95% CI 0.57, 0.82 and OR=0.81; 95% CI 0.67, 0.98, respectively).

Conclusions: Body weight concerns were reported by a sizeable minority of boys and were associated with negative health outcomes. The findings support the need to promote frequent family meals and facilitate open communication in families.

Keywords
Weight concerns
Boys
Family
Dieting
Health

Body weight and body image concerns are common among adolescents^(1–3) and various strategies can be used to achieve an ‘ideal’ body weight or shape. Unhealthy dieting practices among adolescents are associated with numerous physical, psychological and emotional negative outcomes, including poor nutritional status, depressive symptoms, low self-esteem and risk of eating disorders^(4–8). High levels of body dissatisfaction and body image disturbances are also predictive of depressive mood, psychosomatic complaints and disordered or inappropriate eating behaviours^(9–12).

Research on body weight concerns has focused on girls rather than boys and has concentrated on weight-control behaviours such as dieting and mechanisms to lose weight. Dieting is highly prevalent among adolescent girls^(4,13–15) and is described as a normative behaviour for girls⁽¹⁶⁾. Boys do not report or openly discuss weight and body image concerns^(17–22) and dieting is not described as normative for boys. As such, weight concerns and dieting

among boys may result in different or increased risks to physical and emotional health compared with those experienced among girls.

The prevalence of reported weight concerns and related behaviours among boys varies by study, ranging from 9% to 34%^(3,4,8,13,15,23). Differences in methodology and whether weight gain, weight loss or body dissatisfaction is the primary area of interest may explain such variations. Nevertheless, there exists a clear gender difference in the type of weight concerns reported. Compared with girls, boys are more likely to desire an increase in weight gain related to an increase in muscle mass⁽²⁴⁾ and, indeed, muscle-enhancing behaviours are higher now than in the past among boys^(25,26).

Family mealtimes are considered fundamental to the health of adolescents. Family meal frequency is positively associated with dietary outcomes⁽²⁷⁾ and, in general, an inverse relationship between frequency of family meals and (unhealthy) weight-loss behaviours among girls has

*Corresponding author: Email Colette.kelly@nuigalway.ie

been found^(28–30). However the relationship is not clear among boys⁽³¹⁾. Moreover, the influence of family meals on types of weight concerns among boys is not evident in the literature.

While family meals provide an opportunity for parents to provide healthy foods and model healthy behaviours, frequent family meals may also reflect the broader family environment or family dynamics⁽³²⁾. Previous research suggests the importance of considering family environment in adolescent dietary behaviours^(33,34). Dimensions of family dynamics such as parent–child relations^(35,36), family cohesion⁽³⁷⁾, connectedness⁽⁶⁾ and communication^(38,39) have been explored as factors involved in the development of dieting, disordered eating and weight-control behaviours among adolescents. However, these studies have focused on dieting and weight loss rather than on weight-gain behaviours. Family structure and maternal employment, both of which are likely to affect family dynamics, have not previously been the focus of studies on dieting and weight concerns. Mothers and fathers play different roles in young people's development and differences in mother–child and father–child relationships should also be considered. Among thirty-eight countries, boys reported easier communication with their fathers than did girls⁽⁴⁰⁾. Relationship quality between fathers and sons may be important for weight concerns among boys.

From existing research, it is evident that much less is known about weight concerns and weight-control behaviours among boys than girls. The present study explores weight concerns among 10–17-year-old boys in Ireland, in the context of the family, as well as the health outcomes of reporting weight concerns. The present study aimed to: (i) explore the prevalence of weight concerns among boys in Ireland; (ii) describe physical and emotional symptoms in boys reporting weight concerns; and (iii) investigate the relationship between family factors and weight concerns among boys.

Methods

Sample

The present study utilised data from the 2010 Irish Health Behaviour in School-aged Children study, a part of the WHO collaborative study (WHO-HBSC; www.hbcs.org). Sampling was conducted to be representative of the proportion of children in each geographical region of the Republic of Ireland. Children in primary (aged 10–12 years) and post-primary (12–17 years) schools were randomly selected and individual classrooms within these schools were subsequently randomly selected for inclusion. A total of 256 schools took part in the survey and further details on the sampling methodology and recruitment strategy are available⁽⁴¹⁾. The student response rate was 85%. The geographical location and social class of respondents were compared with the 2006 Census and

Table 1 Distribution of male respondents by age group and social class; 2010 Irish Health Behaviour in School-aged Children study

| | SC1–2 (%) | SC3–4 (%) | SC5–6 (%) | n* |
|-------------|-----------|-----------|-----------|------|
| 10–11 years | 41 | 44 | 15 | 644 |
| 12–14 years | 41 | 44 | 14 | 2557 |
| 15–17 years | 46 | 40 | 14 | 1953 |

SC, social class.

*The number of boys is less than the sample used in the present paper due to missing or incomplete social class data.

found to be representative of the population distribution across regions. Slight variations were noted but were expected for social class because the census reports all persons by social class, not all of whom would be parents or guardians of children in the age groups of interest in the study⁽⁴¹⁾. Ethical approval was obtained from the National University of Ireland, Galway Research Ethics Committee.

The current study focuses on the data collected from 6295 boys. There were 108 cases with missing data on weight concerns, resulting in 6187 boys (mean age 14.2 (SD 1.81) years) in the sample for analysis. The percentages of male respondents by age group and social class are shown in Table 1.

Measurements

Weight concerns

To identify participants who had weight concerns, students were asked 'At present are you on a diet or doing something else to lose weight?', with response options 'yes', 'no, because I need to put on weight', 'no, but I should lose some weight' and 'no, my weight is fine'. Students who answered 'yes' and who answered 'no, but I should lose some weight' were classed as having a weight 'loss' concern. Those who answered 'no, because I need to put on weight' were defined as having a weight 'gain' concern. Information on height and weight was also collected to calculate BMI (kg/m²), but self-reported BMI was not included in the present analyses because of the high rate of missing data (60% among boys aged 11–17 years).

Self-rated health and life satisfaction

Self-rated health was assessed by the question 'Would you say your health is...?' and the response options were dichotomised at 'excellent' *v.* 'good', 'fair' or 'poor'. Self-reported happiness was measured by the question 'In general, how do you feel about your life at present?' and the responses were dichotomised at 'very happy' *v.* 'quite happy', 'don't feel very happy' and 'not happy at all'. Children were also asked to rank themselves from 0 to 10 on a life satisfaction ladder^(42,43). This scale was used to identify those with high life satisfaction (response >6).

Physical and emotional symptoms

Psychosomatic symptoms were measured using the HBSC symptom checklist⁽⁴⁴⁾. Children were asked to report the

frequency, in the six months prior to the survey, that they experienced a variety of emotional (feeling low, bad temper, feeling nervous, sleeping difficulties) and physical symptoms (headache, stomach-ache, backache, feeling dizzy). Response options were 'about every day', 'more than once a week', 'about every week', 'about every month' and 'rarely or never'. Children were characterised as symptomatic if they reported two or more symptoms at least once per week (physical and emotional symptoms were analysed separately).

Family factors

Students were asked 'How often do you have breakfast together with your mother or father?' and 'How often do you have an evening meal together with your mother or father?' Response options were: 'never', 'less than once a week', '1–2 days a week', '3–4 days a week', '5–6 days a week' and 'every day'. Responses were categorised as daily *v.* less than daily.

Communication with mother and father was assessed separately with two items: 'How easy is it for you to talk to your mother/father about things that really bother you?' Response options were: 'very easy', 'easy', 'difficult', 'very difficult' and 'don't have or see this person'. Ease of communication variables were dichotomised into 'very easy'/'easy' and 'difficult'/'very difficult', while the response 'don't have or see this person' was recoded to missing.

Maternal employment was determined by children answering 'yes' to the following question: 'Does your mother have a job?' Children were also asked about who they live with in their main home with the following list to choose from: 'mother', 'father', 'stepmother', 'stepfather', 'grandmother', 'grandfather', 'I live in a foster home or children's home' and 'with someone or somewhere else'. Children were dichotomised into those whose father lived in the main family home and those where the father did not.

Statistical analyses

All analyses were conducted separately for those who reported a weight 'gain' and a weight 'loss' concern. Data were analysed by age group, to reflect the developmental literature, and are in line with early, middle and older adolescence⁽⁴⁵⁾. All associations were examined with logistic regression analysis and were expressed by odds ratios. Regression analyses were adjusted for age group and for social class. Associations with weight concerns were investigated for each health indicator (self-rated health, emotional symptoms, physical symptoms, life satisfaction and happiness). Bivariate and multivariate analyses were performed to explore the associations between family factors and weight concerns. The multivariate models were adjusted for all family factors, age group and social class. Confidence intervals were computed at the 95 % level and statistical significance was

established at 5 %. All analyses were conducted using the statistical software package IBM SPSS Statistics 20.0. The reference group was children who reported that their weight was fine.

Results

Overall, 67.3 % of boys reported their weight to be fine, 25.1 % reported a weight 'loss' concern and 7.7 % reported a weight 'gain' concern (Table 2). Weight concerns were associated with age group ($P < 0.000$; Pearson $\chi^2 = 27.065$).

Health indicators and weight concerns

Table 3 shows the association between health indicators and weight 'loss' and weight 'gain' concerns as reported by participating boys. Those who expressed body weight concerns were more likely to report negative health outcomes. Weight 'loss' and weight 'gain' concerns were associated with lower levels of self-rated health, life satisfaction and happiness, as well as with more frequent emotional and physical symptoms. In addition, the associations between all health indicators studied and weight concerns were similar between boys who reported a weight 'loss' concern and those who reported a weight 'gain' concern, with the exception of self-rated health.

Family factors and weight concerns

In Table 4, crude and adjusted odds ratios between family factors and weight concerns are presented. Except for maternal employment and weight 'gain' concerns, an inverse association between all other family factors and weight concerns was evident. In unadjusted analyses all associations were statistically significant, except for the relationship between frequent evening meals and concerns about weight 'loss'.

In the full model, frequent family breakfasts and evening meals were inversely associated with concerns about weight 'loss' and weight 'gain', respectively. Ease of communication with mother was inversely associated with weight concerns. A similar pattern was evident for father-son communication and weight concerns, albeit the statistical significance was lost in the full model for concerns about weight 'gain'.

Table 2 Prevalence of weight concerns among Irish boys by age group (n 6187); 2010 Irish Health Behaviour in School-aged Children study

| | Age group | | | | | |
|-----------------------|-------------|-----|-------------|------|-------------|------|
| | 10–11 years | | 12–14 years | | 15–17 years | |
| Weight concern | % | n | % | n | % | n |
| No weight concern | 72.8 | 561 | 67.0 | 2063 | 65.8 | 1537 |
| Weight 'loss' concern | 22.6 | 174 | 26.0 | 800 | 24.8 | 578 |
| Weight 'gain' concern | 4.7 | 36 | 7.1 | 218 | 9.4 | 220 |

Table 3 Association between health indicators and weight concerns among Irish boys aged 10–17 years (*n* 6187); 2010 Irish Health Behaviour in School-aged Children study

| | Excellent self-reported health | | Frequent emotional symptoms | | Frequent physical symptoms | | High life satisfaction | | Feeling very happy | |
|-----------------------|--------------------------------|------------|-----------------------------|------------|----------------------------|------------|------------------------|------------|--------------------|------------|
| | OR† | 95% CI | OR† | 95% CI | OR† | 95% CI | OR† | 95% CI | OR† | 95% CI |
| Weight 'loss' concern | 0.31 | 0.27, 0.36 | 1.86 | 1.62, 2.15 | 1.78 | 1.50, 2.12 | 0.49 | 0.42, 0.57 | 0.56 | 0.49, 0.77 |
| Weight 'gain' concern | 0.49 | 0.39, 0.62 | 2.25 | 1.79, 2.82 | 2.41 | 1.87, 3.11 | 0.49 | 0.38, 0.62 | 0.48 | 0.39, 0.60 |

†The reference category is the group of boys who reported their weight was fine (OR = 1.00); adjusted for age group and social class.

Table 4 Logistic regression models predicting weight concerns (crude and adjusted odds ratios) among Irish boys aged 10–17 years (*n* 6187) with different family factors; 2010 Irish Health Behaviour in School-aged Children study

| Family factor | Weight 'loss' concern | | | Weight 'gain' concern | | |
|---|-----------------------|--------------|--------------------|-----------------------|--------------|--------------------|
| | Crude OR | Adjusted OR† | 95% CI (adjusted†) | Crude OR | Adjusted OR† | 95% CI (adjusted†) |
| Breakfast with mother/father every day | 0.72*** | 0.80* | 0.66, 0.97 | 0.61** | 0.72 | 0.51, 1.03 |
| Evening meal with mother/father every day | 0.92 | 1.05 | 0.90, 1.22 | 0.70** | 0.77* | 0.60, 0.99 |
| Ease of communication with father | 0.56*** | 0.69*** | 0.57, 0.82 | 0.56*** | 0.77 | 0.57, 1.02 |
| Ease of communication with mother | 0.59*** | 0.74*** | 0.60, 0.90 | 0.53*** | 0.61** | 0.44, 0.82 |
| Father present in the home | 0.77** | 0.81* | 0.67, 0.98 | 0.81* | 1.08 | 0.76, 1.52 |
| Mother working | 0.94* | 0.92 | 0.79, 1.08 | 1.05 | 1.06 | 0.82, 1.34 |

* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$;

†Adjusted for all family factors, age group and social class.

Weight concerns were not associated with maternal employment in the adjusted analysis, while boys whose fathers were present in the home were less likely to report a weight 'loss' concern in the adjusted analysis.

Discussion

The main objectives of the present study were to explore weight concerns among boys in Ireland, to describe physical and emotional symptoms among those with different weight concerns and to investigate the association between family characteristics and weight concerns among boys. It was important to differentiate between weight 'loss' and weight 'gain' concerns because increasing muscle and gaining weight is particularly pertinent to boys^(1,24,46).

Overall, 32.8% of Irish boys aged 10–17 years reported weight concerns. Weight 'loss' concerns (dieting or desire to lose weight; 25.1%) are in line with dieting prevalence among boys reported elsewhere^(3,4). Weight 'gain' concerns (8%) were lower than in studies in the USA (22%), even though the age groups (14.2 (SD 1.8) years *v.* 14.9 (SD 1.7) years) were comparable⁽³⁾. Concerns about muscularity are reported to emerge in mid-adolescence (14–16 years)^(47,48) and an association with age was evident in the present study.

Lower self-rated health, life satisfaction and levels of happiness, in addition to more frequent emotional and physical symptoms, were significantly more likely to be reported by boys with weight concerns, regardless of the

type of concern. Thus boys appear to experience similar types of negative consequences to girls who diet^(4–8). Future research is needed to explore both the type and healthiness of weight-control behaviours employed, particularly as unhealthy dieting among adolescents has been shown to cluster with other health-comprising behaviours^(4,5,49). Whether this association would hold for younger age groups is also worth exploring.

There is no single solution to addressing adolescent weight concerns. Several different and complementary preventive approaches are necessary, such as public policies, school and computer programmes, all of which can play a role in promoting healthy behaviours, reducing weight stigma and building self-esteem among young people⁽⁵⁰⁾. The present paper demonstrates the need for public health action to focus on family factors in addressing weight concerns of boys.

In general, the findings indicate that regular family meals are inversely associated with weight concerns among boys. Our findings are not easily comparable to existing studies, which focused on specific weight-loss behaviours among boys, such as purging, and are in fact equivocal for an effect of frequent family meals on such behaviours^(29,31,51). Nevertheless, family mealtimes are an opportunity to develop strong relationships and a sense of belonging and connectedness. Thus the weakening of the association between mealtimes and weight concerns when other family factors such as communication were included in the analyses may not be surprising. Indeed, this mirrors other work exploring the influence of family connectedness on associations between boys' extreme weight-control

behaviours and family meal frequency⁽²⁸⁾. Given the potential positive relationship between regular family meals and weight concerns, efforts to promote and support frequent mealtimes should be endorsed. The barriers to frequent joint mealtimes for both young people and their parents should be explored as the facilitators required may differ for adults and their children.

Ease of communication with father and mother was inversely associated with weight concerns. While statistical significance was lost in the adjusted analysis for communication with father and weight 'gain' concerns, the direction and magnitude remained. Parental support is related to the well-being of adolescents and lack of support is predictive of a range of negative behaviours and outcomes^(33,52–54). An open relationship between parents and their sons may be another mechanism for protecting against weight concerns. Cross-nationally, poor communication with fathers was positively associated with a desire to lose weight, after controlling for communication with the other parent, age and BMI, among adolescent boys in fourteen out of twenty-four countries examined⁽³⁸⁾. However, poor communication with mothers was rarely associated with a desire to lose weight among boys. Desire to gain weight has not been explored cross-nationally. The potential protective role of an open mother–son relationship on weight concerns, observed here, may reflect the ease with which adolescents share information or indeed teenage concerns with their mothers and that they perceive their mothers as trustworthy⁽⁵⁵⁾. Clearly mother–son and father–son relationships are important for weight concerns among adolescent boys in Ireland.

The presence of the father in the home was inversely associated with weight 'loss' concerns, but not weight 'gain' concerns in the full model. The same pattern was found for communication with fathers. Perhaps the mechanisms through which fathers influence concerns about weight 'gain' differ from those about weight 'loss'. Exploring father's own body image, weight-control behaviours and attitudes to body shape and muscle building may shed some light on the nuances of these relationships. The impact of family structure on adolescent health could be affected by the stability or a recent change in family structure, which was not captured here. In a Swedish study relational content was more important for adolescent well-being than family structure⁽⁵⁶⁾; and among 13- and 15-year-old adolescents in Scotland, family structure was of less importance than the quality of the parent–child relationships on adolescent life satisfaction, although the impact of family structure on life satisfaction did persist for boys⁽⁵⁷⁾. The strong and consistent relationship between communication and weight concerns observed in the present study suggests that this aspect of child–parent relationships should be the focus going forward.

Limitations of the current study include the cross-sectional design, making it impossible to determine causality, and the reliance on self-report data; however, students' responses were anonymous and therefore participants had no reason

to misreport their responses. The health indicators used in the study such as life satisfaction, a cognitive aspect of well-being, and happiness, a facet of well-being, can be influenced by many life experiences and relationships in addition to school-related factors^(58,59). An exploration of other variables contributing to these outcomes as well as physical and emotional symptoms was not possible here. Strengths include the large sample size, inclusion of children at a national level and the focus on boys.

To conclude, body weight concerns are associated with negative health outcomes as reported by adolescent boys in Ireland. The type of weight-control behaviours used by boys in Ireland deserves further attention particularly in light of the emphasis on muscularity in recent times. Boys reporting an open communication with their fathers and mothers and regular family meals were, for the most part, less likely to report a weight concern. The dimensions of relations between parents and adolescent boys deserve further attention so that particular relationship skills can be targeted if required. Qualitative research with fathers and mothers on their perceived role in influencing weight concerns among their sons would also prove insightful.

Acknowledgements

Acknowledgements: The authors acknowledge all of the parents and children who consented and participated, as well as the management authorities, principals and teachers in all schools who participated. HBSC is an international study carried out in collaboration with WHO/EURO. The International Coordinator of HBSC is Professor Candace Currie (University of St. Andrews, Scotland); Data Bank Manager is Professor Oddrun Samdal (University of Bergen). For details, see <http://www.hbsc.org/>. *Financial support:* This work was supported by the Department of Health, Ireland. The Department of Health, Ireland had no role in the design, analysis or writing of this article. *Conflict of interest:* None. *Authorship:* C.K. participated in the design and data collection for the study, conceived of the study questions and drafted the manuscript. A.F., M.S. and J.G. contributed to the statistical analysis. M.M. and S.N.G. participated in the design and data collection for the study and edited the manuscript. All authors have read and approved the final manuscript. *Ethics of human subject participation:* Ethical approval was obtained from the National University of Ireland, Galway Research Ethics Committee.

References

1. Ricciardelli L & McCabe M (2011) Body image development in adolescent boys. In *Body Image: A Handbook of Theory, Research and Clinical Practice*, pp. 85–92 [TF Cash and T Pruzinsky, editors]. New York: Guilford Press.
2. Smolak L (2004) Body image in children and adolescents: where do we go from here? *Body Image* **1**, 15–28.

3. Neumark-Sztainer D, Story M, Hannan PJ *et al.* (2002) Weight-related concerns and behaviors among overweight and nonoverweight adolescents: implications for preventing weight-related disorders. *Arch Pediatr Adolesc Med* **156**, 171–178.
4. Crow S, Eisenberg ME, Story M *et al.* (2006) Psychosocial and behavioral correlates of dieting among overweight and non-overweight adolescents. *J Adolesc Health* **38**, 569–574.
5. French SA, Story M, Downes B *et al.* (1995) Frequent dieting among adolescents: psychosocial and health behaviour correlates. *Am J Public Health* **85**, 695–701.
6. Pesa J (1999) Psychosocial factors associated with dieting behaviors among female adolescents. *J Sch Health* **69**, 196–201.
7. Canpolat BI, Orsel S & Akdemir A (2005) The relationship between dieting and body image, body ideal, self-perception, and body mass index in Turkish adolescents. *Int J Eat Disord* **37**, 150–155.
8. Neumark-Sztainer D & Hannan PJ (2000) Weight-related behaviors among adolescent girls and boys: results from a national survey. *Arch Pediatr Adolesc Med* **154**, 569–577.
9. Stice E, Hayward C, Cameron RP *et al.* (2000) Body-image and eating disturbances predict onset of depression among female adolescents: a longitudinal study. *J Abnorm Psychol* **109**, 438–444.
10. Siegal JM (2002) Body image change and adolescent depressive symptoms. *J Adolesc Res* **17**, 27–41.
11. Thompson AM & Chad KE (2002) The relationship of social physique anxiety to risk for developing an eating disorder in young females. *J Adolesc Health* **31**, 183–189.
12. Paxton SJ, Neumark-Sztainer D, Hannan PJ *et al.* (2006) Body dissatisfaction prospectively predicts depressive mood and low self-esteem in adolescent girls and boys. *J Clin Child Adolesc Psychol* **35**, 539–549.
13. Kann L, Kinchen S, Shanklin S *et al.* (2014) Youth risk behavior surveillance – United States, 2013. *MMWR Surveill Summ* **63**, Suppl. 4, 1–168.
14. Nowak M (1998) The weight-conscious adolescent: body image, food intake, and weight-related behaviour. *J Adolesc Health* **23**, 389–398.
15. Ojala K (2012) Weight-reduction behaviour. In *Social Determinants of Health and Well-Being Among Young People: Health Behaviour in School-Aged Children (HBSC) Study: International Report from the 2009/2010 Survey*, pp. 97–100 [C Currie, C Zanotti, A Morgan *et al.*, editors]. Copenhagen: WHO Regional Office for Europe.
16. Cooper PJ & Goodyer I (1997) Prevalence and significance of weight and shape concerns in girls aged 11–16. *Br J Psychiatry* **171**, 542–544.
17. Bearman S, Presnell K, Martinez E *et al.* (2006) The skinny on body dissatisfaction: a longitudinal study of adolescent girls and boys. *J Youth Adolesc* **35**, 217–229.
18. Hargreaves DA & Tiggemann M (2004) 'Body image is for girls': a qualitative study of boys' body image. *J Health Psychol* **11**, 567–576.
19. Nishina A, Ammon NY, Bellmore AD *et al.* (2006) Body dissatisfaction and physical development among ethnic minority adolescents. *J Youth Adolesc* **35**, 179–191.
20. Presnell K, Bearman SK & Stice E (2004) Risk factors for body dissatisfaction in adolescent boys and girls: a prospective study. *Int J Eat Disord* **36**, 389–401.
21. Ryan TA & Morrison T (2009) Factors perceived to influence young Irish men's body image investment: a qualitative investigation. *Int J Mens Health* **8**, 213–234.
22. McSharry M (2011) *Schooled Bodies?: Negotiating Adolescent Validation Through Press, Peers and Parents*. Stoke-on-Trent: Trentham Books Ltd.
23. Nemeth A (2012) Body image. In *Social Determinants of Health and Well-Being Among Young People: Health Behaviour in School-Aged Children (HBSC) Study: International Report from the 2009/2010 Survey*. pp. 93–96 [C Currie, C Zanotti, A Morgan *et al.*, editors]. Copenhagen: WHO Regional Office for Europe.
24. Field AE, Austin SB, Camargo CA Jr *et al.* (2005) Exposure to the mass media, body shape concerns, and use of supplements to improve weight and shape among male and female adolescents. *Pediatrics* **116**, e214–e220.
25. Eisenberg ME, Wall M & Neumark-Sztainer D (2012) Muscle-enhancing behaviours among adolescent girls and boys. *Pediatrics* **130**, 1019–1026.
26. Westmoreland Corson P & Andersen AE (2002) Body image issues among boys and men. In *Body Image: A Handbook of Theory, Research, and Clinical Practice*, pp. 192–199 [TF Cash and Pruzinsky T, editors]. New York: Guilford Press.
27. Fulkerson JA, Larson N, Horning M *et al.* (2014) A review of associations between family or shared meal frequency and dietary and weight status outcomes across the lifespan. *J Nutr Educ Behav* **46**, 2–19.
28. Neumark-Sztainer D, Wall M, Story M *et al.* (2004) Are family meal patterns associated with disordered eating behaviors among adolescents? *J Adolesc Health* **35**, 350–359.
29. Neumark-Sztainer D, Eisenberg ME & Fulkerson JA (2008) Family meals and disordered eating in adolescents: longitudinal findings from Project EAT. *Arch Pediatr Adolesc Med* **162**, 17–22.
30. Fulkerson JA, Story M, Mellin A *et al.* (2006) Family dinner meal frequency and adolescent development: relationships with developmental assets and high-risk behaviors. *J Adolesc Health* **39**, 337–345.
31. Skeer MR & Ballard EL (2013) Are family meals as good for youth as we think they are? A review of the literature on family meals as they pertain to adolescent risk prevention. *J Youth Adolesc* **42**, 943–963.
32. Welsh EM, French S & Wall M (2011) Examining the relationship between family meal frequency and individual dietary intake: does family cohesion play a role? *J Nutr Educ Behav* **43**, 229–235.
33. Sweeting H & West P (2005) Dietary habits and children's family lives. *J Hum Nutr Diet* **18**, 93–97.
34. Pearson N, MacFarlane A, Crawford D *et al.* (2009) Family circumstance and adolescent dietary behaviours. *Appetite* **52**, 668–674.
35. von Soest T & Wichstrom L (2009) Gender differences in the development of dieting from adolescence to early adulthood: a longitudinal study. *J Res Adolesc* **19**, 509–529.
36. Saling M, Ricciardelli L & McCabe MP (2005) A prospective study of individual factors in the development of weight and muscle concerns among preadolescent children. *J Youth Adolesc* **34**, 651–661.
37. Hill AJ & Franklin JA (1998) Mothers, daughters and dieting: investigating the transmission of weight control. *Br J Clin Psychol* **37**, 3–13.
38. Al Sabbah H, Vereecken C, Elgar F *et al.* (2009) Body weight dissatisfaction and communication with parents among adolescents in 24 countries: international cross-sectional survey. *BMC Public Health* **9**, 52.
39. Vander Wal SJ (2012) The relationship between body mass index and unhealthy weight control behaviors among adolescents: the role of family and peer social support. *Econ Hum Biol* **10**, 395–404.
40. Brooks F, Zaboriskis A, Örkényi A *et al.* (2012) Family: communication with father. In *Social Determinants of Health and Well-Being Among Young People: Health Behaviour in School-aged Children (HBSC) Study: International Report from the 2009/2010 Survey*, pp. 23–26 [C Currie, C Zanotti, A Morgan *et al.*, editors]. Copenhagen: WHO Regional Office for Europe.
41. Gavin A, Molcho M, Kelly C *et al.* (2013) *The HBSC Ireland Trends Report 1998–2010: Child Health Behaviours, Outcomes and Contexts*. Dublin: NUI Galway.

42. Levin KA & Currie C (2014) Reliability and validity of an adapted version of the Cantril Ladder for use with adolescent samples. *Soc Indic Res* **119**, 1047–1063.
43. Cantril H (1965) *The Pattern of Human Concerns*. New Brunswick, NJ: Rutgers University Press.
44. Ravens-Sieberer U, Erhart M, Torsheim T *et al.* (2008) An international scoring system for self-reported health complaints in adolescents. *Eur J Public Health* **18**, 294–299.
45. Shaffer DR & Kipp K (2013) *Developmental Psychology: Childhood and Adolescence*, 9th ed. Belmont, CA: Wadsworth.
46. Jones D & Crawford J (2005) Adolescent boys and body image: weight and muscularity concerns as dual pathways to body dissatisfaction. *J Youth Adolesc* **34**, 629–636.
47. Polce-Lynch M, Myers BJ, Kilmartin CT *et al.* (1998) Gender and age patterns in emotional expression, body image, and self-esteem: a qualitative analysis. *Sex Roles* **38**, 1025–1048.
48. Cortese S, Falissard B, Pigiani Y *et al.* (2010) The relationship between body mass index and body size dissatisfaction in young adolescents: spline function analysis. *J Am Diet Assoc* **110**, 1098–1102.
49. Neumark-Sztainer D, Story M, Dixon LB *et al.* (1998) Adolescents engaging in unhealthy weight control behaviors: are they at risk for other health-compromising behaviors? *Am J Public Health* **88**, 952–955.
50. Cash TF & Smolak L (editors) (2012) *Body Image: A Handbook of Science, Practice and Prevention*. New York: Guildford Press.
51. Haines J, Gillman MW, Rifas-Shiman S *et al.* (2010) Family dinner and disordered eating behaviours in a large cohort of adolescents. *Eat Disorders* **18**, 10–24.
52. Parker JS & Benson MJ (2004) Parent–adolescent relations and adolescent functioning: self-esteem, substance abuse, and delinquency. *Adolescence* **39**, 519–530.
53. Wills TA, Resko JA, Ainette MG *et al.* (2004) Role of parent support and peer support in adolescent substance use: a test of mediated effects. *Psychol Addict Behav* **18**, 122–134.
54. Young JF, Berenson K, Cohen P *et al.* (2005) The role of parent and peer support in predicting adolescent depression: a longitudinal community study. *J Res Adolesc* **15**, 407–423.
55. Tamara D, Afifi A & Aldeis D (2008) Why can't we just talk about it? Parents' and adolescents' conversations about sex. *J Adolesc Res* **23**, 689–721.
56. Låftman SB & Östberg V (2006) The pros and cons of social relations: an analysis of adolescents' health complaints. *Soc Sci Med* **63**, 611–623.
57. Levin KA & Currie C (2010) Family structure, mother–child communication, father–child communication, and adolescent life satisfaction: a cross-sectional multilevel analysis. *Health Educ* **110**, 152–168.
58. Diseth A, Danielsen AG & Samdal O (2012) A path analysis of basic need support, self-efficacy, achievement goals, life satisfaction and academic achievement level among secondary school students. *Educ Psychol* **32**, 335–354.
59. Levin KA, Inchley J, Currie D *et al.* (2012) Subjective health and mental well-being of adolescents and the health promoting school: a cross-sectional multilevel analysis. *Health Educ* **112**, 170–184.