

A CLASS OF ITERATION METHODS FOR THE MATRIX EQUATION
 $AXB = C$

KONGHUA GUO, XIYAN HU AND LEI ZHANG

An iteration method for the matrix equation $AXB = C$ is constructed. By this iteration method, the least-norm solution for the matrix equation can be obtained when the matrix equation is consistent and the least-norm least-squares solutions can be obtained when the matrix equation is not consistent. The related optimal approximation solution is obtained by this iteration method. A preconditioned method for improving the iteration rate is put forward. Finally, some numerical examples are given.

1. INTRODUCTION

The matrix equation problem is an active research topic in computational mathematics, and has been widely applied in various areas, such as structural design, system identification, principal component analysis, exploration and remote sensing, biology, electricity, solid mechanics, molecular spectroscopy, structural dynamics, automatics control theory, vibration theory, and so on.

We use R^n to denote the set of all real vectors of n dimensions, I_n the identity matrix of order n , and $R^{n \times m}$ all $n \times m$ real matrices. Let $\|A\|_F$, A^+ , A^T denote especially the Frobenius norm, the Moore–Penrose generalised inverse, and the transpose of a matrix A . ($\text{tr}(A)$ means the trace of matrix A , $R(A)$ the column space of matrix A), $R^\perp(A)$ the orthogonal complement space of $R(A)$, and for any $A \in R^{m \times n}$, $B \in R^{n \times p}$, $A \otimes B$ means the Kronecker product of the matrices A and B .

The following problems are considered in this paper.

PROBLEM 1.1. Given $A \in R^{m \times n}$, $B \in R^{n \times p}$, $C \in R^{m \times p}$, find $X \in R^{n \times n}$, such that

$$(1) \quad AXB = C$$

PROBLEM 1.2. Suppose Problem 1.1 is consistent, and its solution set is S_E , for $X_0 \in R^{n \times n}$. Find $\hat{X} \in S_E$, such that

$$(2) \quad \|\hat{X} - X_0\|_F = \min_{X \in S_E} \|X - X_0\|_F$$

Received 18th October, 2006

Research supported by National Natural Science Foundation of China(10571047) and Scientific Research Fund of Hunan Provincial Education Department(06C095)

Copyright Clearance Centre, Inc. Serial-fee code: 0004-9727/07 \$A2.00+0.00.

In fact, Problem 1.2 is to find the optimal approximation solution to the given matrix X_0 . In 1955, Penrose obtained necessary and sufficient conditions for solving Problem 1.1 and the general expressions of the solution [10]. Since then Problem 1.1 has been considered in the case of some special solution structures, for example, symmetric, triangular or diagonal solution X . We can refer to Hua [3], Chu [2], Don [4], Magnus [6], Morris and Odell [8], Bjerhammer [1] for more details. Mitra [7] considered common solutions to a pair of linear matrix equations $A_1XB_1 = C_1, A_2XB_2 = C_2$. In these papers, the problem was discussed by using matrix decompositions such as the singular value decomposition, the generalised single valued decomposition, the quotient single valued decomposition and the canonical correlation decomposition. However, it is difficult to apply these methods to solving problems such as finding symmetric solutions of the matrix equation $AXB = C$. In 2005, Y.X. Peng put forward an iteration method for finding symmetric solutions of the matrix equation $AXB = C$ ([9]). The advantage of this iteration method is that when the problem is consistent, its solution can be obtained theoretically within a finite number of steps, and the disadvantage of the method is that the convergence rate can not be analysed.

In this paper, we construct a new iterative method for the matrix equation $AXB = C$, by which we can obtain the least-norm solution of Problem 1.1 when the problem is consistent and obtain the least-norm least-squares solution of Problem 1.1 when the problem is not consistent. Furthermore, we show that the convergence rate of the method is related to the singular value of the matrix A , and so the iteration method can be improved by some preconditioned methods. When the solution set of Problem 1.1 is not empty, Problem 1.2 has a unique solution and we can obtain it by the iteration method.

The paper is organised as follows: In Section 2 we first introduce a new iterative method for finding the matrix equation $AXB = C$ and prove the convergence of the method. In Section 3 we solve Problem 1.2 by using this iteration method. In Section 4 we propose an improvement of the iteration method in order to increase the convergence rate. In the last section, we shall give some numerical examples to verify the method and compare the convergence rate between the original method and the improved method.

2. THE SOLUTION OF PROBLEM 1.1

In this section, we shall introduce a new iteration method for solving Problem 1.1, and then we shall prove the convergence of the iteration method.

ITERATION METHOD 2.1.

- step1: Select $C_0 = C, X_0 = O$;
- step2: Let $\alpha_k = \frac{\|A^T C_k B^T\|_F^2}{\|AA^T C_k B^T B\|_F^2} (k = 0, 1, 2, \dots)$;
- step3: Let $\Delta X_k = \alpha_k A^T C_k B^T, (k = 0, 1, 2, \dots)$;
- step4: If $\Delta X_k = 0$, stop, otherwise, let $X_{k+1} = X_k + \Delta X_k, (k = 0, 1, 2, \dots)$;

step5: Let $C_{k+1} = C_k - A\Delta X_k B$, ($k = 0, 1, 2, \dots$), goto step2.

DEFINITION 2.1: Suppose $A, B \in R^{m \times n}$, then $\text{tr}(A^T B)$ is called the inner product of the matrices A, B , denoted by $\langle A, B \rangle$.

DEFINITION 2.2: Assume $A, B \in R^{m \times n}$. If $\langle A, B \rangle = 0$, that is, $\text{tr}(A^T B) = 0$, then the matrices A, B are called orthogonal each other.

LEMMA 2.1. In the iteration method 2.1, the selection of α_k makes $\|C_{k+1}\|_F$ minimal, and make C_{k+1} and $A\Delta X_k B$ orthogonal.

PROOF: For the iteration method 2.1, we have

$$\begin{aligned} \|C_{k+1}\|_F^2 &= \langle C_k - \alpha_k A^T C_k B^T, C_k - \alpha_k A^T C_k B^T \rangle \\ &= \|C_k\|_F^2 - 2\alpha_k \langle C_k, A A^T C_k B^T B \rangle + \alpha_k^2 \|A A^T C_k B^T B\|_F^2 \end{aligned}$$

From the above expression, we know that the necessary and sufficient conditions of making $\|C_{k+1}\|_F$ the minimal is that

$$\alpha_k = \frac{\|A^T C_k B^T\|_F^2}{\|A A^T C_k B^T B\|_F^2}$$

On the other hand, Let $\langle C_{k+1}, A\Delta X_k B \rangle = 0$, we also have that

$$\alpha_k = \frac{\|A^T C_k B^T\|_F^2}{\|A A^T C_k B^T B\|_F^2}$$

Hence, in Iteration method 2.1, selecting

$$\alpha_k = \|A^T C_k B^T\|_F^2 / \|A A^T C_k B^T B\|_F^2$$

will make C_{k+1} and $A\Delta X_k B$ orthogonal. □

LEMMA 2.2. In the iteration method 2.1, we have $\|C_{k+1}\|_F^2 = \|C_k\|_F^2 - \|A\Delta X_k B\|_F^2$

PROOF: From the step5 of Iteration method 2.1, we have $C_k = C_{k+1} + A\Delta X_k B$, and so, $\|C_k\|_F^2 = \|C_{k+1} + A\Delta X_k B\|_F^2$, according to Lemma 2.1, then we have

$$\|C_k\|_F^2 = \|C_{k+1}\|_F^2 + \|A\Delta X_k B\|_F^2$$

Hence,

$$\|C_{k+1}\|_F^2 = \|C_k\|_F^2 - \|A\Delta X_k B\|_F^2$$
□

DEFINITION 2.3: For $A = (a_{ij})_{m \times n} \in R^{m \times n}$, denote by $\text{vec}(A)$ the following vector containing all the entries of matrix A :

$$\text{vec}(A) = [a_{11}, \dots, a_{1n}, a_{21}, \dots, a_{2n}, \dots, a_{m1}, \dots, a_{mn}]^T,$$

then $\text{vec}(A)$ is called straightening of the matrix A .

It is evident that the transform $A \rightarrow \text{vec}(A)$ gives a linear isomorph of $R^{m \times n} \rightarrow R^{mn}$.

LEMMA 2.3. ([5]) *For any matrices A, B and C in suitable size, we have*

$$\text{vec}(A + B) = \text{vec}(A) + \text{vec}(B), \text{vec}(ABC) = (A \otimes C^T) \text{vec}(B)$$

LEMMA 2.4. *Suppose that the consistent system of linear equations $My = b$ has a solution $y_0 \in R(M^T)$, then y_0 is the least-norm solution of the system of linear equations.*

PROOF: We decompose the matrix M by single valued decomposition:

$$M = U \begin{pmatrix} \Sigma & 0 \\ 0 & 0 \end{pmatrix} V^T = U_1 \Sigma V_1^T$$

where $U = (U_1, U_2)$ and $V = (V_1, V_2)$ are orthogonal matrices. Then the Moore-Penrose generalised inverse of matrix M is

$$M^+ = V_1 \Sigma^+ U_1^T$$

and the general solution of the system of linear equations $My = b$ is

$$y = M^+ b + (I - M^+ M)z$$

where z is an arbitrary vector with suitable size.

Since $M^+ = V_1 \Sigma^+ U_1^T \in R(V_1)$, $(I - M^+ M)z = (I - V_1 V_1^T)z = V_2 V_2^T z \in R(V_2)$, and V_2 and V_1 are orthogonal each other; that is, $\text{tr}(V_2^T V_1) = 0$, then $M^+ b$ is the least-norm solution of the system of linear equations $My = b$.

On the other hand, $M^T = V_1 \Sigma U_1^T$, the solution $y_0 \in R(M^T)$, therefore y_0 is the least-norm solution of the system of linear equations $My = b$.

Obviously, the set of solutions of the system of linear equations $My = b$ is closed convex, and so the least-norm solution of the system is unique. □

Similarly, we have the following lemma.

LEMMA 2.5. *Suppose that the inconsistent system of linear equations $My = b$ has a solution $y_0 \in R(M^T)$, then y_0 is the least-norm least-squares solution of the system of linear equations, and the solution is unique.*

LEMMA 2.6. ([11]) *The matrix equation $AXB = C$ has a unique least-norm solution $X = A^+ C B^+$ when the equation is consistent, and has a unique least-norm least-squares solution $X = A^+ C B^+$ when the equation is not consistent.*

DEFINITION 2.4: Let $A, B \in R^{m \times n}$. If $\cos \theta = \langle A, B \rangle / (\|A\|_F \cdot \|B\|_F)$ ($0 \leq \theta \leq \pi$), then θ is called the included angle of the matrices A, B .

THEOREM 2.1. *The iteration method 2.1 is convergent. Let the maximum singular value and the minimum singular value of the matrix A be σ_1, σ_r , the maximum singular value and the minimum singular value of the matrix B be λ_1, λ_s . Then the convergence rate of the iteration method 2.1 is no less than $-0.5 \ln(1 - (\sigma_r^2 \lambda_s^2)/(\sigma_1^2 \lambda_1^2))$.*

PROOF: . Suppose $\text{rank}(A) = r$, $\text{rank}(B) = s$, and the singular value decompositions of matrix A and matrix B are

$$A = UDV^T = U_1 \Sigma V_1^T, B = PEQ^T = P_1 \Lambda Q_1^T$$

where $U = (U_1, U_2), V = (V_1, V_2), P = (P_1, P_2)$, and $Q = (Q_1, Q_2)$ are orthogonal matrices,, $D = \begin{pmatrix} \Sigma & O \\ O & O \end{pmatrix}, E = \begin{pmatrix} \Lambda & O \\ O & O \end{pmatrix}, \Sigma = \text{diag}(\sigma_1, \sigma_2, \dots, \sigma_r), \sigma_i (i = 1, 2, \dots, r)$ are the singular values of matrix A ; $\Lambda = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_s), \lambda_i (i = 1, 2, \dots, s)$ are the singular values of matrix B .

If Problem 1.1 is consistent, then for C_k there exists a matrix G which makes $C_k = UGQ^T$, where $G = (g_{ij} = (g_1, \dots, g_n), g_i \in R^n, (i = 1, 2, \dots, n), g_i = 0, i > s, g_i \in R, (i, j = 1, 2, \dots, n), g_{ij} = 0, i > r, j > s$

Let θ be the included angle of C_k and $A\Delta X_k B$, then we have that

$$\begin{aligned} \cos(\theta) &= \frac{(C_k, A\Delta X_k B)}{\|C_k\|_F \cdot \|A\Delta X_k B\|_F} = \frac{\|A^T C_k B^T\|_F^2}{\|C_k\|_F \cdot \|AA^T C_k B^T B\|_F} \\ &= \frac{\|VD^T U^T UGQ^T QE^T P\|_F^2}{\|UGQ^T\|_F \cdot \|UDV^T VD^T U^T UGQ^T QE^T P^T PEQ^T\|_F} \\ &= \frac{\|D^T GE\|_F^2}{\|G\|_F \cdot \|DD^T GE^T E\|_F} = \frac{\text{tr}(EG^T DD^T GE^T)}{(\text{tr}(G^T G))^{1/2} \cdot (\text{tr}(E^T EG^T DD^T DD^T GE^T E))^{1/2}} \\ &= \frac{\sum_{j=1}^s \sum_{i=1}^r \sigma_i^2 \lambda_j^2 g_{ij}^2}{(\sum_{i=1}^s g_i^T g_i)^{1/2} \cdot (\sum_{j=1}^r \sum_{i=1}^s \sigma_i^4 \lambda_j^4 g_{ij}^2)^{1/2}} \\ &\geq \frac{\sum_{j=1}^s \sum_{i=1}^r \sigma_i^2 \lambda_j^2 g_{ij}^2}{(\sum_{i=1}^s g_i^T g_i)^{1/2} \cdot (\sum_{j=1}^r \sum_{i=1}^s \sigma_i^2 \lambda_j^2 g_{ij}^2)^{1/2} \sigma_1 \lambda_1} = \frac{(\sum_{j=1}^s \sum_{i=1}^r \sigma_i^2 \lambda_j^2 g_{ij}^2)^{1/2}}{(\sum_{i=1}^s g_i^T g_i)^{1/2} \cdot \sigma_1 \lambda_1} \geq \frac{\sigma_r \lambda_s}{\sigma_1 \lambda_1} \end{aligned}$$

Notice that $\|C_k\|_F = \|C_{k+1}\|_F + \|A\Delta X_k B\|_F$, then we have

$$\|C_{k+1}\|_F = \|C_k\|_F \sin(\theta) \leq \sqrt{1 - \frac{\sigma_r^2 \lambda_s^2}{\sigma_1^2 \lambda_1^2}}$$

Therefore, Iteration method 2.1 is convergent, and the convergence rate of the iteration method(2.1) is no less than $-0.5 \ln(1 - (\sigma_r^2 \lambda_s^2)/(\sigma_1^2 \lambda_1^2))$. □

THEOREM 2.2. *The iteration method 2.1 will converge to the least-norm solution of Problem 1.1 when the problem is consistent and will converge to the least-norm least-square solution of Problem 1.1 when the problem is not consistent.*

PROOF: According to Theorem 2.1, if Problem 1.1 is consistent, we can obtain a solution X^* by Iteration method 2.1, and the solution X^* can be represented as that

$$X^* = A^T Y B^T$$

In the sequel, we shall prove that the X^* is just the least-norm solution of Problem 1.1.

Denote $\text{vec}(X) = x, \text{vec}(X^*) = x^*, \text{vec}(Y) = y, \text{vec}(C) = b$, then the matrix equations $AXB = C$ is equivalent to the system of linear equations

$$(2.1) \quad (A \otimes B^T)x = b$$

Notice that

$$\begin{aligned} x^* &= \text{vec}(X^*) = \text{vec}(A^T Y B^T) = (A^T \otimes B)y \\ &= (A \otimes B^T)^T y \in R((A \otimes B^T)^T) \end{aligned}$$

So x^* is the least-norm solution of the system of linear equation 2.1 by Lemma 2.4, Since the vector operator is isomorphic, X^* is the unique least-norm solution of Problem 1.1.

If Problem 1.1 is not consistent, let $C = C^{(1)} + C^{(2)}$, where $C^{(1)} \in R(A)$ and $C^{(2)} \in R^\perp(A)$. For any $X \in R^{n \times n}$, $C^{(1)} - AXB \in R(A)$, and $C^{(1)} - AXB$ is orthogonal with $C^{(2)}$, so we have that

$$(2.2) \quad \|C - AXB\|_F^2 = \|C^{(1)} - AXB + C^{(2)}\|_F^2 = \|C^{(1)} - AXB\|_F^2 + \|C^{(2)}\|_F^2$$

which means that the sufficient and necessary condition of X being the least-squares solution of $AXB = C$ is that X is the solution of consistent equation $AXB = C^{(1)}$.

From the step5 of Iteration method 2.1, we have

$$(2.3) \quad C_{k+1}^{(1)} + C_{k+1}^{(2)} = C_0^{(1)} + C_0^{(2)} - AX_{k+1}B$$

Noticing that $AX_{k+1}B \in R(A)$, we know $C_{k+1}^{(2)} = C_0^{(2)}$, and (2.3) is equivalent to

$$(2.4) \quad C_{k+1}^{(1)} = C_0^{(1)} - AX_{k+1}B$$

Thus the iteration process is conducted in $R(A)$. Then from the iteration method 2.1, we can obtain the least-norm solution of the consistent equation $AXB = C^{(1)}$. It means that the iteration method will converge to the unique least-norm least-squares solution of Problem 1.1 when the problem is not consistent. □

3. THE SOLUTION OF PROBLEM 1.2

When Problem 1.1 is solvable, it is easy to test that S_E is a closed convex set. Hence we know that for the given $X_0 \in R^{n \times n}$, we can find a unique $\hat{X} \in S_E$ which will make $\|\hat{X} - X_0\|_F = \min_{X \in S_E} \|X - X_0\|_F$. Next we give the iteration method which find the $\hat{X} \in S_E$
 If S_E is not empty, for any $X \in S_E$,

$$AXB = C \Leftrightarrow A(X - X_0)B = C - AX_0B$$

Let $X^* = X - X_0$, $C^* = C - AX_0B$, then solving the problem 1.2 is equivalent to finding the least-norm solution \tilde{X}^* of the consistent matrix equation $AX^*B = C^*$, which can be obtained by using Iteration method 2.1, and the solution of the problem 1.2 can be represented as $\hat{X} = \tilde{X}^* + X_0$.

4. THE IMPROVEMENTS OF THE ITERATION METHOD

From Theorem 2.1, if the ratio $(\sigma_r \lambda_s)/(\sigma_1 \lambda_1)$ is near to 1, then the convergence of Iteration 2.1 will be fast, but if $\sigma_1 \lambda_1 \gg \sigma_r \lambda_s$, the convergence of Iteration Method 2.1 may be slow. To improve its convergence rate we may deal with the equation before solving it by using preconditioning methods.

In this paper, we adopt polynomial preconditioning methods to improve the convergence rate, that is, we transform the original equation $AXB = C$ to the equation $C(A)AXC(B) = C(A)CC(B)$, where $C(A), C(B)$ are polynomial on A, B with low order.

In next section, we shall give some example to verify Iteration method 2.1 and compare the convergence rate between original iteration method and preconditioned method.

5. EXAMPLE

In this section, we denote t as the computing time (unit:second), and k as the number of iterations. The computations were performed using MATLAB, version 6.5.1, under the operation system of Windows Me, and the CPU rate of the machine is 2.40GHz.

EXAMPLE 1. Let

$$A = \begin{pmatrix} 8.2462 & 9.0000 & 9.8954 & 10.3441 & 10.9545 & 1.0000 & 3.7417 & 5.1962 & 6.3246 & 7.2801 & 8.1240 \\ 8.9443 & 9.6437 & 10.2956 & 10.9087 & 3.3166 & 3.6056 & 5.0990 & 6.2450 & 7.2111 & 8.0623 & 8.1854 \\ 9.5917 & 10.2470 & 10.8628 & 3.1623 & 3.4641 & 5.0000 & 6.1644 & 7.1414 & 8.0000 & 8.7750 & 8.8882 \\ 10.1980 & 10.8167 & 3.0000 & 4.6904 & 4.8990 & 6.0828 & 7.0711 & 7.9373 & 8.7178 & 8.8318 & 9.5394 \\ 10.7703 & 2.8284 & 4.5826 & 4.7958 & 6.0000 & 7.0000 & 7.8740 & 8.6603 & 9.3808 & 9.4868 & 10.1489 \\ 2.6458 & 4.4721 & 5.7446 & 5.9161 & 6.9282 & 7.8102 & 8.6023 & 9.3274 & 9.4340 & 10.0995 & 10.7238 \\ 4.3589 & 5.6569 & 5.8310 & 6.8557 & 7.7460 & 8.5440 & 9.2736 & 9.9499 & 10.0499 & 10.6771 & 2.4495 \\ 5.5678 & 6.6332 & 6.7823 & 7.6811 & 8.4853 & 9.2195 & 9.8995 & 10.0000 & 10.6301 & 2.2361 & 4.2426 \\ 6.5574 & 6.7082 & 7.8158 & 8.4261 & 9.1652 & 9.8489 & 10.4881 & 10.5830 & 2.0000 & 4.1231 & 5.4772 \\ 7.4162 & 7.5498 & 8.3866 & 9.1104 & 9.7980 & 10.4403 & 10.5357 & 1.7321 & 4.0000 & 5.3852 & 6.4807 \\ 7.4633 & 8.3066 & 9.0554 & 9.7468 & 10.3923 & 11.0000 & 1.4142 & 3.8730 & 5.2915 & 6.4031 & 7.3485 \end{pmatrix},$$

$B = A$, and let $C =$

904.6936	913.2006	907.3617	881.3187	820.4256	655.7400	751.2413	810.0672	853.1009	883.1039	898.1381
923.8008	932.8133	927.1399	900.7094	838.2387	667.1675	765.4937	826.0158	870.3427	901.3351	916.9743
911.8758	921.1333	915.9313	890.2094	828.5479	656.6647	754.3255	814.3715	858.4038	889.2931	905.0192
892.7852	901.9109	897.0814	872.4388	812.9149	644.4845	739.5227	797.8934	840.6953	870.7534	886.1313
835.6333	843.6018	838.9989	816.6930	763.3729	611.6509	697.6750	750.4725	789.0757	816.0686	829.8106
725.1206	728.6117	722.1176	702.2365	660.9599	563.0691	626.5100	665.8320	694.0489	712.9421	721.5573
784.2758	789.1816	782.7550	760.8806	713.6275	595.9011	669.1322	714.4161	747.1302	769.3535	779.8532
834.0055	840.0351	833.5978	809.9872	757.7157	623.9719	705.2647	755.4724	791.8953	816.8518	828.8779
862.2418	869.0409	862.6897	838.1054	782.7825	638.6341	724.9469	778.2080	816.9464	843.6336	856.6556
890.7232	898.3408	892.1058	866.5343	808.0409	653.1216	744.5845	800.9802	842.1045	870.5854	884.6472
895.9263	903.8165	897.6516	871.8114	812.3613	654.3113	747.2089	804.4829	846.2916	875.3039	889.6836

- (1) Find the least-norm solution of Problem 1.1.
- (2) Let S_E denote the set of all solutions of the matrix equation $AXB = C$, suppose $X_0 =$

1.3701	-4.8758	2.1230	0.7781	3.6417	2.5740	2.4194	4.3440	0.3185	3.9318	-3.0110
-0.6231	-1.6490	2.3485	1.0349	-5.6921	-4.2331	0.0966	-1.6502	3.5918	3.8883	-3.0388
-2.7381	2.0104	-1.1974	-0.5989	1.7949	5.4612	-2.5518	-4.5650	2.7999	3.0393	-3.0322
-3.9526	3.6712	-2.2277	-1.0113	4.2979	-5.3661	1.5700	2.0865	0.2286	1.2513	-3.0349
-4.1968	2.2302	0.6011	0.5023	-4.9471	3.5160	1.1593	4.0253	-3.4762	-1.1058	-3.0181
-2.8254	-3.3452	4.0723	1.6831	2.3066	-1.7267	-3.1906	-2.9813	-3.6434	-3.4390	-2.9897
-0.5413	-3.8440	-3.5421	-3.5699	-0.8585	0.4972	4.6667	-2.1376	1.8821	-4.4475	-2.9977
2.0459	0.3452	-2.6328	5.0286	0.0968	0.0902	-3.5274	3.0677	4.1941	-3.8691	-3.0076
3.8297	3.4886	5.0435	-5.3804	0.3609	0.1758	-1.0414	1.1182	1.6169	-2.1467	-3.0099
4.2991	3.2883	0.0882	4.3508	0.3813	0.1274	4.5894	-3.5804	-2.9543	0.3095	-3.0140
3.3747	-1.3949	-4.6599	-2.8194	-1.3521	-1.0905	-4.1776	0.2709	-4.6121	2.4728	-3.0121

find the solution of Problem 1.2.

- (1) At first, we find the least-norm solution of Problem 1.1 by iteration method 2.1. Let $\epsilon = 1.0e - 10$, and when $\|\Delta X_k\| < \epsilon$, stop the iteration, then we have that

$$X = \begin{pmatrix} 1.0000 & 0.5000 & 0.3333 & 0.2500 & 0.2000 & 0.1667 & 0.1429 & 0.1250 & 0.1111 & 0.1000 & 0.0909 \\ 0.5000 & 0.3333 & 0.2500 & 0.2000 & 0.1667 & 0.1429 & 0.1250 & 0.1111 & 0.1000 & 0.0909 & 0.0833 \\ 0.3333 & 0.2500 & 0.2000 & 0.1667 & 0.1429 & 0.1250 & 0.1111 & 0.1000 & 0.0909 & 0.0833 & 0.0769 \\ 0.2500 & 0.2000 & 0.1667 & 0.1429 & 0.1250 & 0.1111 & 0.1000 & 0.0909 & 0.0833 & 0.0769 & 0.0714 \\ 0.2000 & 0.1667 & 0.1429 & 0.1250 & 0.1111 & 0.1000 & 0.0909 & 0.0833 & 0.0769 & 0.0714 & 0.0667 \\ 0.1667 & 0.1429 & 0.1250 & 0.1111 & 0.1000 & 0.0909 & 0.0833 & 0.0769 & 0.0714 & 0.0667 & 0.0625 \\ 0.1429 & 0.1250 & 0.1111 & 0.1000 & 0.0909 & 0.0833 & 0.0769 & 0.0714 & 0.0667 & 0.0625 & 0.0588 \\ 0.1250 & 0.1111 & 0.1000 & 0.0909 & 0.0833 & 0.0769 & 0.0714 & 0.0667 & 0.0625 & 0.0588 & 0.0556 \\ 0.1111 & 0.1000 & 0.0909 & 0.0833 & 0.0769 & 0.0714 & 0.0667 & 0.0625 & 0.0588 & 0.0556 & 0.0526 \\ 0.1000 & 0.0909 & 0.0833 & 0.0769 & 0.0714 & 0.0667 & 0.0625 & 0.0588 & 0.0556 & 0.0526 & 0.0500 \\ 0.0909 & 0.0833 & 0.0769 & 0.0714 & 0.0667 & 0.0625 & 0.0588 & 0.0556 & 0.0526 & 0.0500 & 0.0476 \end{pmatrix}$$

where, $\sigma_r / \sigma_1 = \lambda_s / \lambda_1 = 4.4456/81.1530$, $t = 7.2210$, $k = 6756$. In comparison, by using the iteration method from the paper [9], we can get the same result with $t = 28.6520$, $k = 18317$.

Secondly, by using preconditioned iteration method, let $C(A)A = I_{11} - 4 \times (0.001 \times A - I_{11})^3 + 3 \times (0.001 \times A)^2$, denote $C(A)A$ as \tilde{A} , $C(A)C$ as \tilde{C} , solving the equation $\tilde{A}X\tilde{A} = \tilde{C}$, we can obtain the same X with the original method, but where, $\sigma_r / \sigma_1 = \lambda_s / \lambda_1 = 4.1228/5.1832$, $t = 0.1000$, $k = 17$. And under the same situation, by using the iteration method from the paper [9], we find that the iteration is not convergent.

- (2) Denotes the set of all solutions of the matrix equation $AXB = C$ in this example as S_E . In order to find the optimal approximate solution to a given matrix X_0 , let $X^* = X - X_0$, $C^* = C - AX_0B$, and $\epsilon = 1.0e - 10$, by iteration method 2.1, when $\|\Delta X_k\| < \epsilon$, stop the iteration, then we can obtain the least-norm solution \tilde{X}^* of the

consistent matrix equation $AX^*B = C^*$, and so the optimal approximation solution \hat{X} to the given matrix X_0 is that

$$\hat{X} = \tilde{X}^* + X_0 = \begin{pmatrix} 1.0000 & 0.5000 & 0.3333 & 0.2500 & 0.2000 & 0.1667 & 0.1429 & 0.1250 & 0.1111 & 0.1000 & 0.0909 & 0.0833 \\ 0.5000 & 0.3333 & 0.2500 & 0.2000 & 0.1667 & 0.1429 & 0.1250 & 0.1111 & 0.1000 & 0.0909 & 0.0833 & 0.0769 \\ 0.3333 & 0.2500 & 0.2000 & 0.1667 & 0.1429 & 0.1250 & 0.1111 & 0.1000 & 0.0909 & 0.0833 & 0.0769 & 0.0714 \\ 0.2500 & 0.2000 & 0.1667 & 0.1429 & 0.1250 & 0.1111 & 0.1000 & 0.0909 & 0.0833 & 0.0769 & 0.0714 & 0.0667 \\ 0.2000 & 0.1667 & 0.1429 & 0.1250 & 0.1111 & 0.1000 & 0.0909 & 0.0833 & 0.0769 & 0.0714 & 0.0667 & 0.0625 \\ 0.1667 & 0.1429 & 0.1250 & 0.1111 & 0.1000 & 0.0909 & 0.0833 & 0.0769 & 0.0714 & 0.0667 & 0.0625 & 0.0588 \\ 0.1429 & 0.1250 & 0.1111 & 0.1000 & 0.0909 & 0.0833 & 0.0769 & 0.0714 & 0.0667 & 0.0625 & 0.0588 & 0.0556 \\ 0.1250 & 0.1111 & 0.1000 & 0.0909 & 0.0833 & 0.0769 & 0.0714 & 0.0667 & 0.0625 & 0.0588 & 0.0556 & 0.0528 \\ 0.1111 & 0.1000 & 0.0909 & 0.0833 & 0.0769 & 0.0714 & 0.0667 & 0.0625 & 0.0588 & 0.0556 & 0.0528 & 0.0500 \\ 0.1000 & 0.0909 & 0.0833 & 0.0769 & 0.0714 & 0.0667 & 0.0625 & 0.0588 & 0.0556 & 0.0528 & 0.0500 & 0.0476 \\ 0.0909 & 0.0833 & 0.0769 & 0.0714 & 0.0667 & 0.0625 & 0.0588 & 0.0556 & 0.0528 & 0.0500 & 0.0476 & 0.0446 \end{pmatrix}$$

EXAMPLE 2. Let A, B is as the same matrix as of the Example 1,

$$C = \begin{pmatrix} 112.9234 & 101.6547 & 97.0850 & 92.5469 & 85.4087 & 68.6364 & 77.4207 & 82.7639 & 86.6531 & 89.3282 & 90.5845 \\ 102.7098 & 99.8743 & 97.4271 & 93.6838 & 86.7587 & 69.3091 & 78.6322 & 84.3071 & 88.4434 & 91.3066 & 92.6865 \\ 97.5420 & 96.8334 & 95.2690 & 92.0052 & 85.3728 & 67.9033 & 77.2982 & 83.0183 & 87.1934 & 90.0975 & 91.5227 \\ 93.6942 & 93.8054 & 92.6886 & 89.7810 & 83.4736 & 66.4063 & 75.8278 & 81.2426 & 85.3412 & 88.1977 & 89.6139 \\ 86.9268 & 87.2927 & 86.4117 & 83.8490 & 78.2639 & 62.9107 & 71.2892 & 76.3890 & 80.1000 & 82.6749 & 83.9463 \\ 75.4483 & 75.3918 & 74.4059 & 72.1537 & 67.8220 & 57.8860 & 64.0435 & 67.8242 & 70.5189 & 72.3014 & 73.0736 \\ 80.6782 & 80.9804 & 80.1255 & 77.7547 & 72.8791 & 60.9908 & 68.1686 & 72.5765 & 75.7453 & 77.8800 & 78.8556 \\ 85.1314 & 85.6985 & 84.9325 & 82.4481 & 77.1117 & 63.6465 & 71.6630 & 76.5880 & 80.1474 & 82.5711 & 83.7116 \\ 87.5688 & 88.3184 & 87.6244 & 85.0862 & 79.4751 & 64.9886 & 73.5321 & 78.7816 & 82.5879 & 85.1971 & 86.4471 \\ 90.0943 & 91.0130 & 90.3820 & 87.7802 & 81.8769 & 66.3291 & 75.4082 & 80.9873 & 85.0452 & 87.8443 & 89.2067 \\ 90.3920 & 91.3856 & 90.7946 & 88.1901 & 82.2090 & 66.3601 & 75.5981 & 81.2772 & 85.4134 & 88.2738 & 89.6746 \end{pmatrix}$$

In this example, the equation $AXB = C$ is not consistent, by using Iteration method 2.1, we can obtain the least-norm least-squares solution of the equation as

$$X = \begin{pmatrix} 0.1013 & 0.0513 & 0.0369 & 0.0279 & 0.0234 & 0.0032 & 0.0155 & 0.0138 & 0.0135 & 0.0116 & 0.0108 \\ 0.0514 & 0.0347 & 0.0293 & 0.0239 & 0.0229 & -0.0040 & 0.0138 & 0.0123 & 0.0129 & 0.0108 & 0.0102 \\ 0.0353 & 0.0269 & 0.0269 & 0.0237 & 0.0264 & -0.0194 & 0.0128 & 0.0114 & 0.0135 & 0.0108 & 0.0104 \\ 0.0270 & 0.0219 & 0.0241 & 0.0219 & 0.0231 & -0.0207 & 0.0116 & 0.0105 & 0.0128 & 0.0102 & 0.0099 \\ 0.0227 & 0.0194 & 0.0300 & 0.0358 & 0.0839 & -0.1108 & 0.0118 & 0.0075 & 0.0197 & 0.0116 & 0.0118 \\ 0.0074 & 0.0053 & -0.0262 & -0.0341 & -0.0952 & 0.2294 & 0.0003 & 0.0036 & -0.0190 & -0.0058 & -0.0079 \\ 0.0152 & 0.0134 & 0.0136 & 0.0121 & 0.0123 & -0.0018 & 0.0086 & 0.0081 & 0.0085 & 0.0074 & 0.0072 \\ 0.0136 & 0.0122 & 0.0130 & 0.0116 & 0.0120 & -0.0042 & 0.0082 & 0.0077 & 0.0084 & 0.0072 & 0.0070 \\ 0.0122 & 0.0111 & 0.0121 & 0.0108 & 0.0113 & -0.0047 & 0.0077 & 0.0073 & 0.0080 & 0.0069 & 0.0067 \\ 0.0112 & 0.0103 & 0.0119 & 0.0107 & 0.0116 & -0.0076 & 0.0074 & 0.0071 & 0.0080 & 0.0068 & 0.0067 \\ 0.0103 & 0.0096 & 0.0111 & 0.0100 & 0.0108 & -0.0075 & 0.0071 & 0.0068 & 0.0077 & 0.0065 & 0.0064 \end{pmatrix}$$

where, $\sigma_r / \sigma_1 = \lambda_s / \lambda_1 = 4.4456/81.1530, t = 7.6710, k = 7272$.

By using preconditioned method as Example 1, we can obtain the same result, but where, $\sigma_r / \sigma_1 = \lambda_s / \lambda_1 = 4.1228/5.1832, t = 0.0200, k = 17$.

From the above two examples, we see that the converge rate of the iteration method is surely related to the singular value of the matrices A, B , and by using preconditioning methods which increase the ratio $(\sigma_r \lambda_s) / (\sigma_1 \lambda_1)$, we can obtain a faster iteration rate.

REFERENCES

- [1] A. Bjerhammer, 'Rectangular reciprocal matrices with special reference to geodetic calculations', *Kung. Tekn. Hogsk. Handl. Stockhol* **45** (1951).
- [2] K-W.E. Chu, 'Symmetric solutions of linear matrix equations by matrix decompositions', *Linear Algebra Appl.* **119** (1989), 35-50.
- [3] H. Dai, 'On the symmetric solutions of linear matrix equations', *Linear Algebra Appl.* **131** (1990), 1-7.
- [4] F.J.H Don, 'On the symmetric solution of a linear matrix equation', *Linear Algebra Appl.* **93** (1987), 1-7.

- [5] G. Li, *Common matrix theories and methods* (Shanghai Science and Technology Press, Shanghai 1983).
- [6] J.R. Magnus, 'L-structured matrices and linear matrix equation', *Linear and Multilinear Algebra* **14** (1983), 67–88.
- [7] S.K. Mitra, 'Common solutions to a pair of linear matrix equations $A_1XB_1 = C_1$ and $A_2XB_2 = C_2$ ', *Proc. Cambridge Philos. Soc.* **74** (1973), 213–216.
- [8] G.R. Morris and P.L. Odell, 'Common solutions for n matrix equations with applications', *J. Assoc. Comput. Mach.* **15** (1968), 272–274..
- [9] Y.X. Peng and X.Y. Hu, 'An iteration method for the symmetric solutions and optimal approximation solution of the matrix equation $AXB = C$ ', *Appl. Math. Comput.* **160** (2005), 163–167.
- [10] R. Penrose, 'A generalized inverse for matrices', *Proc. Cambridge Philos. Soc.* **5** (1955), 406–413.
- [11] X.C. He and W.Y. Sun, *Introduction to generalized inverse of matrices* (Jiangsu Science and Technology Press, Nanjing, 1991).

College of Mathematics and Econometrics
Hunan University
Changsha 410082
Peoples Republic of China
e-mail: khguo@126.com