

Bioinformatic analysis of eosinophil activity and its implications for model and target species

Review

Cite this article: Jenvey C.J, Alenizi D, Almasi F, Cairns C, Holmes A, Sloan S, Stear M.J (2020). Bioinformatic analysis of eosinophil activity and its implications for model and target species. *Parasitology* **147**, 393–400. <https://doi.org/10.1017/S0031182019001768>

Received: 31 October 2019
Revised: 8 December 2019
Accepted: 9 December 2019
First published online: 16 December 2019

Key words:

Bioinformatics; CCR3; eosinophils; EPX; Fc α RI; gastrointestinal nematodes; PRG3; ruminants

Author for correspondence:

C.J. Jenvey, E-mail: C.Jenvey@latrobe.edu.au

C.J. Jenvey , D. Alenizi, F. Almasi, C. Cairns, A. Holmes, S. Sloan and M.J. Stear

Department of Animal, Plant and Soil Sciences, Agribio Centre for Agribioscience, School of Life Sciences, La Trobe University, Bundoora, Victoria, Australia

Abstract

Eosinophils are important immune cells that have been implicated in resistance to gastrointestinal nematode (GIN) infections in both naturally and experimentally infected sheep. Proteins of particular importance appear to be IgA-Fc alpha receptor (Fc α RI), C-C chemokine receptor type 3 (CCR3), proteoglycan 3 (PRG3, major basic protein 2) and EPX (eosinophil peroxidase). We used known human nucleotide sequences to search the ruminant genomes, followed by translation to protein and sequence alignments to visualize differences between sequences and species. Where a sequence was retrieved for cow, but not for sheep and goat, this was used additionally as a reference sequence. In this review, we show that eosinophil function varies among host species. Consequently, investigations into the mechanisms of ruminant immune responses to GIN should be conducted using the natural host. Specifically, we address differences in protein sequence and structure for eosinophil proteins.

Introduction

Host immune responses to gastrointestinal nematodes (GINs) are dominated by a Th2 immune response; involving antibodies and immune cells, such as immunoglobulin A (IgA), IgE, mast cells and eosinophils. In particular, ruminants naturally and experimentally infected with GIN demonstrate an increase in blood and tissue eosinophilia, implying that eosinophils may be an important mediator of host immune responses to GIN. However, both phenotypic and bioinformatic evidence suggest that eosinophil activity against GIN may differ between hosts (Urban *et al.*, 1991; Henderson and Stear, 2006). Bioinformatic analyses on eosinophil-associated proteins were used to explore whether differences in resistance to GIN among species were genetic in origin. Specifically, we addressed differences in protein sequence and structure for eosinophil proteins. These proteins included IgA and its receptor, Fc α RI, interleukin (IL)-5 and its receptor, IL-5R α , eotaxin and its receptor, CCR3, major basic protein (MBP, PRG3) and eosinophil peroxidase (EPX). We used known human nucleotide sequences to search the ruminant genomes (*Bos taurus*, cow; *Ovis aries*, sheep; *Capra hircus*, goat), retrieved sequences (Supplementary Table 1), followed by translation to protein and sequence alignments to visualize differences between sequences and species.

Eosinophils and GIN infections

Eosinophils are a sub-type of granulocyte, along with mast cells, neutrophils and basophils. Following proliferation of eosinophil precursors from the bone marrow, eosinophils traffic to sites of infection and are activated. Once activated, eosinophils undergo degranulation, releasing cytotoxic proteins from secondary granules to protect the host against foreign pathogens. Eosinophils are also involved in immune homeostasis and immunity (Rothenberg and Hogan, 2006; Weller and Spencer, 2017). Eosinophils are found in both blood and tissue, however the gastrointestinal tract contains the largest reservoir of eosinophils in the body (Zuo and Rothenberg, 2007) and only tissue eosinophils degranulate (Blanchard and Rothenberg, 2009). The ability of eosinophils to defend the host against parasitic helminths is suggested by the ability of eosinophils to mediate antibody- (or complement-) dependent cellular cytotoxicity (ADCC) *in vitro* and *in vivo* (Giacomin *et al.*, 2008; Huang *et al.*, 2015), increased numbers of eosinophils during helminth infections, as well as degranulation in close proximity to helminths *in vivo* (Rothenberg and Hogan, 2006).

Eosinophils are not only important for GIN infections of humans and mice, but also of ruminants. In particular, GIN infections ravage sheep and goat populations in temperate regions of the world, and in Australia can cost sheep producers up to \$500 million per year largely in lost productivity (Lane *et al.*, 2015). In particular, *Teladorsagia circumcincta*, *Haemonchus contortus* and *Trichostrongylus colubriformis* are the dominant GIN-infecting small ruminants (Roeder *et al.*, 2013). The typical immune response to GIN is dominated by Th2 immune responses, namely the production of antibodies such as IgG, IgE and IgA, as well as involvement of mast cells and eosinophils, the details of which have previously been covered by a number of reviews (Maizels and Yazdanbakhsh, 2003; Anthony *et al.*,

© The Author(s), 2019. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.




Fig. 1. The goat IgA receptor, *FcαRI*, may be dysfunctional in goats. Homology modelling of human, P24071.1 (A); sheep, XM_027976143.1 (B) and goat, XM_018059931.1 (C) IgA receptors revealed that human and sheep receptors have a similar conformation, however, the goat receptor has a C-terminus that is bent away from the main body of the protein and lacked an alpha helix within the transmembrane domain (blue).

2007; McRae *et al.*, 2015; Motran *et al.*, 2018). Previous research using experimentally and naturally infected sheep have indicated that eosinophils may play an important role in resistance to infection. Investigations into sheep immune responses to *T. circumcincta* (Gruner *et al.*, 1994; Stear *et al.*, 1995, 2002; Henderson and Stear, 2006; Beraldi *et al.*, 2008), *H. contortus* (Rainbird *et al.*, 1998; Gill *et al.*, 2000; Balic *et al.*, 2006; Terefe *et al.*, 2007, 2009) and *T. colubriformis* (Dawkins *et al.*, 1989; Rothwell *et al.*, 1993; Amarante *et al.*, 2007) have all demonstrated increases in eosinophils in resistant animals, resistant breeds and/or in sheep selectively bred for resistance. In addition, differences in numbers of eosinophils and susceptibility to infection have also been observed in goats (Bambou *et al.*, 2013; Basripuzi *et al.*, 2018).

Such findings imply that eosinophils are important cells in ruminant responses to GIN infections. Phenotypic and bioinformatic evidence suggests that there are differences in immune responses to GIN between species. A recent review by Weller and Spencer (2017) discussed a number of unresolved issues when comparing mouse and human eosinophils, namely whether the formation and secretion of eosinophil cytokines is regulated by common mechanisms. In addition, a review by Behm and Ovington (2000) highlighted that IL-5 and eosinophils have different impacts on different helminth infections. Conversely, a review by Meeusen and Balic (2000) suggests that the presence of IL-5 independent eosinophil populations within tissue and peripheral blood may play a role in unnatural nematode-mouse models by increasing resistance to primary infections, and enhancing the development of specific immunity upon subsequent infections. Ultimately, although a number of *in vitro* studies investigating the mechanisms by which eosinophils cause helminth death have been demonstrated, it is not yet clear whether these same mechanisms also occur *in vivo* (Motran *et al.*, 2018). Recent updates to the human, mouse, cow, sheep and goat genomes have provided insights into the possible mechanisms of eosinophil function, which can be used to direct functional studies into the mechanisms of eosinophils during GIN infections of ruminants.

The IgA receptor, *FcαRI*, may be dysfunctional in goats

Immunoglobulin A (IgA) is an antibody that plays a crucial role in the immune function of mucus membranes. TGF- β , together with IL-5, is responsible for class-switching of B lymphocytes into IgA-producing plasma cells (Coffman *et al.*, 1989; Sonoda *et al.*, 1989). This local production of IgA is termed secretory IgA and is the predominant form of IgA in mucosal secretions (van Egmond *et al.*, 2001; Bakema and van Egmond, 2011). Considering the importance of IgA in the protection of mucus membranes, it is unsurprising that studies investigating immune responses to parasitic infections have found associations between IgA and parasite-induced eosinophilia (Muraki *et al.*, 2011). IgA has been demonstrated to be associated with nematode fecundity and peripheral eosinophils, and therefore resistance to helminth

infection in sheep (Gill *et al.*, 1993; Henderson and Stear, 2006; Halliday *et al.*, 2007; Hernández *et al.*, 2016; Fairlie-Clarke *et al.*, 2019). The lack of an effective IgA response has also been implicated in increased susceptibility to parasite infection in goats (Basripuzi *et al.*, 2018). Additionally, *in vitro* studies have indicated that parasite-induced eosinophil cytolysis may be dependent upon IgA binding to its receptor (Ueki *et al.*, 2013).

IgA exists as a monomer of two identical heavy chains and two identical light chains, but can form a dimer (secretory IgA) stabilized by disulphide bonds and a joining (J) chain (Woof and Russell, 2011). In ruminants, most IgA in serum derives from mucosal surfaces and is largely dimeric (Scicchitano *et al.*, 1986). The *FcαRI* (CD89) is a transmembrane receptor and due to heavy glycosylation, eosinophil *FcαRI* is heavier (70–100 kDa) compared to macrophage and neutrophil *FcαRI* (55–75 kDa) (van Egmond *et al.*, 2001). *FcαRI* comprises a ligand-binding alpha chain, which consists of two extracellular Ig-like domains, a transmembrane domain, and a short cytoplasmic tail. Due to the lack of signalling motifs, *FcαRI* must associate with the *FcRγ* chain for signalling and function (van Egmond *et al.*, 2001).

Molecular modelling of the eosinophil IgA receptor indicates that it may be dysfunctional in goats (Basripuzi *et al.*, 2018). Structural modelling of the IgA receptor indicated that the sheep and human receptors had similar structures. However, the goat receptor had a different conformation, with the C-terminus being bent away from the main body of the protein and lacked an alpha helix within the transmembrane domain (approximately 20 amino acids) (Fig. 1). Although the sheep and goat sequences showed high sequence identity (96%), the sheep and goat sequences showed only 55 and 54% sequence identity with the human protein. Additionally, a difference of only three amino acids was observed within the binding domain (EC1) and no differences were observed within the transmembrane domain. This seems to indicate that the loss of secondary structure within the transmembrane domain is most likely a consequence of sequence differences outside of this domain. As yet, no mouse homologue of *FcαRI* has been identified (van Egmond *et al.*, 2001; Decot *et al.*, 2005), thus if eosinophils and IgA interact to control nematodes through ADCC, mice may not be appropriate models for studying eosinophil activity against nematodes infecting humans and ruminants.

Differences in eosinophil responsiveness are not due to IL-5 and IL-5R α

IL-5 is a growth factor and chemoattractant of eosinophils and is involved in the recruitment, activation, degranulation and survival of eosinophils (Lopez *et al.*, 1988; Horie *et al.*, 1996; Fulkerson *et al.*, 2014; Sippel *et al.*, 2018). IL-5 is an important cytokine in the differentiation and activation of anti-parasitic eosinophil responses and has been specifically targeted in mice to demonstrate the *in vivo* role of eosinophils in helminth infections, as




Fig. 2. Eotaxin receptor, CCR3, may be dysfunctional in goats. Homology modelling revealed that the two sheep sequences [Q9N0M0 (B), W5PXW1 (C)] were identical except for two substitutions and all substitutions were at the C-terminal end of the sequence. The two goat sequences [JO419941.2 (A), LOC5316646 (D)] contained two substitutions compared to the sheep sequences. In addition, the goat genome sequence contained a frameshift deletion, corresponding to amino acid 330 (pink). Substitutions are colour coded by the following: aa 354 (blue), aa 356 (red), aa 357 (green) and aa 358 (orange).

discussed in previous reviews (Huang and Appleton, 2016; Meeusen and Balic, 2000). However, the reagents and knock-out models used in these rodent studies are generally not available or suitable for large animal experimentation. Several studies have shown that mice vaccinated with recombinant cytokines can induce autoantibodies that specifically cancel out the activity of the native cytokine *in vivo* (Dalum *et al.*, 1999; Hertz, *et al.*, 2001; Richard *et al.*, 2000). This approach may provide an alternative for the *in vivo* study of immune responses in large animals.

IL-5 acts on target cells by binding to its receptor, IL-5R, which consists of an α and a β subunit, however the α subunit is specific to IL-5R only (McBrien and Menzies-Gow, 2017). IL-5 is a dimeric glycoprotein with a four-helix bundle motif. In complex, IL-5 forms a homodimer which is sandwiched by the IL-5R α . Binding of IL-5 to the receptor alpha subunit results in recruitment of the β subunit to the receptor (Tavernier *et al.*, 1991; Kusano *et al.*, 2012). The human and mouse cDNA code for proteins of 134 and 133 amino acids in length, respectively, and have 70% amino acid sequence identity (Yamaguchi, 1994).

Molecular modelling indicates that it is unlikely that IL-5 and its receptor are responsible for differences in eosinophil responses between sheep and goats. For IL-5, the predicted protein sequences were identical between sheep and goat, while for IL-5R α , the predicted protein sequences differed in 6 out of the 432 amino acids. All of these differences occurred in the first 42 amino acids of the protein, which included within the 15–34 amino acids that aligned with the signal peptide for human IL-5R α , as well as within a region lacking secondary structure. This indicates that the observed sequence differences are unlikely to result in a dysfunctional IL-5 and IL-5R.

The eotaxin receptor, CCR3, contains a frameshift mutation in goats

Eotaxin is a chemoattractant cytokine which is important for promoting eosinophil recruitment and degranulation (Garcia-Zepeda *et al.*, 1996; Davoine and Lacy, 2014), and has been shown to be important for eosinophil recruitment during helminth infections (Rothenberg *et al.*, 1997; Mochizuki *et al.*, 1998; Ruth *et al.*, 1998; Culley *et al.*, 2000; Simons *et al.*, 2005). Eotaxin belongs to the CC chemokine family, which is distinguished by two cysteines immediately adjacent to the N terminus. Eotaxin has been determined to be in equilibrium between a monomer and a dimer at near physiological pH, however, functional eotaxin is present as a monomer (Crump *et al.*, 1998). It exhibits a chemokine-like fold

consisting of three anti-parallel β -strands with an overlying α -helix (Crump *et al.*, 1998). There are three molecules of Eotaxin, CCL11 (Eotaxin-1), CCL24 (Eotaxin-2) and CCL26 (Eotaxin-3); however, Eotaxin-1 is the dominant isoform. The highest levels of Eotaxin-1 are found in the GI system and can be produced by a variety of cells (Kitaura *et al.*, 1996; Ying *et al.*, 1999). Eotaxin-1 is important for the release of eosinophil precursors from the bone marrow (Palframan *et al.*, 1998) and is activated by Th2 cytokines (Mochizuki *et al.*, 1998) and inhibited by Th1 cytokines (Miyamasu *et al.*, 1999; Fukuda *et al.*, 2002). Eotaxin-2 is synthesized and released by mucosal epithelial cells and macrophages, while Eotaxin-3 is produced by epithelial and endothelial cells (Kitaura *et al.*, 1999; Shinkai *et al.*, 1999; Dulkys *et al.*, 2001). Eotaxin-2 and Eotaxin-3 can also recruit eosinophils, but at later stages of infection (>24 h) (Ying *et al.*, 1999; Rosenwasser *et al.*, 2003; Kalomenidis *et al.*, 2005; Schratl *et al.*, 2006).

All Eotaxin isoforms are associated with a single receptor, CCR3, however the receptor binds to the different isoforms with different affinities (Kitaura *et al.*, 1999). CCR3 is very abundant in eosinophils (approximately $40\text{--}400 \times 10^3$ receptors per cell), but is also expressed at lower levels in basophils, mast cells and a subset of Th2 lymphocytes (Sallusto *et al.*, 1997; Ugucioni *et al.*, 1997; Romagnani *et al.*, 1999). CCR3 also binds to other non-eosinophil selective CC chemokines, but with lower affinity compared to Eotaxin-1 (Ponath *et al.*, 1996; Baggiolini *et al.*, 1997; Sabroe *et al.*, 1999). CCR3 is a G-protein-coupled receptor of 335 amino acids in length and shares 63 and 51% sequence homology with CCR1 and CCR2, respectively (Daugherty *et al.*, 1996). The CCR3 gene codes for four cysteine residues, one in each of the extracellular domains, and a serine/threonine-rich cytoplasmic tail, all of which are highly conserved features of chemokine receptors (Ponath *et al.*, 1996). Uniquely, CCR3 contains a cluster of negatively charged amino acids distal to the transmembrane helix IV in the second extracellular loop (Daugherty *et al.*, 1996).

Molecular modelling of the Eotaxin receptor, CCR3, indicates that it may be dysfunctional in goats. The sheep and goat CCR3 protein sequences differed by only two amino acids. The two sheep protein sequences were identical except for two substitutions. The two goat sequences were derived from the goat genome sequence and the other from mRNA extracted from the liver of an Osmanabadi goat. The goat genome sequence contained a frameshift deletion on chromosome 22 at 52,155,650, which corresponded to amino acid 330 of the CCR3 protein (Fig. 2). This deletion was not present in the mRNA; therefore, it is possible that this deletion is merely a




Fig. 3. Major basic protein 2 is the only MBP molecule in ruminants. Homology modelling of MBP-1 and MBP-2 C-type lectin fold revealed high structural similarities between human MBP-2 and ruminant MBP-2 molecules, however, theoretical isoelectric point (pI) computation suggests that ruminant MBP-2 molecules may function similarly to human MBP-1. Amino acids used to calculate theoretical pI are as follows: aspartic acid (green), glutamic acid (orange), histidine (yellow), cysteine (red), tyrosine (pink), lysine (aqua) and arginine (blue). Molecules are presented in the order of percentage of sequence identity to human MBP-1, CR450311.1 (A) (highest to lowest): human MBP-2, NM_006093.4 (B); sheep XM_027979083.1 (C); sheep XM_027979084.1 (D); goat XM_018058941.1 (E); cow NM_001098471.1 (F); sheep XM_027979081.1 (G) and goat XM_018058942 (H).

Table 1. Theoretical isoelectric point (pI) of sequences for MBP molecules from humans, cows, sheep and goats

MBP sequence	Theoretical pI
Human CR450311.1	6.23
Human NM_006093.4	4.69
Cow NM_001098471.1	8.00
Sheep XM_02797908.1	6.42
Sheep XM_027979083.1	6.22
Sheep XM_027979084.1	7.96
Goat XM_018058942	6.43
Goat XM_018058941.1	8.59

Amino acid sequences were entered into the ExPASy Bioinformatics Resource Portal Compute pI/MW Tool (https://web.expasy.org/compute_pi/) for estimating average theoretical pI based upon the pK values of amino acids described in Bjellqvist *et al.* (1993) and Bjellqvist *et al.* (1994). The pK values in these studies were defined by examining polypeptide migration between pH 4.5 and 7.3; therefore, predictions for proteins outside of this pH range (cow NM_001098471.1, sheep XM_027979084.1, goat XM_018058941.1) may not be accurate.

sequencing artefact. Additionally, both goat sequences contain two amino acid substitutions in relation to the sheep sequences. However, it is unlikely that these substitutions would be the cause of any dysfunction, as the same amino acids are present in the same positions in the human CCR3 protein sequence. In any case, more research is necessary to establish if the potential frameshift mutation is polymorphic in goats and whether or not it may contribute to differences in eosinophil responsiveness between relatively resistant and susceptible goat breeds.

MBP-2 is the only MBP molecule present in ruminants

MBP-1 is an abundant granule protein of human eosinophils. Its homologue, MBP-2, is unique to eosinophils (Acharya and Ackerman, 2014). MBP is localized within the crystalline core of the eosinophil and is an important mediator of eosinophil function (Gleich and Adolphson, 1986; Kita, 2011). MBP is highly basic, which results in the binding of MBP to cell membranes. Cytotoxic mechanisms of MBP involve surface interchange to increase cell membrane permeability and interrupting tissue enzyme activity (Ackerman *et al.*, 1985; Gleich and Adolphson, 1986; Swaminathan *et al.*, 2001). MBP has been demonstrated to

be toxic against *Schistosoma mansoni* by disrupting the cell membrane *via* the binding of heparin (Butterworth *et al.*, 1979), as well as being important in the control of *Litomosoides sigmodontis* in mice (Specht *et al.*, 2006). Structurally, MBP is most like C-type lectins, except that it lacks a calcium binding site, and instead binds selectively to heparin and heparin sulphate, glycosaminoglycans and chondroitin sulphate B (Swaminathan *et al.*, 2005; Wagner *et al.*, 2007). Human MBP-1 is 222 amino acids long, consisting of a signalling peptide, pro-peptide and two chains (Swaminathan *et al.*, 2001). It has been suggested that the pro-peptide protects the eosinophil from MBP during transport from the Golgi apparatus to the crystalline core by masking the mature domain, as well as by blocking glycosylated binding sites to inactivate the protein (Swaminathan *et al.*, 2001). The mature domain is highly basic and is the region where carbohydrate recognition occurs. There is a 66% amino acid sequence identity between MBP-1 and MBP-2, with MBP-2 being less basic. MBP-2 contains 10 cysteine residues, eight of which are conserved in MBP-1, including those cysteines that are involved in disulphide bridges. The conservation of these disulphide bridges is consistent with other C-type lectins and is thought to be important for tertiary structure and function (Wagner *et al.*, 2007).

Sequence searches of MBP indicate that only MBP-2 is detectable in ruminant genomes; it may function similarly to MBP-1. A total of three sequences for sheep (XM_027979083.1, XM_027979084.1 and XM_02797908.1), two sequences for goats (XM_018058941.1 and XM_018058942) and one sequence for cows (NM_001098471.1) were retrieved. Of the two human MBP sequences, the ruminant sequences were most similar to MBP-2 (55–58% homology). Homology between the sheep sequences was between 72 and 79%, while homology between the goat sequences was 79%. The highest homology was observed between sheep sequence XM_027979084.1 and goat sequence XM_018058941.1, with 95%, which implies a recent divergence and that sheep and goats may have multiple MBP loci. Typical features of C-type lectins and MBP were conserved in the ruminant sequences, including the C-type lectin fold, disulphide-bonded cysteines, and the heparin binding sites. Despite higher homology between human MBP-2 and the ruminant sequences (Fig. 3), theoretical isoelectric point (pI) calculations for the ruminant sequences were most similar to the pI for human MBP-1, with the pI of all ruminant sequences being within 26% of the pI human MBP-1, compared to within 61% of the pI of MBP-2 (Table 1). It is possible that in the absence of




Fig. 4. Eosinophil peroxidase may be dysfunctional in goats. Homology modelling EPX heavy chain revealed goat EPX does not contain a tyrosine involved which is involved in post-translational modification of eosinophil granule proteins during eosinophil maturation in human eosinophils. The nitrated tyrosine (green) in the human, NM_000502.6 (A); cow, XM_024980582.1 (B) and sheep, contig. AMGL01017333.1 (C) sequences have been replaced by a cysteine (red) in the goat, contig. LWLT0100022.1 (D) sequence.

MBP-1, ruminant MBP-2 may function as MBP-1, being more cytotoxic and abundant. Functional assays are required to determine if this hypothesis is in fact the case.

Goat EPX lacks a nitrosylated tyrosine

Similar to MBP-2, eosinophil peroxidase (EPX) is unique to eosinophils and is the most abundant cationic protein within the matrix of the specific granule of the eosinophil (Acharya and Ackerman, 2014). Human EPX is 715 AA long, located on chromosome 17. It is structurally similar to myeloperoxidase (MPO), which is present in neutrophil-specific granules (Loughran *et al.*, 2008). EPX uses hydrogen peroxide to produce toxic reactive oxygen species, such as hypohalous acids, and is capable of killing parasites, including *S. mansoni* (Auriault *et al.*, 1982), *Toxoplasma gondii* (Locksley *et al.*, 1982) and *L. sigmodontis* (Specht *et al.*, 2006). In addition to eosinophils, mast cells also play a role in parasitic infections. High concentrations of EPX have been demonstrated to result in mast cell lysis. Mast cell lysis is followed by the binding of EPX to mast cell granules to form a complex, which results in the retention of secretory activity on mast cells (Henderson *et al.*, 1980). In addition, a study by Metzler *et al.* (2011) demonstrated donors that were deficient in MPO failed to form neutrophil extracellular traps (NET), indicating that MPO is essential for NET formation. Based on this evidence, it is possible that EPX could be involved in eosinophil extracellular trap (EET) formation, which would implicate eosinophils in the direct control of parasitic infections.

EPX is structurally distinct from the other granule proteins, being a two-chain (55-kDa heavy chain and 12.5-kDa light chain) haemoprotein, although EPX is highly cationic, much like MBP-1 and eosinophil cationic protein. A feature of EPX is that it post-translationally modifies itself *via* the nitrosylation of a specific tyrosine residue (Tyr-488) during synthesis and packaging of the granule proteins into the developing eosinophil (Ulrich *et al.*, 2008). Tyr-488 has also been shown to be surface exposed, which may assist in the production of reactive oxygen species by EPX. Other granule proteins are also nitrosylated by EPX, but it is unclear whether this is important in protection of the host against helminths.

Molecular modelling of EPX indicates that this protein may be dysfunctional in goats. Based on searches using the human (NM_000502.6) and bovine (XM_024980582.1) EPX sequences, no annotated sequences were retrieved from either the sheep or goat genomes. However, short sequences that matched the reference, but not annotated with an associated gene, were retrieved and re-aligned to the reference. The sheep sequence contained one single-nucleotide polymorphism (SNP), while the goat sequence contained two SNP, one of which was in the same location as in the sheep sequence. The sheep and goat sequences were

90.3% homologous, with homology of sheep and goat to the cow sequence being 84.9 and 82.9%, respectively. The cow, sheep and goat sequences all contained MPO-like protein domains, all of which were conserved except for single-residue substitutions within each domain. In addition to the MPO-like domains, the cow and sheep sequences also contained a tryptic peptide fragment, which was conserved in all ruminant sequences except for a single substitution (alignment to human sequence; Arg⁴⁸³ to His⁴⁸³). Of note, the nitrosylated tyrosine was not conserved in the goat sequence and was substituted for a cysteine (Fig. 4). The nitrosylated tyrosine is important for EPX-mediated activities, including the post-translational nitration of eosinophil secondary granule proteins, which in turn, may influence inflammatory responses. The importance of a nitrosylated tyrosine in GIN infections of ruminants has not yet been established; however, the absence of this residue may be responsible for the relative susceptibility of goats to GIN infection. Functional studies are required to determine whether the SNP identified in the sheep and goat sequences may be sequencing error rather than true SNP, as well as to confirm the importance of the nitrosylated tyrosine in resistance and susceptibility to GIN infection.

Directions for future research

Phenotypic and bioinformatic analyses, in combination, are valuable in assessing differences in immune responses between species in order to better direct the design of functional studies. Some eosinophil proteins may be at least partly responsible for the susceptibility of certain ruminant species to GINs. In particular, sequence variation in the IgA receptor, Fc α RI, the Eotaxin receptor, CCR3 and EPX may contribute to the susceptibility of goats to GIN infection. The different conformation of goat Fc α RI, as compared to sheep and human, indicates that it is unlikely that it would be fully effective in antigen-dependent cellular cytotoxicity processes, which have been shown to be important for some helminth infections (Huang *et al.*, 2015). In addition, goat CCR3 may contain a frameshift mutation, which would in turn affect the role of Eotaxin-1 in eosinophil recruitment and degranulation. Goat EPX lacks a nitrosylated tyrosine. In humans, the nitrosylated tyrosine in EPX is important for the post-translational modification of EDGPs during eosinophil maturation. Goat EPX may not be involved in this process. Finally, although MBP-1 is the most abundant and important cationic protein in human eosinophils, it has no orthologue in ruminant genomes. MBP-2 appears to be the only MBP molecule present in ruminants and it may function similarly to human MBP-1. Future research should focus on functional studies to confirm these findings and should involve the extraction of these proteins from the host of interest, followed by sequencing, crystallography

and *in vitro* assays to assess the overall importance of these proteins to the susceptibility of ruminants to GIN.

Perhaps the clearest finding of this review is that bioinformatic analyses indicate that the function of specific eosinophil proteins may vary among host species. Therefore, functional studies of eosinophil activity should be performed in the target species. Extrapolations from one species to another need to be interpreted with great care.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0031182019001768>

Conflict of interest. None.

References

- Acharya KR and Ackerman SJ (2014) Eosinophil granule proteins: form and function. *The Journal of Biological Chemistry* **289**, 17406–17415.
- Ackerman SJ, Gleich GJ, Loegering DA, Richardson BA and Butterworth AE (1985) Comparative toxicity of purified human eosinophil granule cationic proteins for schistosomula of *Schistosoma mansoni*. *The American Journal of Tropical Medicine and Hygiene* **34**, 735–745.
- Amarante AFT, Rocha RA and Bricarello PA (2007) Relationship of intestinal histology with the resistance to *Trichostrongylus colubriformis* infection in three breeds of sheep. *Pesquisa Veterinária Brasileira* **27**, 43–48.
- Anthony RM, Rutitzky LI, Urban JF, Stadecker MJ and Gause WC (2007) Protective immune mechanisms in helminth infection. *Nature Reviews Immunology* **7**, 975–987.
- Auriault C, Capron M and Capron A (1982) Activation of rat and human eosinophils by soluble factor(s) released by *Schistosoma mansoni* schistosomula. *Cellular Immunology* **66**, 59–69.
- Baggiolini M, Dewald B and Moser B (1997) Human chemokines: an update. *Annual Review of Immunology* **15**, 675–705.
- Bakema JE and van Egmond M (2011) The human immunoglobulin A Fc receptor FcαRI: a multifaceted regulator of mucosal immunity. *Mucosal Immunology* **4**, 612.
- Balic A, Cunningham CP and Meeusen ENT (2006) Eosinophil interactions with *Haemonchus contortus* larvae in the ovine gastrointestinal tract. *Parasitology* **28**, 107–115.
- Bambou JC, Larcher T, Cei W, Dumoulin PJ and Mandonnet N (2013) Effect of experimental infection with *Haemonchus contortus* on parasitological and local cellular responses in resistant and susceptible young Creole goats. *BioMed Research International* **2013**, 9.
- Basripuzi NH, Salisi MS, Isa NMM, Busin V, Cairns C, Jenvey C and Stear MJ (2018) Boer goats appear to lack a functional IgA and eosinophil response against natural nematode infection. *Veterinary Parasitology* **264**, 18–25.
- Behm CA and Ovington KS (2000) The role of eosinophils in parasitic helminth infections: insights from genetically modified mice. *Parasitology Today* **16**, 202–209.
- Beraldi D, Craig BH, Bishop SC, Hopkins J and Pemberton JM (2008) Phenotypic analysis of host–parasite interactions in lambs infected with *Teladorsagia circumcincta*. *International Journal for Parasitology* **38**, 1567–1577.
- Bjellqvist B, Hughes GJ, Pasquali C, Paquet N, Ravier F, Sanchez J-C, Frutiger S and Hochstrasser D (1993) The focusing positions of polypeptides in immobilized pH gradients can be predicted from their amino acid sequences. *Electrophoresis* **14**, 1023–1031.
- Bjellqvist B, Basse B, Olsen E and Celis JE (1994) Reference points for comparisons of two-dimensional maps of proteins from different human cell types defined in a pH scale where isoelectric points correlate with polypeptide compositions. *Electrophoresis* **15**, 529–539.
- Blanchard C and Rothenberg ME (2009) Chapter 3 Biology of the eosinophil. In Alt FW (ed.), *Advances in Immunology*, vol. 101. San Diego: Academic Press, pp. 81–121.
- Butterworth AE, Wassom DL, Gleich GJ, Loegering DA and David JR (1979) Damage to *Schistosomula* of *Schistosoma mansoni* induced directly by eosinophil major basic protein. *The Journal of Immunology* **122**, 221–229.
- Coffman RL, Lebnan DA and Shrader B (1989) Transforming growth factor beta specifically enhances IgA production by lipopolysaccharide-stimulated murine B lymphocytes. *The Journal of Experimental Medicine* **170**, 1039–1044.
- Crump MP, Rajarathnam K, Kim K-S, Clark-Lewis I and Sykes BD (1998) Solution structure of Eotaxin, a chemokine that selectively recruits eosinophils in allergic inflammation. *The Journal of Biological Chemistry* **273**, 22471–22479.
- Culley FJ, Brown A, Conroy DM, Sabroe I, Pritchard DI and Williams TJ (2000) Eotaxin is specifically cleaved by hookworm metalloproteases preventing its action *in vitro* and *in vivo*. *The Journal of Immunology* **165**, 6447–6453.
- Dalum I, Butler DM, Jensen MR, Hindersson P, Steinaa L, Waterston AM, Grell SN, Feldmann M, Elsner HI and Mouritsen S (1999) Therapeutic antibodies elicited by immunization against TNF-α. *Nature Biotechnology* **17**, 666–669.
- Daugherty BL, Siciliano SJ, DeMartino JA, Malkowitz L, Sirotina A and Springer MS (1996) Cloning, expression, and characterization of the human eosinophil eotaxin receptor. *The Journal of Experimental Medicine* **183**, 2349–2354.
- Davoine F and Lacy P (2014) Eosinophil cytokines, chemokines, and growth factors: emerging roles in immunity. *Frontiers in Immunology* **5**, 570.
- Dawkins H, Windon R and Eagleson G (1989) Eosinophil responsiveness in sheep selected for high and low responsiveness to *Trichostrongylus colubriformis*. *International Journal for Parasitology* **19**, 199–205.
- Decot V, Woerly G, Loyens M, Loiseau S, Quatannens B, Capron M and Dombrowicz D (2005) Heterogeneity of expression of IgA receptors by human, mouse, and rat eosinophils. *The Journal of Immunology* **174**, 628–635.
- Dulkys Y, Schramm G, Kimmig D, Knöß S, Weyergraf A, Kapp A and Elsner J (2001) Detection of mRNA for Eotaxin-2 and Eotaxin-3 in human dermal fibroblasts and their distinct activation profile on human eosinophils. *Journal of Investigative Dermatology* **116**, 498–505.
- Fairlie-Clarke K, Kaseja K, Sotomaior C, Brady N, Moore K and Stear M (2019) Salivary IgA: a biomarker for resistance to *Teladorsagia circumcincta* and a new estimated breeding value. *Veterinary Parasitology* **269**, 16–20.
- Fukuda K, Yamada N, Fujitsu Y, Kumagai N and Nishida T (2002) Inhibition of eotaxin expression in human corneal fibroblasts by interferon-γ. *International Archives of Allergy and Immunology* **129**, 138–144.
- Fulkerson PC, Schollaert KL, Bouffi C and Rothenberg ME (2014) IL-5 triggers a cooperative cytokine network that promotes eosinophil precursor maturation. *Journal of Immunology* **193**, 4043–4052.
- García-Zepeda EA, Rothenberg ME, Ownbey RT, Celestin J, Leder P and Luster AD (1996) Human eotaxin is a specific chemoattractant for eosinophil cells and provides a new mechanism to explain tissue eosinophilia. *Nature Medicine* **2**, 449–456.
- Giacomin PR, Gordon DL, Botto M, Daha MR, Sanderson SD, Taylor SM and Dent LA (2008) The role of complement in innate, adaptive and eosinophil-dependent immunity to the nematode *Nippostrongylus brasiliensis*. *Molecular Immunology* **45**, 446–455.
- Gill HS, Gray GD, Watson DL and Husband AJ (1993) Isotype-specific antibody responses to *Haemonchus contortus* in genetically resistant sheep. *Parasite Immunology* **15**, 61–67.
- Gill HS, Altmann K, Cross ML and Husband AJ (2000) Induction of T helper 1- and T helper 2-type immune responses during *Haemonchus contortus* infection in sheep. *Immunology* **99**, 458–463.
- Gleich GJ and Adolphson CR (1986) The eosinophilic leukocyte: structure and function. In Dixon FJ (ed.), *Advances in Immunology*, vol. 39. Orlando: Academic Press, pp. 177–253.
- Gruner L, Mandonnet N, Bouix J, Khang JVT, Cabaret J, Hoste H, Kerboeuf D and Barnouin J (1994) Worm population characteristics and pathological changes in lambs after a single or trickle infection with *Teladorsagia circumcincta*. *International Journal for Parasitology* **24**, 347–356.
- Halliday A, Routledge C, Smith S, Matthews J and Smmith W (2007) Parasite loss and inhibited development of *Teladorsagia circumcincta* in relation to the kinetics of the local IgA response in sheep. *Parasite Immunology* **29**, 425–434.
- Henderson NG and Stear MJ (2006) Eosinophil and IgA responses in sheep infected with *Teladorsagia circumcincta*. *Veterinary Immunology and Immunopathology* **112**, 62–66.
- Henderson WR, Chi EY and Klebanoff SJ (1980) Eosinophil peroxidase-induced mast cell secretion. *The Journal of Experimental Medicine* **152**, 265–279.
- Hernández JN, Hernández A, Stear MJ, Conde-Felipe M, Rodríguez E, Piedrafita D and González JF (2016) Potential role for mucosal IgA in modulating *Haemonchus contortus* adult worm infection in sheep. *Veterinary Parasitology* **223**, 153–158.

- Hertz M, Mahalingam S, Dalum I, Klysner S, Mattes J, Neisig A, Mouritsen S, Foster PS and Gautam A (2001) Active vaccination against IL-5 bypasses immunological tolerance and ameliorates experimental asthma. *The Journal of Immunology* **167**, 3792–3799.
- Horie S, Gleich GJ and Kita H (1996) Cytokines directly induce degranulation and superoxide production from human eosinophils. *Journal of Allergy and Clinical Immunology* **98**, 371–381.
- Huang L and Appleton JA (2016) Eosinophils in helminth infection: defenders and dupes. *Trends in Parasitology* **32**, 798–807.
- Huang L, Gebreselassie NG, Gagliardo LF, Ruyechan MC, Lubber KL, Lee NA, Lee JJ and Appleton JA (2015) Eosinophils mediate protective immunity against secondary nematode infection. *The Journal of Immunology* **194**, 283–290.
- Kalomenidis I, Stathopoulos GT, Barnette R, Guo Y, Peebles RS, Blackwell TS and Light RW (2005) Eotaxin-3 and interleukin-5 pleural fluid levels are associated with pleural fluid eosinophilia in post-coronary artery bypass grafting pleural effusions. *Chest* **127**, 2094–2100.
- Kita H (2011) Eosinophils: multifaceted biological properties and roles in health and disease. *Immunological Reviews* **242**, 161–177.
- Kitaura M, Nakajima T, Imai T, Harada S, Combadiere C, Tiffany HL, Murphy PM and Yoshie O (1996) Molecular cloning of human eotaxin, an eosinophil-selective CC chemokine, and identification of a specific eosinophil eotaxin receptor, CC chemokine receptor 3. *The Journal of Biological Chemistry* **271**, 7725–7730.
- Kitaura M, Suzuki N, Imai T, Takagi S, Suzuki R, Nakajima T, Hirai K, Nomiya H and Yoshie O (1999) Molecular cloning of a novel human CC chemokine (eotaxin-3) that is a functional ligand of CC chemokine receptor 3. *The Journal of Biological Chemistry* **274**, 27975–27980.
- Kusano S, Kukimoto-Niino M, Hino N, Ohsawa N, Ikutani M, Takaki S, Sakamoto K, Hara-Yokoyama M, Shirouzu M, Takatsu K and Yokoyama S (2012) Structural basis of interleukin-5 dimer recognition by its α receptor. *Protein Science* **21**, 850–864.
- Lane, J., Jubb, T., Shephard, R., Webb-Ware, J. and Fordyce, G. (2015). Priority list of endemic diseases for the red meat industries. Report No. B.AHE.0010. Meat and Livestock Australia Limited, North Sydney.
- Locksley RM, Wilson CB and Klebanoff SJ (1982) Role for endogenous and acquired peroxidase in the toxoplasmicidal activity of murine and human mononuclear phagocytes. *The Journal of Clinical Investigation* **69**, 1099–1111.
- Lopez AF, Sanderson CJ, Gamble JR, Campbell HD, Young IG and Vadas MA (1988) Recombinant human interleukin 5 is a selective activator of human eosinophil function. *The Journal of Experimental Medicine* **167**, 219–224.
- Loughran NB, O'Connor B, Ó'Fágáin C and O'Connell MJ (2008) The phylogeny of the mammalian heme peroxidases and the evolution of their diverse functions. *BMC Evolutionary Biology* **8**, 101.
- Maizels RM and Yazdanbakhsh M (2003) Immune regulation by helminth parasites: cellular and molecular mechanisms. *Nature Reviews Immunology* **3**, 733–744.
- McBrien C and Menzies-Gow A (2017) The biology of eosinophils and their role in asthma. *Frontiers in Medicine* **4**, 93.
- McRae KM, Stear MJ, Good B and Keane OM (2015) The host immune response to gastrointestinal nematode infection in sheep. *Parasite Immunology* **37**, 605–613.
- Meeusen ENT and Balic A (2000) Do eosinophils have a role in the killing of helminth parasites? *Parasitology Today* **16**, 95–101.
- Metzler KD, Fuchs TA, Nauseef WM, Reumaux D, Roesler J, Schulze I, Wahn V, Papayannopoulos V and Zychlinsky A (2011) Myeloperoxidase is required for neutrophil extracellular trap formation: implications for innate immunity. *Blood* **117**, 953–959.
- Miyamasu M, Yamaguchi M, Nakajima T, Misaki Y, Morita Y, Matsushima K, Yamamoto K and Hirai K (1999) Th1-derived cytokine IFN- γ is a potent inhibitor of eotaxin synthesis *in vitro*. *International Immunology* **11**, 1001–1004.
- Mochizuki M, Bartels J, Mallet AL, Christophers E and Schröder J-M (1998) IL-4 induces eotaxin: a possible mechanism of selective eosinophil recruitment in helminth infection and atopy. *The Journal of Immunology* **160**, 60–68.
- Motran CC, Silvane L, Chiapello LS, Theumer MG, Ambrosio LF, Volpini X, Celas DP and Cervi L (2018) Helminth infections: recognition and modulation of the immune response by innate immune cells. *Frontiers in Immunology* **9**, 12.
- Muraki M, Gleich GJ and Kita H (2011) Antigen-specific IgG and IgA, but not IgE, activate the effector functions of eosinophils in the presence of antigen. *International Archives of Allergy and Immunology* **154**, 119–127.
- Palframan RT, Collins PD, Williams TJ and Rankin SM (1998) Eotaxin induces a rapid release of eosinophils and their progenitors from the bone marrow. *Blood* **91**, 2240–2248.
- Ponath PD, Qin S, Post TW, Wang J, Wu L, Gerard NP, Newman W, Gerard C and Mackay CR (1996) Molecular cloning and characterization of a human eotaxin receptor expressed selectively on eosinophils. *The Journal of Experimental Medicine* **183**, 2437–2448.
- Rainbird M, Macmillan D and Meeusen E (1998) Eosinophil-mediated killing of *Haemonchus contortus* larvae: effect of eosinophil activation and role of antibody, complement and interleukin-5. *Parasite Immunology* **20**, 93–103.
- Richard M, Grecis RK, Humphreys NE, Renaud J-C and Van Snick J (2000) Anti-IL-9 vaccination prevents worm expulsion and blood eosinophilia in *Trichuris muris*-infected mice. *Proceedings of the National Academy of Sciences* **97**, 767–772.
- Roeber F, Jex AR and Gasser RB (2013) Impact of gastrointestinal parasitic nematodes of sheep, and the role of advanced molecular tools for exploring epidemiology and drug resistance – an Australian perspective. *Parasites & Vectors* **6**, 153.
- Romagnani P, De Paulis A, Beltrame C, Annunziato F, Dente V, Maggi E, Romagnani S and Marone G (1999) Tryptase-chymase double-positive human mast cells express the eotaxin receptor CCR3 and are attracted by CCR3-binding chemokines. *The American Journal of Pathology* **155**, 1195–1204.
- Rosenwasser LJ, Zimmermann N, Hershey GK, Foster PS and Rothenberg ME (2003) Chemokines in asthma: cooperative interaction between chemokines and IL-13. *Journal of Allergy and Clinical Immunology* **111**, 227–242.
- Rothenberg ME and Hogan SP (2006) The eosinophil. *Annual Review of Immunology* **24**, 147–174.
- Rothenberg ME, MacLean JA, Pearlman E, Luster AD and Leder P (1997) Targeted disruption of the chemokine eotaxin partially reduces antigen-induced tissue eosinophilia. *The Journal of Experimental Medicine* **185**, 785–790.
- Rothwell TLW, Windon RG, Horsburgh BA and Anderson BH (1993) Relationship between eosinophilia and responsiveness to infection with *Trichostrongylus colubriformis* in sheep. *International Journal for Parasitology* **23**, 203–211.
- Ruth JH, Lukacs NW, Warmington KS, Polak TJ, Burdick M, Kunkel SL, Strieter RM and Chensue SW (1998) Expression and participation of eotaxin during mycobacterial (type 1) and schistosomal (type 2) antigen-elicited granuloma formation. *The Journal of Immunology* **161**, 4276–4282.
- Sabroe I, Hartnell A, Jopling LA, Bel S, Ponath PD, Pease JE, Collins PD and Williams TJ (1999) Differential regulation of eosinophil chemokine signaling via CCR3 and non-CCR3 pathways. *The Journal of Immunology* **162**, 2946–2955.
- Sallusto F, Mackay CR and Lanzavecchia A (1997) Selective expression of the eotaxin receptor CCR3 by human T helper 2 cells. *Science* **277**, 2005–2007.
- Schratl P, Sturm EM, Royer JF, Sturm GJ, Lippe IT, Peskar BA and Heinemann A (2006) Hierarchy of eosinophil chemoattractants: role of p38 mitogen-activated protein kinase. *European Journal of Immunology* **36**, 2401–2409.
- Scicchitano R, Sheldrake R and Husband A (1986) Origin of immunoglobulins in respiratory tract secretion and saliva of sheep. *Immunology* **58**, 315–321.
- Shinkai A, Yoshisue H, Koike M, Shoji E, Nakagawa S, Saito A, Takeda T, Imabepu S, Kato Y, Hanai N, Anazawa H, Kuga T and Nishi T (1999) A novel human CC chemokine, eotaxin-3, which is expressed in IL-4-stimulated vascular endothelial cells, exhibits potent activity toward eosinophils. *The Journal of Immunology* **163**, 1602–1610.
- Simons JE, Rothenberg ME and Lawrence RA (2005) Eotaxin-1-regulated eosinophils have a critical role in innate immunity against experimental *Brugia malayi* infection. *European Journal of Immunology* **35**, 189–197.
- Sippel V, Pierlot GM, Renault B, Groenen PMA and Strasser DS (2018) Activation of IL5R and CRTH2 on human eosinophils elicit a similar molecular response and reveal a synergistic effect. *European Journal of Molecular & Clinical Medicine* **5**, 1–11.
- Sonoda E, Matsumoto R, Hitoshi Y, Ishii T, Sugimoto M, Araki S, Tominaga A, Yamaguchi N and Takatsu K (1989) Transforming growth factor beta induces IgA production and acts additively with interleukin 5 for IgA production. *The Journal of Experimental Medicine* **170**, 1415–1420.
- Specht S, Saefel M, Arndt M, Endl E, Dubben B, Lee NA, Lee JJ and Hoerauf A (2006) Lack of eosinophil peroxidase or major basic protein

- impairs defense against murine filarial infection. *Infection and Immunity* **74**, 5236–5243.
- Stear MJ, Bishop SC, Doligalska M, Duncan JL, Holmes PH, Irvine J, McCririe L, McKellar QA, Sinski E and Murray M** (1995) Regulation of egg production, worm burden, worm length and worm fecundity by host responses in sheep infected with *Ostertagia circumcincta*. *Parasite Immunology* **17**, 643–652.
- Stear MJ, Henderson NG, Kerr A, McKellar QA, Mitchell S, Seeley C and Bishop SC** (2002) Eosinophilia as a marker of resistance to *Teladorsagia circumcincta* in Scottish Blackface lambs. *Parasitology* **124**, 553–560.
- Swaminathan GJ, Weaver AJ, Loegering DA, Checkel JL, Leonidas DD, Gleich GJ and Acharya KR** (2001) Crystal structure of the eosinophil major basic protein at 1.8 Å: an atypical lectin with a paradigm shift in specificity. *Journal of Biological Chemistry* **276**, 26197–26203.
- Swaminathan GJ, Myszka DG, Katsamba PS, Ohnuki LE, Gleich GJ and Acharya KR** (2005) Eosinophil-granule major basic protein, a C-type lectin, binds heparin. *Biochemistry* **44**, 14152–14158.
- Tavernier J, Devos R, Cornelis S, Tuypens T, Van der Heyden J, Fiers W and Plaetinck G** (1991) A human high affinity interleukin-5 receptor (IL5R) is composed of an IL5-specific α chain and a β chain shared with the receptor for GM-CSF. *Cell* **66**, 1175–1184.
- Terefe G, Griesz C, Prevot F, Bergeaud J-P, Dorchies P, Brunel J-C, Francois D, Fourquaux I and Jacquet P** (2007) *In vitro* pre-exposure of *Haemonchus contortus* L3 to blood eosinophils reduces their establishment potential in sheep. *Veterinary Research* **38**, 647–654.
- Terefe G, Lacroux C, Prévot F, Grisez C, Bergeaud JP, Bleuart C, Dorchies P, Foucras G and Jacquet P** (2009) Eosinophils in *Haemonchus contortus*-infected resistant and susceptible breeds of sheep: abomasal tissue recruitment and *in vitro* functional state. *Veterinary Parasitology* **165**, 161–164.
- Ueki S, Melo RCN, Ghiran I, Spencer LA, Dvorak AM and Weller PF** (2013) Eosinophil extracellular DNA trap cell death mediates lytic release of free secretion-competent eosinophil granules in humans. *Blood* **121**, 2074–2083.
- Uguccioni M, Mackay CR, Ochensberger B, Loetscher P, Rhis S, LaRosa GJ, Rao P, Ponath PD, Baggiolini M and Dahinden CA** (1997) High expression of the chemokine receptor CCR3 in human blood basophils: role in activation by eotaxin, MCP-4, and other chemokines. *The Journal of Clinical Investigation* **100**, 1137–1143.
- Ulrich M, Petre A, Youhnovski N, Prömm F, Schirle M, Schumm M, Pero RS, Doyle A, Checkel J, Kita H, Thiyagarajan N, Acharya KR, Schmid-Grendelmeier P, Simon H-U, Schwarz H, Tsutsui M, Shimokawa H, Bellon G, Lee JJ, Przybylski M and Döring G** (2008) Post-translational tyrosine nitration of eosinophil granule toxins mediated by eosinophil peroxidase. *The Journal of Biological Chemistry* **283**, 28629–28640.
- Urban JF, Jr, Katona IM, Paul WE and Finkelman FD** (1991) Interleukin 4 is important in protective immunity to a gastrointestinal nematode infection in mice. *Proceedings of the National Academy of Sciences USA* **88**, 5513–5517.
- van Egmond M, Damen CA, van Spriel AB, Vidarsson G, van Garderen E and van de Winkel JGJ** (2001) IgA and the IgA Fc receptor. *Trends in Immunology* **22**, 205–211.
- Wagner LA, Ohnuki LE, Parsawar K, Gleich GJ and Nelson CC** (2007) Human eosinophil major basic protein 2: location of disulfide bonds and free sulfhydryl groups. *The Protein Journal* **26**, 13–18.
- Weller PF and Spencer LA** (2017) Functions of tissue-resident eosinophils. *Nature Reviews Immunology* **17**, 746.
- Woof JM and Russell MW** (2011) Structure and function relationships in IgA. *Mucosal Immunology* **4**, 590.
- Yamaguchi Y** (1994) Interleukin-5. *Gan To Kagaku Ryoho* **21**, 1287–1293.
- Ying S, Robinson DS, Meng Q, Barata LT, McEuen AR, Buckley MG, Walls AF, Askenase PW and Kay AB** (1999) C-C chemokines in allergen-induced late-phase cutaneous responses in atopic subjects: association of eotaxin with early 6-hour eosinophils, and of eotaxin-2 and monocyte chemoattractant protein-4 with the later 24-hour tissue eosinophilia, and relationship to basophils and other C-C chemokines (monocyte chemoattractant protein-3 and RANTES). *The Journal of Immunology* **163**, 3976–3984.
- Zuo L and Rothenberg ME** (2007) Gastrointestinal eosinophilia. *Immunology and Allergy Clinics of North America* **27**, 443–455.