

The background of the advertisement is a dark, monochromatic image of several jellyfish. One large jellyfish is in the upper center, with its bell and tentacles clearly visible. To its left is another jellyfish, and below it is a smaller one. In the bottom right corner, another jellyfish is partially visible. The overall aesthetic is scientific and naturalistic.

Life Sciences

Books and Journals from
Cambridge University Press

Cambridge is one of the leading publishers in ecology and conservation biology and publishes high quality texts and research across the breadth of the life sciences, focusing particularly on animal behaviour, biological anthropology, evolutionary biology, computational and systems biology, as well as statistics and professional development titles for biologists.


We also have an extensive portfolio of established journals in agriculture, ecology and conservation, and animal science.

For further details visit:
cambridge.org/core-life-sciences

Cambridge
Core



CAMBRIDGE
UNIVERSITY PRESS



Access
leading
journals in
your subject

Cambridge Core

Explore today at [cambridge.org/core](https://www.cambridge.org/core)

Cambridge Core



CAMBRIDGE
UNIVERSITY PRESS

THE MARINE BIOLOGICAL ASSOCIATION OF THE UNITED KINGDOM

TERMS OF MEMBERSHIP FOR 2018

| MBA MEMBERSHIP (prices per annum) | Price from 1st March 2018 |
|--------------------------------------|------------------------------------|
| Young Marine Biologist (YMB) | £12.50 |
| Student Member | £25.00 |
| Associate Member | £40.00 (direct debit discount £5) |
| Professional Member, Mem.MBA | £45.00 (direct debit discount £5) |
| MBA Fellow, FMBA | £120.00 (direct debit discount £5) |
| Institutional Member | £340.00 |

| JOURNAL SUBSCRIPTION (MBA price added to the above membership fees, with the exception of YMB as no Journal option for this category) | |
|--|--------|
| Online Journal | £20.00 |
| Print & Online Journal | £80.00 |

The Association is a learned society founded in 1884 and incorporated by Royal Charter in 2013. The Marine Biological Association (MBA) provides a unified, clear, independent voice on behalf of the marine biological community. In 1888, the MBA opened the Plymouth laboratory at Citadel Hill.

The MBA is also a registered charity with its key aims being to promote scientific research into all aspects of life in the sea, including the environment on which it depends, and to disseminate to the public the knowledge gained. The Patron of the Association is His Royal Highness the Duke of Edinburgh, who is also an Honorary Fellow. The other Honorary Fellows are His Serene Highness Prince Albert of Monaco, Dr Sylvia Earle, Sir Tim Hunt and Professor James Lovelock.

The work of the Association is controlled by the Council elected annually by its subscribing members.

The Marine Biological Association, under the direction of its Council, has undertaken research in all branches of marine science and many of the results have been published in this Journal. An account of the history of the first hundred years of the Association and Journal may be found in Volume 67 (pp. 463 and 465).

Since its foundation the Association has been supported by grants from Research Councils and by subscriptions and donations

from private members, universities, learned societies, the Fishmongers' Company and other public bodies.

The Marine Biological Association is a very active scientific learned society with a large international membership. The MBA also manages an internationally renowned research programme.

Subscriptions can be paid by cash/cheque, direct debit from a UK bank account or credit card. *Subscriptions are reduced by £5 per annum if paying by direct debit.

Members of the Association have a wide range of rights and privileges including the use of post-nominals (FMBA for MBA Fellows, Mem.MBA for Professional Members) and being entitled to receive The Marine Biologist Magazine. The Commissioners of Inland Revenue have approved the Association for the purposes of Section 16, Finance Act, 1958, and that the whole of the annual subscriptions paid by a member who qualifies for relief under the section will be allowable as a deduction from his emoluments assessable to income tax.

All correspondence should be addressed to: The Membership Secretary, Marine Biological Association, The Laboratory, Citadel Hill, Plymouth, PL1 2PB, UK (Tel. +44 (0) 1752 426493; e-mail: membership@mba.ac.uk).

PUBLISHED BY THE PRESS SYNDICATE OF THE UNIVERSITY OF CAMBRIDGE
The Pitt Building, Trumpington Street, Cambridge, CB2 1RP, United Kingdom

This journal issue has been printed on FSC-certified paper and cover board. FSC is an independent, non-governmental, not-for-profit organization established to promote the responsible management of the world's forests. Please see www.fsc.org for information.

Printed by Bell and Bain Limited, Glasgow

CAMBRIDGE UNIVERSITY PRESS

Journals Fulfillment Department, UPH, Shaftesbury Road, Cambridge CB2 8BS, United Kingdom
1 Liberty Plaza, Floor 20, New York, NY 10006, USA
477 Williamstown Road, Port Melbourne, VIC 3207, Australia
C/ Orense, 4, planta 13, 28020, Madrid, Spain
Dock House, The Waterfront, Cape Town 8001, South Africa

CONTENTS

| | |
|--|------|
| Lewis, J. Editorial | 1217 |
| Corrales-Ugalde, M., Quesada, A.J., Naranjo-Elizondo, B. and Cortés, J. New records of gelatinous zooplankton from an oceanic island in the Eastern Tropical Pacific | 1219 |
| Westgarth-Smith, A.R. Ocean acidification needs more publicity as part of a strategy to avoid a global decline in calcifier populations | 1227 |
| Brash, J.M., Cook, R.L., Mackenzie, C.L. and Sanderson, W.G. The demographics and morphometries of biogenic reefs: important considerations in conservation management | 1231 |
| Mecho, A., Aguzzi, J., de Mol, B., Lastras, G., Ramirez-Llodra, E., Bahamon, N., Company, J.B. and Canals, M. Visual faunistic exploration of geomorphological human-impacted deep-sea areas of the north-western Mediterranean Sea | 1241 |
| Gordon, J.M. and Knights, A.M. Revisiting Connell: competition but not as we know it | 1253 |
| Liu, Q., Lu, B., Song, X., Li, Y., Gao, Y., Li, H., Wang, M., Shao, H., Warren, A., Chen, X., Jiang, Y. and Lin, X. Recovery of ciliated protozoan communities in response to environmental change in a shrimp-farming pond in southern China | 1263 |
| Plotkin, A., Gerasimova, E. and Rapp, H.T. Polymastiidae (Porifera: Demospongiae) of the Nordic and Siberian Seas | 1273 |
| El Asri, F., Zidane, H., Errhif, A., Tamsouri, M.-N., Maanan, M., Malouli Idrissi, M. and Martin, D. Polychaete diversity and assemblage structure in the Oualidia Lagoon, Moroccan Atlantic coast | 1337 |
| de León-González, J.A., Méndez, N. and Navedo, J.G. <i>Laonereis watsoni</i> (Annelida, Nereididae), a new species from western Mexico | 1347 |
| Bigatti, G., Diaz de Vivar, M.E., Cumplido, M., Nieto Vilela, R.A., Avaro, M., Sastre, V. and Gil, M. Fatty acids and contaminants in edible marine gastropods from Patagonia | 1355 |
| Simone, L.R.L. Convergence with naticids: phenotypic phylogenetic study on some Antarctic littorinoideans, with description of the zerotulid new genus <i>Pseudonatica</i> , and its presence in Brazil (Mollusca, Caenogastropoda) | 1365 |
| Markaida, U., Flores, L., Arias, E. and Mora, E. Reproduction and population structure of <i>Octopus mimus</i> fished in a Marine Protected Area of Ecuador | 1383 |
| Gates, A.R., Sheader, M., Williams, J.A. and Hawkins, L.E. Infection with cerebral metacercariae of microphallid trematode parasites reduces reproductive output in the gammarid amphipod <i>Gammarus insensibilis</i> (Stock 1966) in UK saline lagoons | 1391 |
| Sepahvand, V., Rastegar-Pouyani, N. and Kihara, T.C. Two new species of <i>Clausidium</i> copepods (Crustacea, Poecilostomatoida) associated with ghost shrimps from Iran | 1401 |
| Crickenberger, S. and Wetthey, D.S. Reproductive physiology, temperature and biogeography: the role of fertilization in determining the distribution of the barnacle <i>Semibalanus balanoides</i> | 1411 |
| Weiss, R., Torrecilla, Z., González-Ortegón, E., González-Tizón, A.M., Martínez-Lage, A. and Schubart, C.D. Genetic differentiation between Mediterranean and Atlantic populations of the common prawn <i>Palaemon serratus</i> (Crustacea: Palaemonidae) reveals uncommon phylogeographic break | 1425 |
| Kornienko, E.S., Golubinskaya, D.D., Korn, O.M. and Sharina, S.N. The complete description of larval stages of the lobster shrimp <i>Leonardsaxius amurensis</i> (Kobjakova, 1937) (Decapoda: Axiidea: Axiidae) identified by DNA barcoding | 1435 |
| Villegas-Hernández, H., Poot-López, G.R., López-Rocha, J.A., González-Salas, C. and Guillen-Hernández, S. Abundance and catchability estimates of the Atlantic blue crab <i>Callinectes sapidus</i> based on mark-recapture data from the northern Yucatan Peninsula | 1455 |
| Pinoni, S.A., Jerez-Cepa, I., López Mañanes, A.A. and Mancera Romero, J.M. The euryhaline crab <i>Uca tangeri</i> showed metabolic differences to sex and environmental salinity | 1465 |
| Castillo-Hidalgo, G., Plaza, G., Díaz-Astudillo, M. and Landaeta, M.F. Seasonal variations in early life traits of <i>Sindoscopus australis</i> (Blennioidei: Dactyloscopidae): hatching patterns, larval growth and bilateral asymmetry of otoliths | 1477 |
| Jawad, L.A., Fricke, R. and Näslund, J. Comparative osteology of the family Tripterygiidae (Teleostei: Blenniiformes) | 1487 |
| Ortega Garcia, S., Arizmendi-Rodríguez, D.I. and Zúñiga-Flores, M.S. Striped marlin (<i>Kajikia audax</i>) diet variability off Cabo San Lucas, B.C.S., Mexico during El Niño–La Niña events | 1513 |
| Guimarães, S.M., Tavares, D.C. and Monteiro-Neto, C. Incidental capture of sea turtles by industrial bottom trawl fishery in the Tropical South-western Atlantic | 1525 |