

# Relativistic jets in Narrow-Line Seyfert 1

L. Foschini<sup>1</sup>, E. Angelakis<sup>2</sup>, G. Bonnoli<sup>1</sup>, G. Calderone<sup>3</sup>, M. Colpi<sup>3</sup>,  
F. D'Ammando<sup>4</sup>, D. Donato<sup>5</sup>, A. Falcone<sup>6</sup>, L. Fuhrmann<sup>2</sup>,  
G. Ghisellini<sup>1</sup>, G. Ghirlanda<sup>1</sup>, M. Hauser<sup>7</sup>, Y. Y. Kovalev<sup>2,8</sup>,  
L. Maraschi<sup>1</sup>, E. Nieppola<sup>9</sup>, J. Richards<sup>10</sup>, A. Stamerra<sup>11</sup>,  
G. Tagliaferri<sup>1</sup>, F. Tavecchio<sup>1</sup>, D. J. Thompson<sup>5</sup>, O. Tibolla<sup>12</sup>,  
A. Tramacere<sup>13</sup>, S. Wagner<sup>7</sup>

<sup>1</sup>INAF – Osservatorio Astronomico di Brera, 23807 Merate (LC), Italy  
email: luigi.foschini@brera.inaf.it

<sup>2</sup>Max-Planck-Institut für Radioastronomie, 53121 Bonn, Germany

<sup>3</sup>University of Milano Bicocca, 20100 Milano, Italy

<sup>4</sup>INAF – IASF-Palermo, 90146, Palermo, Italy

<sup>5</sup>NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA

<sup>6</sup>Penn State University, University Park, PA 16802, USA

<sup>7</sup>Landessternwarte, Universität Heidelberg, Königstuhl, D 69117 Heidelberg, Germany

<sup>8</sup> Astro Space Center of the Lebedev Physical Institute, 117997 Moscow, Russia

<sup>9</sup>Metsähovi Radio Observatory, FIN-02540 Kylmala, Finland

<sup>10</sup>California Institute of Technology, Pasadena, CA 91125, USA

<sup>11</sup>University of Siena, 53100, Siena, Italy

<sup>12</sup>University of Würzburg, 97074, Würzburg, Germany

<sup>13</sup>INTEGRAL Science Data Centre, CH-1290, Versoix, Switzerland


**Abstract.** Narrow-Line Seyfert 1 (NLS1) class of active galactic nuclei (AGNs) is generally radio-quiet, but a small percent of them are radio-loud. The recent discovery by *Fermi*/LAT of high-energy  $\gamma$ -ray emission from 4 NLS1s proved the existence of relativistic jets in these systems. It is therefore important to study this new class of  $\gamma$ -ray emitting AGNs. Here we report preliminary results about the observations of the July 2010  $\gamma$ -ray outburst of PMN J0948+0022, when the source flux exceeded for the first time  $10^{-6}$  ph cm<sup>-2</sup> s<sup>-1</sup> ( $E > 100$  MeV).

**Keywords.** Galaxies: jets – Galaxies: Seyfert – Gamma-rays: observations

---

The recent discovery of variable  $\gamma$ -ray emission from 4 NLS1s revealed the presence of a third class of  $\gamma$ -ray AGNs (Abdo *et al.* 2009a). This poses intriguing questions to the current knowledge of relativistic jet systems and on how these structures are generated. One of these sources, PMN J0948 + 0022 ( $z = 0.5846$ ) is classified a typical NLS1, but it displays also strong, compact and variable radio emission, with inverted spectrum, suggesting the possibility of the presence of a relativistic jet (Zhou *et al.* 2003). The confirmation came with the detection of high-energy variable  $\gamma$  rays by *Fermi*/LAT (Abdo *et al.* 2009b, Foschini *et al.* 2010a). A multiwavelength campaign performed in March-July 2009 displayed coordinated variability at all frequencies, thus confirming that the source detected by *Fermi* is indeed the high-energy counterpart of PMN J0948 + 0022 (Abdo *et al.* 2009c).

PMN J0948 + 0022, the first NLS1 detected at  $\gamma$  rays, was soon followed by three more (Abdo *et al.* 2009a). Their main differences with respect to blazars and radio galaxies are the optical spectrum and the radio morphology, which is quite compact and



**Figure 1.** *Fermi*/LAT lightcurve (0.1 – 100 GeV) of PMN J0948 + 0022 in July-August 2010, with 3 days time bin. Blue points are detection with  $TS > 10$  ( $\sim 3\sigma$ ), while the others are  $2\sigma$  upper limits. LAT data have been analyzed as described in Foschini *et al.* (2010b).

without extended structures. These characteristics point to systems with relatively low masses of the central black hole ( $10^{6-8} M_{\odot}$ ) and high accretion rates (up to 90% of the Eddington value), common for NLS1s, but not for blazars or radio galaxies. In addition, since the  $\gamma$ -ray NLS1s seem similar to blazars, i.e. small viewing angles, there should be a parent population with the jet viewed at large angles (as blazars vs radio galaxies). The first source of this type has been recently found (PKS 0558 – 504, Gliozzi *et al.* 2010). Therefore, it seems that NLS1s could be a low mass set of systems “parallel” to blazars and radio galaxies. A key question was the power released by jets of NLS1s. The early observations and the 2009 MW campaign have shown that the maximum luminosity reached by PMN J0948 + 0022, the most powerful of these NLS1s, is  $\sim 10^{47}$  erg  $\text{s}^{-1}$  (0.1–100 GeV). On the other hand, blazars can reach greater luminosities ( $\sim 10^{49}$  erg  $\text{s}^{-1}$  in the case of 3C 454.3, e.g. Foschini *et al.* 2010b). The question was answered in July 2010, when PMN J0948+0022 underwent a strong outburst (Donato *et al.* 2010, Foschini 2010c) with a peak flux of  $\sim 10^{-6}$  ph  $\text{cm}^{-2}$   $\text{s}^{-1}$  (0.1 – 100 GeV), corresponding to a luminosity of  $\sim 10^{48}$  erg  $\text{s}^{-1}$  (Fig. 1). Even if the source position was too close to the Sun for a full MW campaign, some coverage was obtained. Further details will be available in a forthcoming paper.

## References

- Abdo A. A. *et al.*, 2009a, *ApJ* 707, L142.  
 Abdo A. A. *et al.*, 2009b, *ApJ* 699, 976.  
 Abdo A. A. *et al.*, 2009c, *ApJ* 707, 727.  
 Donato D. *et al.*, 2010, *ATel* 2733, 12 July 2010.  
 Foschini L. *et al.*, 2010a, in: *Proceedings of Accretion and ejection in AGN: a global view*. L. Maraschi, G. Ghisellini, R. Della Ceca & F. Tavecchio eds, ASP Conference Series 427, San Francisco, in press (arXiv:0908.3313).  
 Foschini L. *et al.*, 2010b, *MNRAS*, in press (arXiv:1004.4518).  
 Foschini L., 2010c, *ATel* 2752, 22 July 2010.  
 Gliozzi M. *et al.*, 2010, *ApJ* 717, 1243.  
 Zhou H. Y. *et al.*, 2003, *ApJ* 584, 147.