


# Nearby kinematic wiggles from LEGUE

J. L. Carlin<sup>1</sup>, H. J. Newberg<sup>1</sup>, L. Deng<sup>2</sup>, J. Delaunay<sup>1</sup>, D. Gole<sup>1,3</sup>,  
 K. Grabowski<sup>1</sup>, C. Liu<sup>2</sup>, Y. Xu<sup>2</sup>, F. Yang<sup>2</sup>, and H. Zhang<sup>2</sup>

<sup>1</sup>Rensselaer Polytechnic Institute, Troy, NY, USA, [carlij@rpi.edu](mailto:carlij@rpi.edu), <sup>2</sup>National Astronomical Observatories, Chinese Academy of Sciences, Beijing, <sup>3</sup>SUNY Geneseo, Geneseo, NY, USA

In its first two observing seasons, the LEGUE (LAMOST Experiment for Galactic Understanding and Exploration; Deng *et al.*, Zhao *et al.* 2012) survey has obtained  $\sim 1.7$  million science-quality spectra. We apply corrections to the PPMXL proper motions (PMs; Roeser *et al.* 2010) as a function of position, as determined from the measured PMs of extragalactic objects discovered in LAMOST spectra (see Fig. 1, left and center panels). LAMOST radial velocities and corrected PMs are used to derive 3D space velocities for  $\sim 480,000$  F-stars (assuming  $M_V = 4$  to derive distances). The right panel of Fig. 1 shows the radial component of Galactic cylindrical velocities ( $V_R$ ) for stars between  $7.8 < R_{GC} < 9.8$  kpc (with  $R_{\odot,GC} = 7.8$  kpc) as a function of height ( $Z$ ) and angle ( $\theta$ ) from the Galactic  $X$ -axis. Each dot represents the average position of stars in a  $200 \times 200$  pc box, color-coded by the mean  $V_R$  of those stars. Assuming circular rotation,  $V_R$  should be zero. This is true on average for  $\theta > 0^\circ$  (3rd Galactic quadrant), but not for  $\theta < 0^\circ$ . The velocities are also asymmetric across the Galactic plane for  $\theta < 0^\circ$  (2nd quadrant), with most positions  $\langle V_R \rangle > 0$  above the disk (radially outward), and  $\langle V_R \rangle < 0$  below the disk. Similar structure to this apparent “shearing” motion has been seen in RAVE (e.g., Williams *et al.* 2013; Siebert *et al.* 2012), and SDSS (Widrow *et al.* 2012).



**Figure 1.** *Left, center panels:* Proper motion residuals, which should be zero, for extragalactic objects identified with LAMOST spectra, as a function of RA along two declination stripes. Blue curves show a polynomial fit used to correct the PPMXL PMs. *Right panel:* Average  $V_R$  velocities (color-coding of points) for the selected stars as a function of  $(Z, \theta)$  position. For  $\theta < 0^\circ$ , stars above the plane are predominantly moving away from the Galactic center, while those below are moving inward (i.e.,  $V_R < 0$ ).

## References

- Deng, L.-C., Newberg, H. J., Liu, C., *et al.* 2012, *RAA*, 12, 735  
 Roeser, S., Demleitner, M., & Schilbach, E. 2010, *AJ*, 139, 2440  
 Siebert, A., Famaey, B., Binney, J., *et al.* 2012, *MNRAS*, 425, 2335  
 Widrow, L. M., Gardner, S., Yanny, B., Dodelson, S., & Chen, H.-Y. 2012, *ApJL*, 750, L41  
 Williams, M. E. K., Steinmetz, M., Binney, J., *et al.* 2013, *arXiv:1302.2468*  
 Zhao, G., Zhao, Y.-H., Chu, Y.-Q., Jing, Y.-P., & Deng, L.-C. 2012, *RAA*, 12, 723