

## A REMARK ON HOMOGENEOUS CONVEX DOMAINS

SATORU SHIMIZU

### §0. Introduction

In this note, by a homogeneous convex domain in  $\mathbf{R}^n$  we mean a convex domain  $\Omega$  in  $\mathbf{R}^n$  containing no complete straight lines on which the group  $G(\Omega)$  of all affine transformations of  $\mathbf{R}^n$  leaving  $\Omega$  invariant acts transitively. Let  $\Omega$  be a homogeneous convex domain. Then  $\Omega$  admits a  $G(\Omega)$ -invariant Riemannian metric which is called the canonical metric (see [11]). The domain  $\Omega$  endowed with the canonical metric is a homogeneous Riemannian manifold and we denote by  $I(\Omega)$  the group of all isometries of it. A homogeneous convex domain  $\Omega$  is called reducible if there is a direct sum decomposition of the ambient space  $\mathbf{R}^n = \mathbf{R}^{n_1} \times \mathbf{R}^{n_2}$ ,  $n_i > 0$ , such that  $\Omega = \Omega_1 \times \Omega_2$  with  $\Omega_i$  a homogeneous convex domain in  $\mathbf{R}^{n_i}$ ; and if there is no such decomposition, then  $\Omega$  is called irreducible.

The purpose of this note is to prove the following:

**THEOREM.** *Let  $M$  be a homogeneous Riemannian manifold whose universal covering is isometric to a homogeneous convex domain  $\Omega$  in  $\mathbf{R}^n$  endowed with the canonical metric. If  $\Omega$  is irreducible and not affinely equivalent to a convex cone, then  $M$  is simply connected, that is,  $M$  itself is isometric to  $\Omega$ .*

It is already known in [2] that an analogous fact holds for a homogeneous bounded domain in  $\mathbf{C}^n$ .

We prove the above theorem along the same line as in [2] by using results of Tsuji [9], [10].

The author would like to thank Professor Tsuji for his helpful advices.

### §1. The center of a group of affine automorphisms of $\Omega$

First we discuss the connection between the irreducibilities of a homogeneous convex domain and the cone fitted onto it. For the purpose we

---

Received November 13, 1984.

need the notion of  $T$ -algebras. The details for it can be found in [11].

Let  $\Omega$  be a homogeneous convex domain in  $\mathbf{R}^n$  and  $V$  the cone fitted onto it, that is,

$$V = \{(\lambda x, \lambda) \in \mathbf{R}^n \times \mathbf{R} \mid x \in \Omega, \lambda > 0\}.$$

Note that  $V$  is a homogeneous convex cone in  $\mathbf{R}^{n+1}$  (cf. the proof of Proposition 2 in this section). By a theorem of Vinberg [11], we may assume that  $\Omega = \Omega(\mathfrak{A})$  and  $V = V(\mathfrak{A})$ , where  $\mathfrak{A} = \sum_{1 \leq i, j \leq r} \mathfrak{A}_{ij}$  is a  $T$ -algebra of rank  $r$  ( $r \geq 2$ ) and the notations  $V(\mathfrak{A})$  and  $\Omega(\mathfrak{A})$  bear the same meanings as in [9], [10]. We put  $\dim \mathfrak{A}_{ij} = n_{ij}$ . A criterion for  $\Omega$  and  $V$  to be irreducible can be given in terms of the  $T$ -algebra  $\mathfrak{A}$  as follows:

(i) (Tsuji [10])  $\Omega = \Omega(\mathfrak{A})$  is irreducible if and only if, for every pair  $(i, j)$  of indices with  $1 \leq i \leq j \leq r - 1$ , there exists a series  $i_0, i_1, \dots, i_p$  of indices such that  $1 \leq i_\alpha \leq r - 1$  ( $0 \leq \alpha \leq p$ ),  $i_0 = i$ ,  $i_p = j$  and  $n_{i_{\alpha-1}i_\alpha} \neq 0$  ( $1 \leq \alpha \leq p$ ).

(ii) (Asano [1])  $V = V(\mathfrak{A})$  is irreducible if and only if, for every pair  $(i, j)$  of indices with  $1 \leq i \leq j \leq r$ , there exists a series  $i_0, i_1, \dots, i_p$  of indices such that  $1 \leq i_\alpha \leq r$  ( $0 \leq \alpha \leq p$ ),  $i_0 = i$ ,  $i_p = j$  and  $n_{i_{\alpha-1}i_\alpha} \neq 0$  ( $1 \leq \alpha \leq p$ ).

**PROPOSITION 1.** *In the above notation, if  $\Omega$  is irreducible and not affinely equivalent to a convex cone, then  $V$  is irreducible.*

*Proof.* Since  $\Omega = \Omega(\mathfrak{A})$  is not affinely equivalent to a convex cone by assumption, it follows from the definition of  $\Omega(\mathfrak{A})$  that there exists an index  $i$  such that  $1 \leq i \leq r - 1$  and  $n_{ir} \neq 0$ . By (i) and (ii), this implies that  $V$  is irreducible. q.e.d.

*Remark.* If  $\Omega$  is a convex cone, then the cone  $V$  fitted onto  $\Omega$  is reducible. In fact, one has  $V = \Omega \times \mathbf{R}_+$ , where  $\mathbf{R}_+$  denotes the cone of positive real numbers.

We fix notations. Let  $G$  be a group. For a subset  $H$  of  $G$ ,  $C_G(H)$  denotes the centralizer of  $H$  in  $G$ , and the center of  $G$  is denoted simply by  $C(G)$ . When  $G$  is a topological group, the connected component of  $G$  containing the identity element is denoted by  $G^\circ$ . The unit element of a group is denoted by  $e$ . The identity matrix of degree  $n$  is denoted by  $1_n$ .  $A(n, \mathbf{R})$  denotes the group of all affine transformations of  $\mathbf{R}^n$ .

The aim of this section is to prove the following:

**PROPOSITION 2.** *Let  $\Omega$  be an irreducible homogeneous convex domain*

in  $\mathbf{R}^n$  which is not affinely equivalent to a convex cone. If a subgroup  $G$  of  $G(\Omega)$  acts transitively on  $\Omega$ , then one has  $C_{A(n, \mathbf{R})}(G) = \{e\}$  and hence  $C(G) = \{e\}$ . In particular, one has  $C(G(\Omega)) = C(G(\Omega)^*) = \{e\}$ .

For the proof, we need the following result:

(iii) (Rothaus [7]). Let  $V$  be an irreducible homogeneous convex cone in  $\mathbf{R}^n$ . If a subgroup  $G$  of  $G(V)$  acts transitively on  $V$ , then one has  $C_{GL(n, \mathbf{R})}(G) = \{\lambda 1_n \mid \lambda \in \mathbf{R}\}$ .

*Proof of Proposition 2.* Let  $V$  be the cone fitted onto  $\Omega$ . Let  $\rho$  denote the group homomorphism

$$A(n, \mathbf{R}) \ni a \longmapsto \begin{pmatrix} f(a) & q(a) \\ 0 & 1 \end{pmatrix} \in GL(n + 1, \mathbf{R}),$$

where  $f(a)$  and  $q(a)$  denote, respectively, the linear and the translation parts of  $a \in A(n, \mathbf{R})$ . Then one has  $\rho(G(\Omega)) \subset G(V)$ . The pair  $(\rho, \iota)$  of the group homomorphism  $\rho: G(\Omega) \rightarrow G(V)$  and the natural embedding  $\iota: \Omega \rightarrow V$  given by  $\iota(x) = (x, 1)$  is equivariant, that is,  $\iota(ax) = \rho(a)\iota(x)$  for all  $a \in G(\Omega)$ ,  $x \in \Omega$ . Since  $G$  acts transitively on  $\Omega$  by assumption, this shows that the subgroup  $G' = \rho(G) \cdot \{\lambda 1_{n+1} \mid \lambda > 0\}$  of  $G(V)$  acts transitively on  $V$ . By Proposition 1,  $V$  is an irreducible homogeneous convex cone in  $\mathbf{R}^{n+1}$ . Therefore, using (iii), we see  $C_{GL(n+1, \mathbf{R})}(G') = \{\lambda 1_{n+1} \mid \lambda \in \mathbf{R}\}$ . Let  $a \in C_{A(n, \mathbf{R})}(G)$ . Then one has  $\rho(a) \in C_{GL(n+1, \mathbf{R})}(G')$ . Hence  $\rho(a)$  is a scalar matrix and this implies  $a = e$  by the definition of  $\rho$ . q.e.d.

A homogeneous convex domain  $\Omega(n)$  in  $\mathbf{R}^n$  ( $n \geq 2$ ) defined by

$$\Omega(n) = \{(x^1, \dots, x^n) \in \mathbf{R}^n \mid x^1 > (x^2)^2 + \dots + (x^n)^2\}$$

is called the elementary domain. Every elementary domain is irreducible and not affinely equivalent to a convex cone. The following result is known:

(iv) (Tsuji [9]). Let  $\Omega$  be an irreducible homogeneous convex domain which is not affinely equivalent to the elementary domain. Then one has  $I(\Omega)^* = G(\Omega)^*$ .

Combining (iv) with Proposition 2, we obtain

**LEMMA.** *Let  $\Omega$  be an irreducible homogeneous convex domain which is affinely equivalent to neither a convex cone nor the elementary domain. If a connected Lie subgroup  $G$  of  $I(\Omega)$  acts transitively on  $\Omega$ , then one has  $C(G) = \{e\}$ .*

*Remark.* The above lemma remains valid for the elementary domain (cf. the proof of our theorem in the next section and Corollary 2 in Section 3).


**§2. Proof of Theorem**

First, suppose  $\Omega$  is affinely equivalent to the elementary domain  $\Omega(n)$ . Then, since  $\Omega(n)$  endowed with the canonical metric is of negative sectional curvature (see, e.g., [8]),  $M$  is a connected homogeneous Riemannian manifold of negative sectional curvature. Hence our assertion follows from [6, Theorem 8.3, p. 105].

Next, suppose  $\Omega$  is not affinely equivalent to the elementary domain. We set  $G = I(M)^\circ$ . Then one has a natural identification  $M = G/K$ , where  $K$  is an isotropy subgroup of  $G$  at some point of  $M$ . Let  $\tilde{G}'$  be the universal covering group of  $G$  and let  $\pi$  be the covering projection of  $\tilde{G}'$  onto  $G$ . Then one has  $M \simeq \tilde{G}'/\pi^{-1}(K)$  and  $\Omega \simeq \tilde{G}'/\tilde{K}'$ , where  $\tilde{K}' = \pi^{-1}(K)^\circ$ . We put

$$\begin{aligned} \Delta_0 &= \{g \in \tilde{G}' \mid g \cdot y = y \text{ for all } y \in \Omega\}, \\ \Delta &= \{g \in \tilde{G}' \mid g \cdot x = x \text{ for all } x \in M\}. \end{aligned}$$

It follows that  $\Delta_0 \subset \Delta$ . We note that, since  $G$  acts effectively on  $M$ ,  $\Delta$  is a discrete subgroup of  $\tilde{G}'$ . Put  $\tilde{G}'/\Delta_0 = \tilde{G}$  and  $\tilde{K}'/\Delta_0 = \tilde{K}$ . Then  $\tilde{G}$  is a connected Lie subgroup of  $I(\Omega)$ , and one has  $\Omega \simeq \tilde{G}/\tilde{K}$ . Moreover, one has the following commutative diagram:



Since  $\ker \pi' \subset C(\tilde{G})$ , we see by the lemma in the previous section that  $\pi'$  is an isomorphism of  $\tilde{G}$  onto  $G$ . Therefore  $\pi'^{-1}(K)$  is compact, because so is  $K$ . It is easy to see  $\tilde{K} = \pi'^{-1}(K)^\circ$ , and hence  $\tilde{K}$  is compact. Since  $\Omega$  is a cell (see [11]) and since  $\Omega \simeq \tilde{G}/\tilde{K}$ ,  $\tilde{K}$  is a maximal compact subgroup of  $\tilde{G}$ . Therefore one has  $\tilde{K} = \pi'^{-1}(K)$ , and this implies  $\Omega \simeq \tilde{G}/\tilde{K} \simeq G/K = M$ . q.e.d.

**§3. Corollaries and Remarks**

An affine manifold  $M$  of dimension  $n$  is a manifold which admits an atlas  $\{(U_\alpha, \phi_\alpha)\}$  such that each coordinate change  $\phi_\alpha \circ \phi_\beta^{-1}$  is an affine trans-

formation of  $\mathbb{R}^n$  (cf. [5]). A diffeomorphism  $f$  of  $M$  is called an affine transformation of  $M$  if it is affine with respect to the atlas  $\{(U_\alpha, \phi_\alpha)\}$ , that is, if each transformation  $\phi_\alpha \circ f \circ \phi_\beta^{-1}$  is an affine transformation of  $\mathbb{R}^n$ , and  $M$  is called homogeneous if the group  $G(M)$  of all affine transformations of  $M$  acts transitively on it. Note that a domain  $\Omega$  in  $\mathbb{R}^n$  is naturally an affine manifold and the group  $G(\Omega)$  defined in the introduction coincides with the one defined above.

**COROLLARY 1.** *Let  $M$  be a homogeneous affine manifold whose universal covering is affinely equivalent to a homogeneous convex domain  $\Omega$  in  $\mathbb{R}^n$ . If  $\Omega$  is irreducible and not affinely equivalent to a convex cone, then  $M$  is simply connected, that is,  $M$  itself is affinely equivalent to  $\Omega$ .*

*Proof.* Let  $\Gamma$  be the covering transformation group of the covering  $\Omega \rightarrow M$ . By assumption,  $\Gamma$  is a subgroup of  $G(\Omega)$ , and hence the canonical metric of  $\Omega$  is  $\Gamma$ -invariant. With respect to the induced Riemannian metric,  $M$  is a homogeneous Riemannian manifold. Indeed, since every element of  $G(M)$  lifts to an element of  $G(\Omega) \subset I(\Omega)$ ,  $G(M)$  acts as an isometry group, and its action on  $M$  is transitive by assumption. Thus the theorem shows that  $M$  is simply connected. q.e.d.

**COROLLARY 2.** *Let  $\Omega$  be an irreducible homogeneous convex domain which is not affinely equivalent to a convex cone. If a Lie subgroup  $G$  of  $I(\Omega)$  acts transitively on  $\Omega$ , then one has  $C(G) = \{e\}$ . In particular, one has  $C(I(\Omega)) = \{e\}$ .*

*Proof.* If  $\Omega$  is affinely equivalent to the elementary domain, then this is a direct consequence of [6, Theorem 8.4, p. 107] (cf. Proof of Theorem). Otherwise, the proof goes as follows: Since  $C(G) \subset C(\bar{G})$ , where  $\bar{G}$  is the closure of  $G$  in  $I(\Omega)$ , we may assume that  $G$  is a closed subgroup of  $I(\Omega)$ . The subgroup  $C(G)$  of  $I(\Omega)$  is discrete. Indeed, using the lemma in Section 1, we see  $C(G)^\circ \subset C(G^\circ) = \{e\}$ . The same reasoning as in the proof of [6, Theorem 8.4] yields that  $C(G)$  acts properly discontinuously and freely on  $\Omega$  and the quotient space  $C(G)\backslash\Omega$  is a homogeneous Riemannian manifold with respect to the induced Riemannian metric. By the theorem,  $C(G)\backslash\Omega$  is simply connected. Hence we conclude that  $C(G) = \{e\}$ . q.e.d.

*Remark 1.* In our theorem and Corollary 1, the assumption that  $\Omega$  is not affinely equivalent to a convex cone can not be removed. Indeed, let  $\Omega$  be a homogeneous convex cone in  $\mathbb{R}^n$  and put  $M = \Gamma\backslash\Omega$ , where  $\Gamma =$

$\{2^k 1_n \mid k \in \mathbb{Z}\} \subset G(\Omega)$ . Since  $\Gamma \subset C(G(\Omega))$ , the transitive action of  $G(\Omega)$  on  $\Omega$  induces a transitive action of  $G(\Omega)$  on  $M = \Gamma \backslash \Omega$  as an affine transformation group. This implies that  $M$  is a homogeneous affine manifold whose universal covering is affinely equivalent to  $\Omega$ . Therefore  $M$  is also a homogeneous Riemannian manifold whose universal covering is isometric to  $\Omega$  endowed with the canonical metric. However  $M$  is clearly not simply connected.

*Remark 2.* Consider the following problem:

Let  $M$  be an  $n$ -dimensional homogeneous affine manifold which is projectively hyperbolic in the sense of Kobayashi [4]. Then, is  $M$  a homogeneous convex domain in  $\mathbb{R}^n$ ?

This is an affine analogue of Kobayashi's problem concerning homogeneous hyperbolic (complex) manifolds (cf. [3, Problem 12, p. 133]).

Since the intrinsic distance of  $M$  is complete, the universal covering of  $M$  is affinely equivalent to a convex domain in  $\mathbb{R}^n$  containing no complete straight lines (see [5]). Therefore Corollary 1 shows that the answer to the above problem is affirmative when the universal covering  $\Omega$  of  $M$  is irreducible (note that  $\Omega$  is necessarily homogeneous) and not affinely equivalent to a convex cone.

#### REFERENCES

- [ 1 ] H. Asano, On the irreducibility of homogeneous convex cones, *J. Fac. Sci. Univ. Tokyo*, **15** (1968), 201–208.
- [ 2 ] S. Kaneyuki, Homogeneous bounded domains and Siegel domains, *Lect. Notes in Math.*, **241**, Springer, 1971.
- [ 3 ] S. Kobayashi, *Hyperbolic Manifolds and Holomorphic Mappings*, Marcel Dekker, New York, 1970.
- [ 4 ] —, Intrinsic distances associated with flat affine or projective structures, *J. Fac. Sci. Univ. Tokyo*, **24** (1977), 129–135.
- [ 5 ] —, Projectively invariant distances for affine and projective structures, in *Differential Geometry*, Banach Center Publications, Vol. 12, PWN-Polish Scientific Publishers, Warsaw, 1983, 127–152.
- [ 6 ] — and K. Nomizu, *Foundations of Differential Geometry*, Vol. II, Wiley-Interscience, New York, 1968.
- [ 7 ] O. Rothaus, The construction of homogeneous convex cones, *Ann. of Math.*, **83** (1966), 358–376.
- [ 8 ] H. Shima, Homogeneous convex domains of negative sectional curvature, *J. Differential Geom.*, **12** (1977), 327–332.
- [ 9 ] T. Tsuji, On the group of isometries of an affine homogeneous convex domain, *Hokkaido Math. J.*, **13** (1984), 31–50.
- [10] —, The irreducibility of an affine homogeneous convex domain, *Tohoku Math. J.*, **36** (1984), 203–216.

- [11] E. B. Vinberg, The theory of convex homogeneous cones, *Trans. Moscow Math. Soc.*, **12** (1963), 340–403.

*Mathematical Institute  
Tohoku University  
Sendai, 980  
Japan*