

## MOLECULAR ABUNDANCES IN THE SGR A MOLECULAR CLOUD

Y. C. MINH<sup>1</sup>, W. M. IRVINE<sup>2</sup>, and P. FRIBERG<sup>3,4</sup>

<sup>1</sup> *Inst. of Space Sci. and Astronomy, Daejon 305-348, Korea*

<sup>2</sup> *FCRAO, Univ. of Massachusetts, Amherst, MA 01003, USA*

<sup>3</sup> *Onsala Space Observatory, S-439 00 Onsala, Sweden*

<sup>4</sup> *Joint Astronomy Centre, Hilo, HI 96720, USA*

**ABSTRACT.** We have obtained column densities for  $\text{HCO}^+$ ,  $\text{HCO}$ ,  $\text{HCS}^+$ ,  $\text{C}_3\text{H}_2$ ,  $\text{HC}_5\text{N}$ ,  $\text{SiO}$ ,  $\text{OCS}$ ,  $\text{HCOOH}$ ,  $\text{CH}_3\text{CH}_2\text{OH}$ , and  $\text{CH}_3\text{CCH}$  toward Sgr A. The fractional abundance of  $\text{SiO}$  relative to molecular hydrogen in Sgr A is comparable to that for the Orion plateau,  $\sim 10^{-7} - 10^{-8}$ , which may be a typical value for hot clouds. The abundances of  $\text{HCO}$ ,  $\text{CH}_3\text{CH}_2\text{OH}$  and  $\text{CH}_3\text{CCH}$  all appear to be enhanced relative to other molecular clouds such as Sgr B2.

The molecular abundances and chemistry of the Sgr A molecular cloud have not been as well characterized as those of Sgr B2, but the strong activities and shocks of the Galactic center could affect the clouds in Sgr A more efficiently because of their greater proximity. This may result in a unique chemistry of the Sgr A clouds, such as has been suggested from observations of  $\text{HCO}_2^+$  (Minh et al. 1991a; Paper 1).

Data were obtained in 1988 June with the Swedish-ESO 15 m telescope in Chile. Telescope parameters and observing method were included in Paper 1. Observed molecules, transitions, and rest frequencies are listed in Table 1. We have obtained data for the clouds observed in  $\text{NH}_3$  (Gusten et al. 1981) and in  $\text{HCO}_2^+$  (Paper 1). Column densities were determined assuming optically thin emission and an apparent background radiation temperature of 10 K (cf. Paper 1 and Minh et al. 1991b).

In Figure 1 we plot the fractional abundances relative to  $\text{H}_2$  for the molecules observed toward M-0.13-0.08, and also those for TMC-1<sup>2</sup> and Sgr B2, and for Orion(KL) from the tabulations of Irvine et al. (1987), and Blake et al. (1987), respectively, for comparison. The fractional abundance of  $\text{SiO}$  at Sgr A is derived to be  $\sim 10^{-7} - 10^{-8}$  relative to molecular hydrogen which is similar to that of the Orion plateau. The high  $\text{SiO}$  abundance could be explained by high-temperature or shock chemistry (Ziurys et al. 1989). It is also possible, however, that an enhanced abundance of elemental Si comes from the disruption of silicate grains by shocks in the Galactic center region, which can lead naturally to an enhanced  $\text{SiO}$  abundance.

TABLE 1. Observed molecules.

Molecule (Trans.)	Rest Frequency (GHz)
$H^{13}CO^+$ (1-0)	86.75429
$HC^{18}O^+$ (1-0)	85.16226
$HCO$ ( $1_{01}-0_{00}^a$ )	86.67082
$HCS^+$ (2-1)	85.34790
$C_3H_2$ ( $2_{12}-1_{01}$ )	85.33889
$HC_5N$ (32-31)	85.20135
$SiO$ (2-1)	86.84700
$^{29}SiO$ (2-1)	85.75913
$OCS$ (7-6)	85.13911
$HCOOH$ ( $4_{14}-3_{13}$ )	86.54613
$CH_3CH_2OH$ ( $6_{06}-5_{15}$ )	85.26547
$CH_3CCH$ (J=5-4 K=0)	85.45730
$CH_3CCH$ (J=6-5 K=0)	102.54798
$CH_3OH$ ( $3_1-4_0^A$ )	107.01385

<sup>a</sup>For the (3/2-1/2 2-1) trans.




Figure. 1 Abundances relative to molecular hydrogen on a logarithmic scale for Sgr A, Sgr B2, TMC-1 and the Orion extended ridge and the plateau. Data for Sgr B2 and TMC-1 from Irvine et al. (1987), and for Orion(KL) from Blake et al. (1987).

The fractional abundances of several molecules observed here, in particular  $HCO$ ,  $CH_3CH_2OH$ , and  $CH_3CCH$ , appear to be enhanced relative to values for other sources (Figure 1). It is interesting that the production of  $CH_3CH_2OH$  and  $CH_3CCH$  probably involves relatively hydrogenated species such as  $C_2H_4$  or  $CH_4$  (Millar et al. 1991; Millar & Freeman 1984); this might suggest the influence of grain chemistry or high temperature reactions.

We conclude that a rich chemistry exists in Sgr A, which could partly be a result of the energetic processes of the Galactic center region, such as shocks, UV radiation, and also the possible interaction of the neutral and the ionized gas around the nucleus.

- Blake G.A., Sutton E.C., Masson C.R., Phillips T.G., 1987, *ApJ* 315, 621  
 Gusten R., Walmsley C.M. Pauls T., 1981, *A&A* 103, 197  
 Irvine W.M., Goldsmith P.F., Hjalmanson A., 1987, *Interstellar Processes* eds. D.J. Hollenbach, H.A. Thronson, Jr., D.Reidel, p. 561  
 Millar T.J., Freeman A., 1984, *MNRAS* 207, 405  
 Millar T.J., Herbst E., Charnley B., 1991, *ApJ* 369, 147  
 Minh Y.C., Brewer M.K., Irvine W.M., Friberg P., Johansson L.E.B., 1991a, *A&A* 244, 470 ; Paper 1  
 Minh Y.C., Irvine W.M., Friberg P., 1991b, submitted to *A&A*  
 Ziurys L.M., Friberg P., Irvine W.M., 1989, *ApJ* 343, 201