

Anthropogenic food: an emerging threat to polar bears

TOM S. SMITH, ANDREW E. DEROCHER, RACHEL L. MAZUR, GEOFF YORK
MEGAN A. OWEN, MARTYN OBBARD, EVAN S. RICHARDSON and STEVEN C. AMSTRUP

Abstract Supplemental food from anthropogenic sources is a source of conflict with humans for many wildlife species. Food-seeking behaviours by black bears *Ursus americanus* and brown bears *Ursus arctos* can lead to property damage, human injury and mortality of the offending bears. Such conflicts are a well-known conservation management issue wherever people live in bear habitats. In contrast, the use of anthropogenic foods by the polar bear *Ursus maritimus* is less common historically but is a growing conservation and management issue across the Arctic. Here we present six case studies that illustrate how negative food-related interactions between humans and polar bears can become either chronic or ephemeral and unpredictable. Our examination suggests that attractants are an increasing problem, exacerbated by climate change-driven sea-ice losses that cause increased use of terrestrial habitats by bears. Growing human populations and increased human visitation increase the likelihood of human–polar bear conflict. Efforts to reduce food conditioning in polar bears include attractant management, proactive planning and adequate resources for northern communities to reduce conflicts and improve human safety. Permanent removal of unsecured sources of nutrition, to reduce food conditioning, should begin immediately at the local level as this will help to reduce polar bear mortality.

Keywords Anthropogenic food, Arctic, human–bear conflict, polar bear, *Ursus maritimus*

Introduction

Globally, human–wildlife conflict is an increasing threat to species conservation. Wildlife conflicts include crop damage, livestock depredation and human injury (Naughton-Treves, 1998; Madden, 2004; Acharya et al., 2016; Nyhus, 2016). With climate warming-driven shifts in resource availability and species distributions there is an increased potential for human–wildlife conflict (LeDee et al., 2020). However, addressing such conflict is a complex process that involves many stakeholders with various perspectives and objectives. Therefore, the approaches taken to reduce conflicts differ widely (Dickman, 2010; White & Ward, 2010).

Most conflicts with wildlife are associated with habitat loss or modification, changes in resource availability, increasing human populations, incursion into wildlife habitats and the provision of anthropogenic food that can supplement energy input (Nyhus, 2016; Plaza & Lambertucci, 2017). Supplemental food can have both positive and negative effects on wildlife, including changes in distribution, body condition, survival, reproduction and population dynamics (Lunn & Stirling, 1985; Stringham, 1989; Becker et al., 2015; Lillie et al., 2019). For example, polar bears feeding on subsistence-harvested bowhead whale *Balaena mysticetus* carcasses ingest low-fat and high-protein foods that could result in increases in energetic expenditures and lead to health problems (Rode et al., 2021). Although anthropogenic food can benefit wildlife, it often brings animals near to humans and could result in conflict. Management is particularly challenging when access to anthropogenic food results in conflicts with carnivores because of the combination of both perceived and real threats to people and their property (Naughton-Treves et al., 2003; Treves & Karanth, 2003; Morehouse & Boyce, 2017).

Conflicts between people and bears have occurred wherever and whenever their distributions overlap (Rajpurohit & Krausman, 2000; Dai et al., 2020; Krofel et al., 2020) and often because of bears being attracted to anthropogenic foods (Lunn & Stirling, 1985; Craighead et al., 1995). Unsecured food and food waste can lead food-

TOM S. SMITH (Corresponding author, [ORCID iD](https://orcid.org/0000-0001-8384-2454), tom_smith@byu.edu) Wildlife and Wildlands Conservation Program, Brigham Young University, Provo, USA

ANDREW E. DEROCHER ([ORCID iD](https://orcid.org/0000-0002-1104-7774)) Department of Biological Sciences, University of Alberta, Edmonton, Canada

RACHEL L. MAZUR ([ORCID iD](https://orcid.org/0000-0002-8050-7543)) Inventory and Monitoring Division, Olympic National Park, Port Angeles, USA

GEOFF YORK ([ORCID iD](https://orcid.org/0000-0001-9963-3921)) and STEVEN C. AMSTRUP* ([ORCID iD](https://orcid.org/0000-0001-8030-3906)) Polar Bears International, Bozeman, USA

MEGAN A. OWEN ([ORCID iD](https://orcid.org/0000-0003-1655-3972)) San Diego Zoo Wildlife Alliance, Escondido, USA

MARTYN OBBARD ([ORCID iD](https://orcid.org/0000-0003-2064-0155)) Ontario Ministry of Natural Resources and Forestry, DNA Building, Trent University, Peterborough, Canada

EVAN S. RICHARDSON ([ORCID iD](https://orcid.org/0000-0003-3172-5592)) Wildlife Research Division, Science and Technology Branch, Environment and Climate Change Canada, Winnipeg, Canada

*Also at: Department of Zoology and Physiology, University of Wyoming, Laramie, USA

Received 8 November 2021. Revision requested 24 January 2022.

Accepted 21 February 2022. First published online 20 July 2022.

conditioned bears to associate humans with food rewards (Herrero et al., 2005), resulting in property damage, human injury and death (Herrero, 2018).

The relationship between black bears *Ursus americanus* and brown bears *Ursus arctos* (including grizzly bears) and anthropogenic attractants is well documented in the continental USA (Wilson et al., 2005; Mazur, 2010). When the US government established a national park and associated visitor facilities, dumps were built to handle waste. Bears then began foraging at these anthropogenic food sources (Craighead et al., 1995). As bears became food-conditioned, conflicts escalated, resulting in human injuries and property damage (Craighead et al., 1995; Mazur & Seher, 2008; Herrero, 2018). Attempts to control human–bear conflict were often conducted in the absence of an understanding of the underlying causal factors (Haroldson et al., 2008; Elfström et al., 2014). Although public education, hazing, relocation and waste control actions were implemented, abrupt dump closures resulted in increased conflicts as food-conditioned bears sought alternative sources of anthropogenic foods. In Yellowstone National Park, dump closures resulted in a marked increase in brown bear mortality and an associated population decline (Craighead et al., 1995; Haroldson et al., 2008). Once habituated, food-conditioned bears might damage property and injure people to obtain food (McCarthy & Seavoy, 1994; Mazur & Seher, 2008; Elfström et al., 2014; Lillie et al., 2018). Additional challenges arise from parent–offspring learning in bears that can result in anthropogenic food use in new generations (Lunn & Stirling, 1985; Mazur & Seher, 2008). Ultimately, preventing bears from gaining access to anthropogenic food is necessary to reduce conflicts and associated bear mortalities.

Because of their low population density and dependence on remote sea-ice habitats, conflicts between people and polar bears *Ursus maritimus* have been less frequent than those between people and black and brown bears (Krofchuk et al., 2020). Although less studied, human–polar bear conflicts have a long association with anthropogenic foods (Lunn & Stirling, 1985; Dyck, 2006; Rogers et al., 2015). Polar bears on land are often food-deprived, and sea-ice loss driven by climate warming has lengthened the durations that polar bears spend on land (Castro de la Guardia et al., 2013; Stern & Laidre, 2016). Longer ice-free seasons increase the energetic stresses on fasting polar bears (Castro de la Guardia et al., 2013; Hamilton et al., 2014; Molnár et al., 2020) and can lead to more polar bears seeking food near human settlements (Stirling & Derocher, 1993; Towns et al., 2009; Heemsink et al., 2020).

Here we examine case studies of polar bear attraction to anthropogenic food and its role as a source of human–bear conflict. We conclude with a consideration of the conservation and management implications of anthropogenic food reliance.

Case studies

Polar bears are distributed across the Arctic in 19 subpopulations within the jurisdictions of five countries (the USA, Canada, Greenland, Norway and Russia; IUCN/SSC Polar Bear Specialist Group, 2018) and are categorized as Vulnerable on the IUCN Red List (Wiig et al., 2015). We examine human–polar bear conflicts associated with anthropogenic food sources in six areas with a range of social and environmental conditions (Fig. 1).

Churchill, Manitoba, Canada

The western Hudson Bay polar bear subpopulation spends most of the year on the annual sea ice of Hudson Bay. When the ice melts in summer, the entire subpopulation is forced onto land. As sea ice reforms in autumn, polar bears migrate northward. This seasonal phenomenon brings them close to the town of Churchill (population c. 900 people) and often results in human–polar bear conflicts (Towns et al., 2009).

Polar bear attraction and use of anthropogenic food increased when a military base and a beluga *Delphinapterus leucas* whaling station were established in 1942, increasing food attractants in and near Churchill (Stirling et al., 1977). Use of a dump near the town was dominated by subadults and family groups (Lunn & Stirling, 1985). By the 1960s food-conditioned polar bears at Churchill became a public safety concern as polar bears damaged property and inflicted injuries on people whilst in search of anthropogenic food beyond the dump (Fleck & Herrero, 1988; Struzik, 2014). Although supplemental food from the dump was deemed to have minimal survival or reproductive benefit, polar bears returned to the dump habitually every year (Lunn & Stirling, 1985). This study also noted that in Nunavut, the jurisdiction north of Manitoba, polar bears that used the Churchill dump were twice as likely to be hunted by Inuit, for subsistence, compared to polar bears that did not use the dump.

Government-sponsored programmes dedicated to dealing with Churchill's growing polar bear problem (Polar Bear Control Program in 1969 and Polar Bear Alert Program after 1984) were initiated to ensure human safety and the protection of property (Kearney, 1989). Changes to polar bear management policies in the 1980s resulted in an effort to reduce the number of polar bears using the dump habitually, through hazing, relocation, temporary housing of polar bears at a holding facility until the Hudson Bay froze, transportation to zoos or other captive facilities, and lethal removal (Towns et al., 2009; Struzik, 2014; Heemsink et al., 2020). The Churchill town dump was closed in 2005 (Towns et al., 2009) and there was no notable increase in the number of human–polar bear conflicts near Churchill. However, ascribing cause and effect to these various actions is not possible because of a concurrent decline in subpopulation abundance and changing




FIG. 1 Locations of polar bear visits to dumps (Table 1) to exploit anthropogenic food sources. The case study locations are indicated in bold.

management practices, such as dump closures (Heemskerk et al., 2020).

The Churchill experience with dump closure resulted in markedly different outcomes from those seen with the dump closures in Yellowstone National Park. In Churchill, no increase in human–polar bear conflicts occurred post-dump closure because a well-developed plan and infrastructure were in place to address such conflicts, the most problematic food-conditioned polar bears had been removed before dump closure and the subpopulation was declining in abundance because of climate warming. Closing the dump, securing waste within a polar bear-proof facility and public education decreased damage and human injury.

Arviat, Nunavut, Canada

Human–polar bear conflict has increased in recent years in some Nunavut communities. This has been attributed to increased human presence on the land and increased polar bear abundance; polar bears have spent more time onshore because of delayed sea-ice formation driven by climate warming (Dowsley & Wenzel, 2008). Arviat is a coastal community of western Hudson Bay (population c. 3,000 people) in Nunavut, Canada. Polar bears in this area are part of the western Hudson Bay subpopulation, which declined from c. 1,200 to c. 800 polar bears during 1987–2011 (Lunn et al., 2016). Despite this decline in abundance,

human–polar bear conflicts have increased over the past several decades.

Arviat has an open dump that provides anthropogenic food for polar bears. As a result, the number of polar bears using the dump has grown each year since record keeping began in the 1960s (Savikataaq, 2014). A deterrent programme instituted in 2010 has reduced the destruction of problem polar bears but the dump remains a significant attractant. Delayed sea-ice formation in Hudson Bay (Castro de la Guardia et al., 2017) has led to reduced body condition of polar bears (Stirling et al., 1999; Scullo et al., 2016) and could be responsible for the increased use of the dump as polar bears migrate northwards along the western coast (Towns et al., 2009). This case study illustrates how a subpopulation of polar bears that is declining in abundance because of the loss of sea-ice habitat could change movement patterns in response to human food waste, resulting in increased human–polar bear conflicts where few have occurred historically.

Belushya Guba, Novaya Zemlya, Russia

In February 2019, Novaya Zemlya, a remote Russian archipelago between the Barents Sea and the Kara Sea, experienced a ‘mass invasion’ of polar bears (TASS Russian News Agency, 2019). Reports indicated that 52 polar bears had been drawn to a dump near the village of Belushya Guba (population c. 2,000 people). For 2 weeks, dozens of

TABLE 1 Locations, type of food and dates of polar bears *Ursus maritimus* visiting dumps across the Arctic.

Location (by country)	Type of food	Date	Source
Canada			
Churchill, Manitoba	Waste dump	Ongoing	Lunn & Stirling (1985)
Paulatuk, Northwest Territories	Waste dump	2000s	Joint Secretariat (2015)
Tuktoyaktuk, Northwest Territories	Waste dump	c. 2014	Joint Secretariat (2015)
Arviat, Nunavut	Waste dump	9 Apr. 2009	Toth (2019)
Naujuut, Nunavut	Field Camp	Aug. 2018	Canadian Broadcasting Corporation (2018)
Rankin Inlet, Nunavut	Waste dump	Undated, recent	Canadian Wildlife Service (2009)
Resolute Bay, Nunavut	Waste dump	2003	Dyck (2006)
Kashechewan, Ontario	Waste dump	July 2016	Canadian Broadcasting Corporation (2016)
Moose Factory, Ontario	Waste dump	Dec. 2015	Montgomery (2017)
Makkovik, Québec	Waste dump	Feb. 2019	Canadian Broadcasting Corporation (2019)
Norway			
Gåshamna, Svalbard	Field Camp	Aug. 1998	D. Vongraven, pers. comm. (2022)
Russia			
Belushya Guba, Arkhangelsk	Waste dump	27 Nov. 2019	Nilsen (2019)
Leningradsky, Chukotka	Waste dump	May 2020	Ryabikova (2020)
Ryrkaypiy, Chukotka	Waste dump	Dec. 2019	Odynova (2019)
Egvekinot, Chukotka	Dump truck	Mar. 2016	Hooper (2016)
Norilsk, Krasnoyarsk Krai	Waste dump	June 2019	Bever (2019)
Bely Island, Yamalo-Nenets Autonomous Okrug	Rubbish container	Dec. 2019	Siberian Times (2019)
Vilkitskiy Island, Yamalo-Nenets Autonomous Okrug	Rubbish container	Dec. 2019	Siberian Times (2019)
USA			
Kaktovik, Alaska	Whale remains	Ongoing	Miller et al. (2006)

polar bears congregated at the open dump. Some bears entered the village, attempting to gain access to homes and civic buildings. Consequently, local Russian authorities declared a state of emergency on 16 February 2019 that lasted 10 days (TASS Russian News Agency, 2019). Although polar bears had visited the dump in previous years, this event was unprecedented and likely the result of a lengthened on-shore fasting period because of the loss of sea ice in the region. From the perspective of human–polar bear conflict, the situation at Belushya Guba illustrates that changes in the level of polar bear use of anthropogenic food can occur rapidly. There is concern that it is probable large numbers of polar bears will return to the area in the future unless access to anthropogenic food sources is eliminated.

Ryrkaypiy, Chukotka, Russia

In early December 2019, 60 polar bears congregated at a dump near the village of Ryrkaypiy (population c. 600 people) in Russia's Chukotka region (Odynova, 2019). Ryrkaypiy is 0.8 km from the Chukchi Sea. The open dump is a few hundred metres from the village. Polar

bears had unrestricted access to the dump and congregated there until sea-ice formation occurred in late autumn. Both young and old polar bears were observed at the dump, and all were reported to be of below-average body condition. Delayed sea-ice formation and poor body condition probably account for polar bear use of this dump (Odynova, 2019). These polar bears also entered the village, damaged property and threatened inhabitants. Consequently, all public events were cancelled, children were not allowed outside and polar bear monitors worked to keep the polar bears out of the village. Once the sea ice formed and thickened, polar bears returned to the Chukchi Sea. This case illustrates that sea-ice conditions and polar bear body condition are predictors of human–polar bear conflicts and, as in other areas, dumps are an attraction to polar bears and can increase the numbers of problem animals in settlements.

Kaktovik, Alaska, USA

Located on the North Slope of Alaska, Kaktovik is a predominantly Inupiat community (population c. 180 people), with a history of subsistence harvest of bowhead whales. The remains of butchered bowhead whale carcasses left on

beaches where butchering has occurred are a rich food source for a portion of the Southern Beaufort Sea polar bear subpopulation, ranging from minor dietary contributions (Boucher et al., 2019) to 15% of their diet (Bourque et al., 2020). Approximately 16% of the Southern Beaufort Sea polar bear subpopulation visits whale carcasses annually at Kaktovik (Lillie et al., 2019).

Polar bears that were once drawn to the now protected Kaktovik community dump are attracted by bowhead whale remains following hunter harvesting. As many as 90 polar bears drawn from up to 160 km away gather near Kaktovik each autumn to feed on whale carcasses. A similar situation occurs in Barrow, Alaska (Herreman & Peacock, 2013), and this is probably related to increased land use by the Southern Beaufort Sea polar bear subpopulation and the availability of bowhead whale remains from subsistence hunts (Miller et al., 2006). As in other examples where human-mediated food creates an attraction, whale remains represent a significant source of energy to polar bears, yet polar bears feeding on whale remains could suffer negative health effects (Rode et al., 2021) and they often enter the community, damage property and threaten people. Persistent and potentially dangerous polar bears that have become accustomed to entering the community are sometimes euthanized because of public safety concerns. A growing concern is the development of polar bear viewing tourism in the town. Such tourism provides an economic incentive to retain whale carcasses near town, presenting management challenges. This case study illustrates the challenges facing northern communities where traditional harvesting activities interact with climate warming-driven shifts in polar bear distributions and with their increased reliance on anthropogenic food.

First Nations coastal communities in Ontario, Canada

The Southern Hudson Bay subpopulation of polar bears lives on the sea ice of south-eastern Hudson Bay and James Bay to the south. However, during the ice-free season these polar bears move onto islands and coastal areas of James Bay and the Hudson Bay coast of Ontario. First Nations communities (Fort Severn, Peawanuck, Kashechewan, Attawapiskat and Moosonee/Moose Factory, population c. 250–3,500 people in each) are located along the western coast of James Bay and near the Hudson Bay coast. Human–polar bear conflicts related to attractants occur periodically at seasonal camps and communities in these areas (Lemelin et al., 2010). Spring goose hunting, during which hunters live in small cabins or tent camps along the coast, often results in conflict as this is when bears may come off the ice and approach camps. On shore, polar bears raid caches of harvested geese and are subsequently killed to protect the harvest. In the late summer or autumn before sea ice

forms, some polar bears visit dumps near these communities. Occasionally, polar bears are found south of the area that is typically occupied during the ice-free season (Obbard & Middel, 2012). In July 2016 an adult female polar bear and two yearlings were found in the Kashechewan community dump (Canadian Broadcasting Corporation, 2016). One of the yearlings was shot when the family group entered town. Farther south, in December 2015 a single polar bear was shot at the Moose Factory dump (2 km north of the community) in the interest of public safety (Canadian Broadcasting Corporation, 2015), and in December 2020 an adult female polar bear and two yearlings were translocated from the Moose Factory dump (Moose Cree First Nation, 2020).

Polar bear responses to small-scale attractants

The above case studies illustrate the relationship between polar bears and large-scale attractants in the form of unsecured dumps or supplemental food. Dumps affect polar bears by providing an abundant, spatially and temporally predictable source of nutrition that is often exploited during lengthy periods when natural food sources are not available. However, human–polar bear conflict has a long and often unrecorded history involving much smaller-scale, ephemeral attractants such as those at hunting camps in Ontario. Additional examples include camps supporting resource extraction (e.g. oil and gas), tourism and research activities. Even in communities with secure dumps, polar bears could be attracted to animal carcasses stored outside homes, cooking odours, other human-related scents and areas where dogs are yarded. The following cases highlight situations in which polar bears were attracted to small-scale attractants, with one resulting in two polar bears being killed and the other resulting in a human fatality along with two polar bear fatalities.

On 5 July 1998, near Gåshamna in Hornsund, Svalbard, a group of 17 ecotourists and researchers established a field camp. On 8 August 1998, a polar bear described as appearing hungry and aggressive was observed near the cooking tent used by the group. The bear approached to within 20 m of people and was shot and killed. On the following day a second polar bear was observed near the camp's rubbish burning site. When this bear rapidly approached two people, it was shot at a distance of 6–10 m. Attempts to deter these polar bears with noise makers and warning gunshots failed. A necropsy found rubbish in the stomach of one of the polar bears.

In August 2018, on White Island near the town of Naujaat, Nunavut, three subsistence hunters were approached by a female polar bear and her yearling at their camp (Rogers, 2018). The adult female attacked and killed one person before the others, who had been lightly injured,

could kill both bears. These polar bears were reported to have been in good condition (Rogers, 2018). It is possible, but uncertain, that anthropogenic scents (e.g. food, harvested animals) at the camp could have attracted these polar bears.

These cases highlight that both human and polar bear fatalities have been, and probably will continue to be, associated with attractants at camps. With increasing tourism in the Arctic (Runge et al., 2020), negative interactions with polar bears are likely to increase.

Discussion

Unsecured dumps and other sources of anthropogenic food can represent sources of nutrition for polar bears when climate warming-driven sea-ice loss makes natural food less available. Polar bears often remain at these sources until they are driven off, access is barred or natural food becomes available. We have shown that human conflicts with polar bears are often associated with attractants. Although human-inflicted mortality of bears drawn to anthropogenic wastes is a management concern, disease transmission rates are also a concern because bears congregate in close proximity to each other and to other scavenging species when drawn to dumps. Little is known about disease transmission rates amongst congregating bears, but transmission rates are elevated where bears congregate at anthropogenic food sources (Atwood et al., 2017; Whiteman et al., 2018). There are numerous examples of other species experiencing higher rates of disease transmission and outbreak when congregated at anthropogenic food sources (Sorensen et al., 2014).

Food obtained from dumps is a poor substitute for the bears' natural diet (Plaza & Lambertucci, 2017; Rode et al., 2021). Polar bears cannot subsist entirely on terrestrial food because it lacks adequate levels of fat (Rode et al., 2015a), and although food ingested at dumps was found to increase polar bear body mass when compared to polar bears that did not use dumps, reproductive and survival rates did not increase (Lunn & Stirling, 1985). As indiscriminate foragers, polar bears often consume non-food waste contaminated with food residues. This includes plastics (e.g. bread bags, nappies, food waste in rubbish bags), metals (e.g. aluminium foils with food residues, tin cans, batteries), wood (e.g. toothpicks, skewers) and other materials that either contain food (e.g. ceramics) or were coated with food. Instances of polar bears experiencing serious health issues or dying because of ingested materials at dumps have been reported (Lunn & Stirling, 1985; Dickie, 2019; A.E. Derocher, pers. obs., 2020; R. Stimmelmayr, pers. obs., 2020).

Polar bears that feed on anthropogenic food have higher mortality rates than those that do not (Lunn & Stirling, 1985; McCarthy & Seavoy, 1994). Although not all food-conditioned bears attack humans, individuals that do attack

are often food-conditioned (Herrero, 2018). When feeding on anthropogenic food, a link between human scent and anthropogenic food is established. Upon entering a camp and detecting a human scent, those same polar bears occasionally enter tents in search of the food they expect to be present (Herrero, 2018). Such polar bears are often shot either by campers or by the authorities. Translocation is generally not an option for food-conditioned bears because they often resume their search for anthropogenic food when released (Hopkins & Kalinowski, 2013). For these reasons, the management of anthropogenic scents (i.e. attractant security) should be a high priority throughout the range of polar bears.

In summary, we have presented the following concerns associated with the use of anthropogenic food by polar bears: (1) Various forms of anthropogenic food can exacerbate human–polar bear conflict by attracting polar bears to come into close contact with people. (2) If anthropogenic food becomes unavailable or is reduced, food-conditioned polar bears may seek alternative sources, leading them to investigate nearby settlements. (3) Polar bears consuming anthropogenic food may be exposed to harmful chemicals and pollutants, could ingest non-food items or could be exposed to novel diseases or parasites. (4) Feeding on anthropogenic food disrupts the natural movement and distribution of polar bears. (5) Polar bears seeking anthropogenic food may form unnatural aggregations, with negative consequences for both polar bears and people. (6) Variations in the size, temporal, and spatial dynamics of attractants necessitates diverse management strategies. (7) Affected communities are small and often lack the resources to manage anthropogenic attractants.

Problems of food conditioning in polar bears have been reported in all five polar bear range states (the USA, Russia, Canada, Greenland and Norway; Wilder et al., 2017; Fig. 1, Table 1). As polar bears spend increasing amounts of time on land as a result of climate warming-driven sea-ice loss, the likelihood of them seeking anthropogenic food increases (Stirling & Derocher, 1993; Derocher et al., 2004). Along with decreasing body reserves and reduced survival as a result of polar bears spending less time on ice (Molnár et al., 2020), human–polar bear conflict may become an increasing source of polar bear mortality if not addressed. Unless anthropogenic sources of energy are secured, this could represent an increasing threat to the viability of some polar bear subpopulations and decrease the safety of people living in affected regions.

Although the overarching threat from climate warming-driven habitat loss will take many years to resolve, managing anthropogenic waste should be addressed now. Work with black and brown bears has shown that once access to anthropogenic food is eliminated, bears return to their natural feeding strategies (Baruch-Mordo et al., 2014; Lewis et al., 2015) and human–bear conflicts decrease.

We do not need innovative solutions to the problem of anthropogenic waste attracting polar bears, but rather an immediate and concerted effort throughout the Arctic to apply what is already known. Education, the implementation of polar bear-proof methods of waste storage, law enforcement and the provision of adequate resources at the community level are required to mitigate this potentially increasing problem. All communities noted here were small, and thus have limited resources to deal with human–polar bear conflict. In addition, the human population is increasing across the Arctic. For example, the population in Nunavut, Canada, is projected to grow by 31% from 2014 to 2035 (Heleniak, 2020). Therefore, waste will increase, as will the potential for human–polar bear conflict associated with anthropogenic food.

Historically, the attraction of polar bears to anthropogenic food has been uncommon because of infrequent overlap of polar bear feeding areas with human settlements. Although polar bears are usually born on land, they spend most of their lives far from people, on sea ice, where they feed on marine mammals that provide an energy-rich diet (Stirling & McEwan, 1975; DeMaster & Stirling, 1981). Polar bears are largely driven ashore as a result of the annual cycle of melting sea ice or in search of maternal denning habitats (Knudsen, 1978; Latour, 1981), during which time they use up their fat stores as a source of energy (Derocher et al., 1990; Ramsay et al., 1991). In general, polar bears do not continue to use anthropogenic food once sea ice becomes available again (Lunn & Stirling, 1985).

As the global climate warms, the temporal and spatial overlap between polar bears and people is increasing. Reductions in both sea-ice habitat and prey availability are predicted to become more prevalent across broader reaches of the Arctic (Stirling & Derocher, 2012). Consequently, polar bears will be drawn to anthropogenic food in more locations across the Arctic and for longer periods, thus threatening their survival and human safety (Rode et al., 2015b). This growing conflict will be most prominent in locations where human settlements already overlap with areas where polar bears wait for sea-ice formation. In some regions of their range, polar bears are now spending more time waiting on shore for the sea ice to freeze than before 2000 (Castro de la Guardia et al., 2017). Consequently, polar bears are now spending more time searching for alternative food sources while land-bound. Naturally occurring terrestrial foods, such as beached whale carcasses, bird eggs and fish, are too limited in distribution and abundance to substitute for the usual marine mammal diet of polar bears (Rode et al., 2015b). Therefore, polar bears will ultimately be drawn into settlements where nutrient-dense anthropogenic food is often readily available. This is particularly true in areas where once-small settlements and associated dumps are now expanding (Odynova, 2019). The unprecedented polar bear incursion into Novaya

Zemlya underscores how quickly a shift in resource use and dependence can arise. Because these situations have high potential for human–polar bear conflict, resulting in both human injury and the removal of polar bears from declining subpopulations, the negative implications of anthropogenic food for polar bear conservation is a growing concern.

We recommend that localities implement the following actions to address this growing problem: (1) better document incidents of food-seeking behaviour (i.e. monitor the number of polar bears seeking food at dumps and other sources), (2) examine the patterns and timing of food-seeking behaviour, (3) document the age and sex composition of food-seeking polar bears, (4) test methods for keeping food from polar bears, (5) assess the effectiveness of deterrence methods used on food-seeking polar bears, (6) assess the efficacy of relocation methods (e.g. helicopter relocation), (7) examine recidivism rates amongst food-seeking polar bears (i.e. food conditioning), and (8) develop a means of predicting when polar bears could seek anthropogenic food (e.g. sea-ice break-up, ice formation, body condition).

Acknowledgements This paper developed from a meeting in Churchill, Manitoba, in autumn 2019 where the issue of dump use by polar bears arose. We thank Dan Cox for suggesting exploration of this issue and Polar Bears International for arranging this meeting. This study did not receive funding from any agency.

Author contributions All authors contributed to the various components of this work.

Conflicts of interest None.

Ethical standards No animals were handled during this study. Research conducted in this paper complies with all ethical standards in the jurisdictions of the authors and abided by the *Oryx* guidelines on ethical standards.

References

- ACHARYA, K.P., PAUDEL, P.K., NEUPANE, P.R. & KOHL, M. (2016) Human–wildlife conflicts in Nepal: patterns of human fatalities and injuries caused by large mammals. *PLOS ONE*, 11, e0161717.
- ATWOOD, T.C., DUNCAN, C., PATYK, K.A., NOL, P., RHYAN, J., MCCOLLUM, M. et al. (2017) Environmental and behavioral changes may influence the exposure of an Arctic apex predator to pathogens and contaminants. *Scientific Reports*, 7, 13193.
- BARUCH-MORDO, S., WILSON, K.R., LEWIS, D.L., BRODERICK, J., MAO, J.S. & BRECK, S.W. (2014) Stochasticity in natural forage production affects use of urban areas by black bears: implications to management of human–bear conflicts. *PLOS ONE*, 9, e85122.
- BECKER, D.J., STREICKER, D.G. & ALTIZER, S. (2015) Linking anthropogenic resources to wildlife–pathogen dynamics: a review and meta-analysis. *Ecology Letters*, 18, 483–495.
- BEVER, L. (2019) A starving polar bear wandered into a Russian city, scavenging for food. *The Washington Post*, 19 June 2019. washingtonpost.com/world/2019/06/19/starving-polar-bear-wandered-into-city-scavenging-food-hundreds-miles-home [accessed 22 January 2022].

- BOUCHER, N.P., DEROCHER, A.E. & RICHARDSON, E.S. (2019) Space use patterns affect stable isotopes of polar bears (*Ursus maritimus*) in the Beaufort Sea. *Polar Biology*, 42, 1581–1593.
- BOURQUE, J., ATWOOD, T.C., DIVOKY, G.J., STEWART, C. & MCKINNEY, M.A. (2020) Fatty acid-based diet estimates suggest ringed seal remain the main prey of southern Beaufort Sea polar bears despite recent use of onshore food resources. *Ecology and Evolution*, 10, 2093–2103.
- CANADIAN BROADCASTING CORPORATION (2015) Polar bear found feeding at Moose Factory dump shot dead. *CBC News*, 22 December 2015. cbc.ca/news/canada/sudbury/polar-bear-moose-factory-1-3376268 [accessed 5 June 2021].
- CANADIAN BROADCASTING CORPORATION (2016) Polar bear shot after wandering through Kashechewan. *CBC News*, 21 July 2016. cbc.ca/news/canada/sudbury/polar-bear-cub-shot-kashechewan-1-3688025 [accessed 22 January 2022].
- CANADIAN BROADCASTING CORPORATION (2018) Survivor recalls deadly encounter with polar bears. *CBC News*, 29 August 2018. cbc.ca/news/canada/north/polar-bear-kimmirut-nunavut-death-1-4803481 [accessed 22 January 2022].
- CANADIAN BROADCASTING CORPORATION (2019) Polar bear captured near Makkovik dump, 2nd still roaming the area. *CBC News*, 1 February 2019. cbc.ca/news/canada/newfoundland-labrador/polar-bears-makkovik-warning-1.5002244 [accessed 22 January 2022].
- CANADIAN WILDLIFE SERVICE (2009) *Consultations on the Proposed Listing of the Polar Bear as Special Concern under the Species at Risk Act*. Conducted February–April 2009. Canadian Wildlife Service, Iqaluit, Nunavut.
- CASTRO DE LA GUARDIA, L., DEROCHER, A.E., MYERS, P.G., TERWISSCHA VAN SCHELTINGA, A.D. & LUNN, N.J. (2013) Future sea ice conditions in western Hudson Bay and consequences for polar bears in the 21st century. *Global Change Biology*, 19, 2675–2687.
- CASTRO DE LA GUARDIA, L., MYERS, P.G., DEROCHER, A.E., LUNN, N.J. & TERWISSCHA VAN SCHELTINGA, A.D. (2017) Sea ice cycle in western Hudson Bay, Canada, from a polar bear perspective. *Marine Ecology Progress Series*, 564, 225–233.
- CRAIGHEAD, J.J., SUMMER, J.S. & MITCHELL, J.A. (1995) *The Grizzly Bears of Yellowstone: Their Ecology in the Yellowstone Ecosystem, 1959–1992*. Island Press, Washington, DC, USA.
- DAI, Y.C., HACKER, C.E., ZHANG, Y.G., LI, Y., LI, J., XUE, Y.D. & LI, D.Q. (2020) Conflicts of human with the Tibetan brown bear (*Ursus arctos pruinosus*) in the Sanjiangyuan region, China. *Global Ecology and Conservation*, 22, e01039.
- DEMASTER, D.P. & STIRLING, I. (1981) *Ursus maritimus*. *Mammalian Species*, 145, 1–7.
- DEROCHER, A.E., LUNN, N.J. & STIRLING, I. (2004) Polar bears in a warming climate. *Integrative and Comparative Biology*, 44, 163–176.
- DEROCHER, A.E., NELSON, R.A., STIRLING, I. & RAMSAY, M.A. (1990) Effects of fasting and feeding on serum urea and creatinine levels in polar bears. *Marine Mammal Science*, 6, 196–203.
- DICKIE, G. (2019) Polar bear's plastic diets are a growing problem. *Hakai Magazine*, 5 April 2019. hakaimagazine.com/news/polar-bears-plastic-diets-a-growing-problem [accessed 28 January 2022].
- DICKMAN, A.J. (2010) Complexities of conflict: the importance of considering social factors for effectively resolving human–wildlife conflict. *Animal Conservation*, 13, 458–466.
- DOWSLEY, M. & WENZEL, G. (2008) 'The time of the most polar bears': a co-management conflict in Nunavut. *Arctic*, 61, 177–189.
- DYCK, M.G. (2006) Characteristics of polar bears killed in defense of life and property in Nunavut, Canada, 1970–2000. *Ursus*, 17, 52–62.
- ELFSTRÖM, M., DAVEY, M.L., ZEDROSSER, A., MÜLLER, M., DE BARBA, M., STØEN, O.-G. et al. (2014) Do Scandinavian brown bears approach settlements to obtain high-quality food? *Biological Conservation*, 178, 128–135.
- FLECK, S. & HERRERO, S. (1988) *Polar Bear–Human Conflicts*. Contract report for Parks Canada and GNWT, contract 502/85/23. Parks Canada, Calgary, Canada.
- HAMILTON, S.G., CASTRO DE LA GUARDIA, L., DEROCHER, A.E., SAHANATIEN, V., TREMBLAY, B. & HUARD, D. (2014) Projected polar bear sea ice habitat in the Canadian Arctic archipelago. *PLOS ONE*, 9, e113746.
- HAROLDSON, M.A., SCHWARTZ, C.C. & GUNTHER, K.A. (2008) Grizzly bears in the greater Yellowstone ecosystem: from garbage, controversy, and decline to recovery. *Yellowstone Science*, 16, 13–24.
- HEEMSKERK, S., JOHNSON, A.C., HEDMAN, D., TRIM, V., LUNN, N.J., MCGEACHY, D. & DEROCHER, A.E. (2020) Temporal dynamics of human–polar bear conflicts in Churchill, Manitoba. *Global Ecology and Conservation*, 24, e01320.
- HELENIAK, T. (2020) The future of the Arctic populations. *Polar Geography*, 44, 136–155.
- HERREMAN, J. & PEACOCK, E. (2013) Polar bear use of a persistent food subsidy: insights from non-invasive genetic sampling in Alaska. *Ursus*, 24, 148–163.
- HERRERO, S. (2018) *Bear Attacks: Their Causes and Avoidance*, 3rd edition. Globe Pequot Press, Guilford, USA.
- HERRERO, S., SMITH, T., DEBRUYN, T.D., GUNTHER, K. & MATT, C.A. (2005) Brown bear habituation to people – safety, risks, and benefits. *Wildlife Society Bulletin*, 33, 362–373.
- HOOPER, B. (2016) Polar bear and cubs make dump truck their personal playground. *UPI*, 21 March 2016. upi.com/Odd_News/2016/03/21/Polar-bear-and-cubs-make-dump-truck-their-personal-playground/8391458583503 [accessed 22 January 2022].
- HOPKINS, J.B. & KALINOWSKI, S.T. (2013) The fate of transported American black bears in Yosemite National Park. *Ursus*, 24, 120–126.
- IUCN/SSC POLAR BEAR SPECIALIST GROUP (2018) 2016 status report on the world's polar bear subpopulations. In *Polar Bears: Proceedings of the 18th Working Meeting of the IUCN/SSC Polar Bear Specialist Group*, 7–11 June 2016, Anchorage, Alaska, USA (eds G.M. Durner, K.L. Laidre & G.S. York), pp. 1–21. IUCN, Gland, Switzerland and Cambridge, UK.
- JOINT SECRETARIAT (2015) *Inuvialuit and Nunuq: A Polar Bear Traditional Knowledge Study*. Joint Secretariat, Inuvialuit Settlement Region, Inuvik, Canada.
- KEARNEY, S.R. (1989) The polar bear alert program at Churchill, Manitoba. In *Bear–People Conflicts: Proceedings of a Symposium on Management Strategies* (ed. M. Bromley), pp. 83–92. Northwest Territories Department of Renewable Resources, Yellowknife, Northwest Territories, Canada.
- KNUDSEN, B. (1978) Time budgets of polar bears (*Ursus maritimus*) on North Twin Island, James Bay, during summer. *Canadian Journal of Zoology*, 56, 1627–1628.
- KROFEL, M., ELFSTRÖM, M., AMBARLI, H., BOMBieri, G., GONZÁLEZ-BERNARDO, E., JERINA, K. et al. (2020) Human–bear conflicts at the beginning of the twenty-first century: patterns, determinants, and mitigation measures. In *Bears of the World: Ecology, Conservation and Management* (eds M. Melletti & V. Penteriani), pp. 213–226. Cambridge University Press, Cambridge, UK.
- LATOUR, P.B. (1981) Spatial relationships and behavior of polar bears (*Ursus maritimus* Phipps) concentrated on land during the ice-free season of Hudson Bay. *Canadian Journal of Zoology*, 59, 1763–1774.
- LEDEE, O.E., HANDLER, S.D., HOVING, C.L., SWANSTON, C.W. & ZUCKERBERG, B. (2020) Preparing wildlife for climate change: how far have we come? *Journal of Wildlife Management*, 85, 7–16.
- LEMELIN, R.H., DOWSLEY, M., WALMARK, B., SIEBEL, F., BIRD, L., HUNTER, G. et al. (2010) Wabusk of the Omushkegouk: Cree–polar bear (*Ursus maritimus*) interactions in northern Ontario. *Human Ecology*, 38, 803–815.

- LEWIS, D.L., BARUCH-MORDO, S., WILSON, K.R., BRECK, S.W., MAO, J.S. & BRODERICK, J. (2015) Foraging ecology of black bears in urban environments: guidance for human–bear conflict mitigation. *Ecosphere*, 6, 1–18.
- LILLIE, K.M., GESE, E.M., ATWOOD, T.C. & CONNER, M.M. (2019) Use of subsistence-harvested whale carcasses by polar bears in the southern Beaufort Sea. *Arctic*, 72, 404–412.
- LILLIE, K.M., GESE, E.M., ATWOOD, T.C. & SONSTHAGEN, S.A. (2018) Development of on-shore behavior among polar bears (*Ursus maritimus*) in the southern Beaufort Sea: inherited or learned? *Ecology and Evolution*, 8, 7790–7799.
- LUNN, N.J. & STIRLING, I. (1985) The significance of supplemental food to polar bears during the ice-free period of Hudson Bay. *Canadian Journal of Zoology*, 63, 2291–2297.
- LUNN, N.J., SERVANTY, S., REGEHR, E.V., CONVERSE, S.J., RICHARDSON, E. & STIRLING, I. (2016) Demography of an apex predator at the edge of its range: impacts of changing sea ice on polar bears in Hudson Bay. *Ecological Applications*, 26, 1302–1320.
- MADDEN, F. (2004) Creating coexistence between humans and wildlife: global perspectives on local efforts to address human–wildlife conflict. *Human Dimensions of Wildlife*, 9, 247–257.
- MAZUR, R.L. (2010) Does aversive conditioning reduce human–black bear conflict? *Journal of Wildlife Management*, 74, 48–54.
- MAZUR, R.L. & SEHER, V. (2008) Socially learned foraging behaviour in wild black bears, *Ursus americanus*. *Animal Behaviour*, 75, 1503–1508.
- MCCARTHY, T.M. & SEAOY, R.J. (1994) Reducing non-sport losses attributable to food conditioning: human and bear behavior modification in an urban environment. *Bears: Their Biology and Management*, 9, 75–84.
- MILLER, S., SCHLIEBE, S. & PROFFITT, K. (2006) *Demographics and Behavior of Polar Bears Feeding on Bowhead Whale Carcasses at Barter and Cross Islands, Alaska*. OCS Study MMS 2006-14. Report by U.S. Fish & Wildlife Service for Minerals Management Service. Marine Mammals Management Office, Anchorage, USA.
- MOLNÁR, P.K., BITZ, C.M., HOLLAND, M.M., KAY, J.E., PENK, S.R. & AMSTRUP, S.C. (2020) Fasting season length sets temporal limits for global polar bear persistence. *Nature Climate Change*, 10, 732–738.
- MONTGOMERY, M. (2017) Polar bears, climate change, a new problem for some northern communities. *Radio Canada International*, 16 March 2017. rcinet.ca/en/2017/03/16/polar-bears-climate-change-a-new-problem-for-some-northern-communities [accessed 22 January 2022].
- MOOSE CREE FIRST NATION (2020) Polar bear updates. moosecree.com/polar-bear-update [accessed 5 June 2020].
- MOREHOUSE, A.T. & BOYCE, M.S. (2017) Troublemaking carnivores: conflicts with humans in a diverse assemblage of large carnivores. *Ecology and Society*, 22, 4.
- NAUGHTON-TREVES, L. (1998) Predicting patterns of crop damage by wildlife around Kibale National Park, Uganda. *Conservation Biology*, 12, 156–168.
- NAUGHTON-TREVES, L., GROSSBERG, R. & TREVES, A. (2003) Paying for tolerance: rural citizens' attitudes toward wolf depredation and compensation. *Conservation Biology*, 17, 1500–1511.
- NILSEN, T. (2019) Plastic waste on polar bears' menu a growing problem. *The Barents Observer*, 27 November 2019. thebarentsobserver.com/en/ecology/2019/11/plastic-waste-polar-bears-menu-growing-problem [accessed 22 January 2022].
- NYHUS, P.J. (2016) Human–wildlife conflict and coexistence. *Annual Review of Environment and Resources*, 41, 143–171.
- OBBARD, M.E. & MIDDEL, K.R. (2012) Bounding the southern Hudson Bay polar bear subpopulation. *Ursus*, 23, 134–144.
- ODYNNOVA, A. (2019) Dozens of hungry polar bears are causing problems for a Russian village. *CBS News*, 6 December 2019. cbsnews.com/news/polar-bears-ryrkaypiy-russia-evacuation-suggested-in-chukotka-village-today-2019-12-06 [accessed 21 January 2020].
- PLAZA, P.I. & LAMBERTUCCI, S.A. (2017) How are garbage dumps impacting vertebrate demography, health, and conservation? *Global Ecology and Conservation*, 12, 9–20.
- RAJPURHIT, K.S. & KRAUSMAN, P.R. (2000) Human–sloth–bear conflicts in Madhya Pradesh, India. *Wildlife Society Bulletin*, 28, 393–399.
- RAMSAY, M.A., NELSON, R.A. & STIRLING, I. (1991) Seasonal changes in the ratio of serum urea to creatinine in feeding and fasting polar bears. *Canadian Journal of Zoology*, 69, 298–302.
- RODE, K.D., ROBBINS, C.T., NELSON, L. & AMSTRUP, S.C. (2015a) Can polar bears use terrestrial foods to offset lost ice-based hunting opportunities? *Frontiers in Ecology and the Environment*, 13, 138–145.
- RODE, K.D., ROBBINS, C.T., STRICKER, C.A., TARAS, B.D. & TOLLEFSON, T.N. (2021) Energetic and health effects of protein overconsumption constrain dietary adaptation in an apex predator. *Scientific Reports*, 11, 15309.
- RODE, K.D., WILSON, R.R., REGEHR, E.V., ST MARTIN, M., DOUGLAS, D.C. & OLSON, J. (2015b) Increased land use by Chukchi Sea polar bears in relation to changing sea ice conditions. *PLOS ONE*, 10, e0142213.
- ROGERS, M.C., PEACOCK, E., SIMAC, K., O'DELL, M.B. & WELKER, J.M. (2015) Diet of female polar bears in the southern Beaufort Sea of Alaska: evidence for an emerging alternative foraging strategy in response to environmental change. *Polar Biology*, 38, 1035–1047.
- ROGERS, S. (2018) Polar bears that killed Nunavut hunter were healthy, GN says. *Nunatsiaq News*, 6 September 2018. nunatsiaq.com/stories/article/65674polar_bears_involved_in_naujaat_attack_were_all_healthy_gn_says [accessed 28 January 2022].
- RUNGE, C.A., DAIGLE, R.M. & HAUSNER, V.H. (2020) Quantifying tourism booms and the increasing footprint in the Arctic with social media data. *PLOS ONE*, 15, e0227189.
- RYABIKOVA, V. (2020) How Russians deal with over-curious bears. *Russia Beyond*, 3 June 2020. rbth.com/lifestyle/332280-how-russians-deal-with-over-curious-bears [accessed 22 January 2022].
- SAVIKATAAQ, J.J. (2014) *Operation Arviat Polar Bear Summary Report*. Arviat, Nunavut, Canada.
- SCIULLO, L., THIEMANN, G.W. & LUNN, N.J. (2016) Comparative assessment of metrics for monitoring the body condition of polar bears in western Hudson Bay. *Journal of Zoology*, 300, 45–58.
- SIBERIAN TIMES (2019) T-34 polar bear video explained: scientists marked the predator in 'safe paint'. *The Siberian Times*, 7 December 2019. siberiantimes.com/other/others/news/t-34-polar-bear-video-explained-scientists-marked-the-predator-in-safe-paint [accessed 22 January 2022].
- SORENSEN, A., VAN BEEST, F.M. & BROOK, R.K. (2014) Impacts of wildlife baiting and supplemental feeding on infectious disease transmission risk: a synthesis of knowledge. *Preventive Veterinary Medicine*, 113, 356–363.
- STERN, H.L. & LAIDRE, K.L. (2016) Sea-ice indicators of polar bear habitat. *The Cryosphere*, 10, 1–15.
- STIRLING, I. & DEROCHER, A.E. (1993) Possible impacts of climatic warming on polar bears. *Arctic*, 46, 240–245.
- STIRLING, I. & DEROCHER, A.E. (2012) Effects of climate warming on polar bears: a review of the evidence. *Global Change Biology*, 18, 2694–2706.
- STIRLING, I. & MCEWAN, E.H. (1975) The calorific value of whole ringed seals (*Phoca hispida*) in relation to polar bear (*Ursus*

- maritimus*) ecology and hunting behavior. *Canadian Journal of Zoology*, 53, 1021–1027.
- STIRLING, I., JONKEL, C., SMITH, P., ROBERTSON, R. & CROSS, D. (1977) The ecology of the polar bear (*Ursus maritimus*) along the western coast of Hudson Bay. *Canadian Wildlife Service Occasional Paper*, 33, 1–64.
- STIRLING, I., LUNN, N.J. & IACOZZA, J. (1999) Long-term trends in the population ecology of polar bears in western Hudson Bay in relation to climatic change. *Arctic*, 52, 294–306.
- STRINGHAM, S.F. (1989) Demographic consequences of bears eating garbage in dumps: an overview. In *Bear-People Conflicts: Proceedings of a Symposium on Management Strategies* (ed. M. Bromley), pp. 35–42. Northwest Territories Department of Renewable Resources, Yellowknife, Canada.
- STRUZIK, E. (2014) *Arctic Icons: How the Town of Churchill Learned to Love its Polar Bears*. Fitzhenry & Whiteside, Markham, Canada.
- TASS RUSSIAN NEWS AGENCY (2019) Emergency declared in Novaya Zemlya archipelago over polar bear ‘invasion’. TASS, 9 February 2019. tass.com/emergencies/1043985 [accessed 20 January 2021].
- TOTH, K. (2019) Polar bears in North are getting into more garbage. *CBC News*, 9 April 2019. cbc.ca/news/canada/north/plastics-in-polar-bear-stomachs-1.5089982 [accessed 25 January 2022].
- TOWNS, L., DEROCHER, A.E., STIRLING, I., LUNN, N.J. & HEDMAN, D. (2009) Spatial and temporal patterns of problem polar bears in Churchill, Manitoba. *Polar Biology*, 32, 1529–1537.
- TREVES, A. & KARANTH, K.U. (2003) Human–carnivore conflict and perspectives on carnivore management worldwide. *Conservation Biology*, 17, 1491–1499.
- VON STACKELBERG, M. (2016) Polar bear shot after wandering through Kashechewan. *CBC News*, 21 July 2016. cbc.ca/news/canada/sudbury/polar-bear-cub-shot-kashechewan-1.3688025 [accessed 5 June 2021].
- WHITE, P.C.L. & WARD, A.I. (2010) Interdisciplinary approaches for the management of existing and emerging human–wildlife conflicts. *Wildlife Research*, 37, 623–629.
- WHITEMAN, J.P., HARLOW, H.J., DURNER, G.M., REGEHR, E.M., AMSTRUP, S.C. & BEN-DAVID, M. (2018) Heightened immune system function in polar bears using terrestrial habitats. *Physiological and Biochemical Zoology*, 92, 1–11.
- WIIG, Ø., AMSTRUP, S., ATWOOD, T., LAIDRE, K., LUNN, N., OBBARD, M. et al. (2015) *Ursus maritimus*. In *The IUCN Red List of Threatened Species* 2015. dx.doi.org/10.2305/IUCN.UK.2015-4.RLTS.T22823A14871490.en.
- WILDER, J.M., VONGRAVEN, D., ATWOOD, T., HANSEN, B., JESSEN, A., KOCHNEV, A. et al. (2017) Polar bear attacks on humans: implications of a changing climate. *Wildlife Society Bulletin*, 41, 537–547.
- WILSON, S.M., MADEL, M.J., MATTSON, D.J., GRAHAM, J.M., BURCHFIELD, J.A. & BELSKY, J.M. (2005) Natural landscape features, human-related attractants, and conflict hotspots: a spatial analysis of human–grizzly bear conflicts. *Ursus*, 16, 117–129.