

# Jansky VLA S-band view of H $\alpha$ emitters (HAEs) associated with a protocluster 4C23.56 at $z = 2.5$

Minju Lee<sup>1,2</sup>, Kenta Suzuki<sup>3</sup>, Kotaro Kohno<sup>3</sup>, Yoichi Tamura<sup>3</sup>,  
Daisuke Iono<sup>2</sup>, Bunyo Hatsukade<sup>2</sup>, Kouichiro Nakanishi<sup>2</sup>,  
Ichi Tanaka<sup>2</sup>, Tadayuki Kodama<sup>2</sup>, Kenichi Tadaki<sup>2</sup>, Soh Ikarashi<sup>2,4</sup>,  
Junko Ueda<sup>2</sup>, Hideki Umehata<sup>3</sup>, Toshiki Saito<sup>1,2</sup> and Ryohei Kawabe<sup>2</sup>

<sup>1</sup>Department of Astronomy, The University of Tokyo,  
7-3-1 Hongo, Bunkyo-ku, Tokyo 133-0033, Japan

<sup>2</sup>National Observatory of Japan, 2-21-1 Osawa,  
Mitaka, Tokyo 181-0015, Japan  
email: minju.lee@nao.ac.jp

<sup>3</sup>Institute of Astronomy, The University of Tokyo,  
2-21-1 Osawa, Mitaka, Tokyo, 181-0015, Japan

<sup>4</sup>European Southern Observatory, Karl-Schwarzschild-Str. 2,  
D-85748 Garching, Germany

**Abstract.** We present recent results on Karl Jansky Very Large Array (JVLA) deep S-band (2-4 GHz) observation towards a protocluster 4C23.56 at redshift  $z \sim 2.5$ . The protocluster 4C23.56 is known to have a significant over density ( $\sim 5$  times) of star-burst galaxies selected to be H $\alpha$  line-bright by a Subaru narrow band imaging. Now we have found 25 HAEs associated with the protocluster. These starburst HAEs are likely to become massive ellipticals at  $z = 0$  in a cluster. Various other galaxy populations also reside in this field and the fact makes the field very unique as a tool to understand galaxy formation in a over dense region. Subsequent deep 1100- $\mu$ m continuum surveys by the ASTE 10-m dish have discovered that several submillimeter bright galaxies (SMGs) coincide with HAEs, suggesting HAEs undergoing dusty starbursts. As star formation rates (SFRs) of HAEs might have been underestimated, we use radio being resistant to dust extinction. We investigate the correlation between  $\text{SFR}_{1.4\text{GHz}}$  and  $\text{SFR}_{\text{H}\alpha}$  for radio index  $\alpha = 0.8$  to see if the correlation holds for the sources and to check the number of dusty star forming galaxies. Our final results will allow us to evaluate quantitatively how the galaxy formation channel may be different under the condition of over-densities.

**Keywords.** galaxies: active – galaxies: clusters: general – surveys – galaxies: evolution – galaxies: formation – galaxies: high-redshift – radio continuum: galaxies – galaxies: starburst

---

## 1. Introduction

Clusters of galaxies, being the most massive structures bounded with gravity in the Universe, provide us an exclusive window to investigate the galaxy formation and evolution in the over-dense region. The progenitors of clusters, protoclusters, are unsettled systems having over-densities than the fields, which is becoming present-day clusters, perhaps by merging and intense star formation, although only a handful to them are identified up to now. A protocluster 4C23.56 is at redshift  $z = 2.48$ , where a group of HAEs are found from the Subaru narrow band (NB) survey (MAHALO; Mapping H $\alpha$  and Lines of Oxygen with Subaru). The total number of the detected HAEs is 25 up to now (Tanaka *et al.*, in prep; Tanaka *et al.* (2011)). The protocluster is very unique in that it has various galaxy populations in addition to HAEs, e.g. distant red


galaxies (DRGs; Kajisawa *et al.* (2006)), extremely red objects (EROs; Knopp & Chambers (1997)), mid-infrared sources (Galametz *et al.* (2012), Mayo *et al.* (2012)) and that there are multi-wavelength ancillary data sets available. Sub-mm single dish telescope ASTE/AzTEC revealed that four of SMGs were overlapped with some HAEs (Suzuki *et al.* in prep; Zeballos *et al.* in prep). However, we could barely pin down the counterparts of SMGs at other wavelengths due to the coarse resolution with ASTE compared to other optical/NIR surveys as well as their redshifts. The overlap of SMGs suggests that dusty star bursts may be onset in HAEs making even the  $H\alpha$  emissions be diminished, leading the underestimation of SFRs. So, we targeted a protocluster associated to the radio galaxy 4C23.56 with Karl Jansky Very Large Array (JVLA) at S-band (2-4 GHz).

## 2. Results and conclusions

We have reached the r.m.s. level to detect HAEs having  $SFR > 100 - 400 M_{\odot} \text{yr}^{-1}$  ( $3\sigma$ ) and, as a result, the detection number of JVLA counterparts was seven out of 25. We have also applied stacking analysis to reveal the averaged properties of undetected sources but no the features detected at  $3\sigma$ . The figure 1 shows the correlation between 1.4 GHz radio continuum and  $H\alpha$  emission which was derived from JVLA 3 GHz (this work) and Subaru NB observations where we assumed radio spectral index  $\alpha = 0.8$  to convert 3 GHz information to 1.4 GHz luminosity. We speculate five HAEs among the rest are undergoing dusty starbursts, of which SFRs have been underestimated. Otherwise, these would represent AGN. Adding the Spitzer/MIPS  $24\mu\text{m}$  data and PdBI CO(5-4) observation, one HAE is likely to be undergoing a merger with a heavily obscured starburst. Differentiating the (dusty) starburst and the AGN for the rest will be remained as future works by doing SED fittings and comparison with X-ray observation (Lee *et al.* in prep). Further follow-up observations with ALMA/JVLA will provide more information on the physical properties on each galaxy in this protocluster and illuminate the processes how massive galaxy formation and evolution is affected by its surrounding environment.

## References

- Galametz, A., Stern, D., De Breuck, C., *et al.* 2012, *ApJ*, 749, 169  
 Knopp, G. P. & Chambers, K. C. 1997, *ApJS*, 109, 367  
 Kajisawa, M., Kodama, T., Tanaka, I., Yamada, T., & Bower, R. 2006, *MNRAS*, 371, 577  
 Mayo, J. H., Vernet, J., De Breuck, C., *et al.* 2012, *A&A*, 539, A33  
 Tanaka, I., Breuck, C. D., Kurk, J. D., *et al.* 2011, *PASJ*, 63, 415



**Figure 1.** Radio and  $H\alpha$  emission correlation of a protocluster associated with 4C23.56. The detection rate of HAE counterparts in radio is  $\sim 30\%$ . The radio galaxy 4C23.56 itself is on top right of this figure and well-above the correlation representing a radio excess by AGN. All JVLA detections of HAE counterparts, except for one is above the correlation which suggests dusty starbursts of HAEs and/or AGN.