

# GEMS: The destiny of Blue Spheroidal Galaxies

Boris Häußler<sup>1</sup>, Eric F. Bell<sup>1</sup>, Marco Barden<sup>1</sup>, Daniel H. McIntosh<sup>2</sup>,  
Hans-Walter Rix<sup>1</sup>, Andrea Borch<sup>3</sup>, Steven V. W. Beckwith<sup>4,5</sup>,  
John A. R. Caldwell<sup>6</sup>, Catherine Heymans<sup>7</sup>, Knud Jahnke<sup>1</sup>,  
Shardha Jogee<sup>8</sup>, Sergey E. Koposov<sup>1</sup>, Klaus Meisenheimer<sup>1</sup>,  
Chien Y. Peng<sup>4</sup>, Sebastian F. Sánchez<sup>9</sup>, Rachel S. Somerville<sup>1</sup>,  
Lutz Wisotzki<sup>10</sup>, and Christian Wolf<sup>11</sup>

<sup>1</sup>Max-Planck-Institut für Astronomie, Königstuhl 17, 69117, Heidelberg, Germany

<sup>2</sup>Department of Astronomy, University of Massachusetts, 710 North Pleasant Street, Amherst, MA 01003, USA

<sup>3</sup>Astronomisches Recheninstitut, Mönchhofstraße 12-14, 69120, Heidelberg, Germany

<sup>4</sup>Space Telescope Science Institute, 3700 San Martin Dr., Baltimore, MD 21218, USA

<sup>5</sup>Johns Hopkins University, 3400 North Charles Street, Baltimore, MD 21218, USA

<sup>6</sup>University of Texas, McDonald Observatory, Fort Davis, TX 79734, USA

<sup>7</sup>Department of Physics and Astronomy, The University of British Columbia, 6224 Agricultural Road Vancouver, V6T 1Z1 Canada

<sup>8</sup>University of Texas at Austin, 1, University Station C1400, Austin, TX 78712-0259, USA

<sup>9</sup>Centro Astronomico Hispano Aleman de Calar Alto, C/Jesus Durban Remon 2-2, Almeria, E-04004, Spain

<sup>10</sup>Universität Potsdam, Am Neuen Palais 10, 14469, Potsdam, Germany

<sup>11</sup>Department of Physics, Denys Wilkinson Bldg., University of Oxford, Keble Road, Oxford, OX1 3RH, UK

email: boris@mpia.de

One of the key predictions of hierarchical galaxy formation models is that a significant fraction of elliptical galaxies form in late merging events. One of the most important diagnostics of such an assembly is the existence of blue spheroidal galaxies, which have spheroid-dominated morphologies and blue colors indicating recent star formation, as an intermediate step in the evolution of elliptical galaxies.

We present results from the GEMS (Rix *et al.* 2004) survey showing the properties of these galaxies derived from 2-D galaxy fitting using GALFIT (Peng *et al.* 2002) of the  $\sim 8000$  galaxies with photometric redshifts in the 28x28 HST mosaic. For the first time we were able to divide the observed population of blue spheroidal galaxies into sub-populations of different stellar masses.

We find that massive blue spheroidals are likely to be the progenitors of red sequence galaxies. In contrast, low-mass blue spheroidals have half-light radii considerably in excess of those measured for low-mass present day elliptical galaxies; instead, they have larger sizes similar to present-day disk-dominated systems with substantial bulges.

## References

- Peng, C.Y., Ho, L.C., Impey, C.D., & Rix, H., 2002, *AJ*, 124:266-293.  
Rix, H., Barden, M., Beckwith, S.V.W., Bell, E.F., Caldwell, J.A.R., Häußler, B., Jahnke, K., Jogee, S., McIntosh, D.H., Meisenheimer, K., Peng, C.Y., Sanchez, S.F., Somerville, R.S., Wisotzki, L., & Wolf, C., 2004, *ApJS*, 152:163.