

Effects of a high-protein, low-carbohydrate v. high-protein, moderate-carbohydrate weight-loss diet on antioxidant status, endothelial markers and plasma indices of the cardiometabolic profile

Alexandra M. Johnstone^{1*}, Gerald E. Lobley¹, Graham W. Horgan², David M. Bremner¹, Claire L. Fyfe¹, Philip C. Morrice¹ and Garry G. Duthie¹

¹University of Aberdeen, Rowett Institute of Nutrition and Health, Aberdeen AB21 9SB, Scotland, UK

²Biomathematics and Statistics Scotland, Rowett Institute of Nutrition and Health, Aberdeen AB21 9SB, Scotland, UK

(Received 23 July 2010 – Revised 17 December 2010 – Accepted 20 December 2010 – First published online 27 April 2011)

Abstract

There are concerns that weight-loss (WL) diets based on very low carbohydrate (LC) intake have a negative impact on antioxidant status and biomarkers of cardiovascular and metabolic health. Obese men (n 16) participated in a randomised, cross-over design diet trial, with food provided daily, at approximately 8.3 MJ/d (approximately 70% of energy maintenance requirements). They were provided with two high-protein diets (30% of energy), each for a 4-week period, involving a LC (4% carbohydrate) and a moderate carbohydrate (MC, 35% carbohydrate) content. Body weight was measured daily, and weekly blood samples were collected. On average, subjects lost 6.75 and 4.32 kg of weight on the LC and MC diets, respectively ($P < 0.001$, SED 0.350). Although the LC and MC diets were associated with a small reduction in plasma concentrations of retinol, vitamin E (α -tocopherol) and β -cryptoxanthin ($P < 0.005$), these were still above the values indicative of deficiency. Interestingly, plasma vitamin C concentrations increased on consumption of the LC diet ($P < 0.05$). Plasma markers of insulin resistance ($P < 0.001$), lipaemia and inflammation ($P < 0.05$, TNF- α and IL-10) improved similarly on both diets. There was no change in other cardiovascular markers with WL. The present data suggest that a LC WL diet does not impair plasma indices of cardiometabolic health, at least within 4 weeks, in otherwise healthy obese subjects. In general, improvements in metabolic health associated with WL were similar between the LC and MC diets. Antioxidant supplements may be warranted if LC WL diets are consumed for a prolonged period.

Key words: Weight loss: Antioxidant status: Metabolic profile: CVD: High-protein diets

There has been considerable public interest in the use of high-protein, low-carbohydrate (LC) weight-loss (WL) diets, as this type of diet can lead to modest fat loss, at least in the short term^(1,2). Despite widespread use, there are limited data on the possible consequences of this type of restrictive dietary regimen on indices of metabolic dysfunction and CVD risk, such as endothelial function, inflammatory markers and antioxidant status. Specifically, one major nutritional concern is the necessity of a limited intake of fruit and vegetables⁽³⁾ to achieve the LC requirement. This is counter-intuitive to current healthy eating advice, as inadequate intakes of fruits and vegetables are a risk factor for the development of CVD⁽⁴⁾ and certain cancers⁽⁵⁾. This is associated, in part, with a dietary inadequacy of antioxidant micronutrients from plant-based foods. Moreover, it is known that LC intake has an impact on faecal bacterial populations, and this may have implications in the longer term for gut health⁽⁶⁾.

Although the efficacy and safety of high-protein, LC diets⁽⁷⁾ has been questioned, most work has specifically examined the impact of the high protein component on health, for example kidney function⁽⁸⁾ and bone health⁽⁹⁾. There are limited data that directly compare high-protein LC and high-protein, moderate carbohydrate (MC) WL diets on which to make recommendations. For example, it would be of interest to establish whether it is necessary to encourage antioxidant vitamin supplement(s) use while consuming a LC WL diet. Alternatively, recommended daily allowances of vitamins and micronutrients may be met by judicious inclusion of mixed fruits and vegetables, even when LC intake is met. The present study therefore focuses on comparing two iso-energetic high-protein diets, where the fat and carbohydrate contents have been altered. We hypothesised that the LC diet would have a negative effect on markers of antioxidant

Abbreviations: LC, low carbohydrate; M, maintenance; MC, moderate carbohydrate; WL, weight loss.

* **Corresponding author:** Dr A. M. Johnstone, fax +44 1224 716686; email alex.johnstone@abdn.ac.uk

status, and the metabolic and cardiovascular profile in obese men undergoing WL, over a 4-week period.

Subjects and methods

A total of eighteen obese males with a BMI of $>30 \text{ kg/m}^2$ were recruited by newspaper advertisement to participate in a WL diet trial. Thus, the subjects were non-randomly selected individuals who were sufficiently motivated to actively respond to the request for volunteers. Because a subgroup of subjects underwent radioactive positron emission tomography scans (data reported separately), we were ethically constrained to recruit men >50 years, and the final recruitment profile reflects this older age grouping. Women of childbearing age were excluded from the study, on ethical advice, because of the radiation dose. During recruitment, the subjects underwent a medical examination, and their general practitioners were contacted to confirm medical suitability to participate in the study. The present study was conducted according to the guidelines laid down in the Declaration of Helsinki, and all procedures involving human subjects were approved by the NHS North of Scotland Research Ethics Committee. Written informed consent was obtained from all subjects.

Study protocol

Participants were resident in the Human Nutrition Unit at the Rowett Institute of Nutrition and Health, Aberdeen, but were allowed to leave the Unit to attend their workplace. All food and drink were supplied by dietetic staff in the Unit. Subjects consumed a fixed intake maintenance (M) diet supplied at $\text{RMR (MJ/d)} \times 1.5$ for 1 week. They then consumed two WL diets, each for a 4-week period, which provided approximately 70% of energy maintenance requirements as high protein, LC and high protein, MC. The LC diet was designed to be ketogenic, and the MC diet was non-ketogenic. The order of treatments was randomised in a within-subject, cross-over design, whereby half of the subjects started on the LC diet and the other half on the MC diet.

Formulation and preparation of the diets

The composition of each meal, in terms of energy, fat, carbohydrate and protein, was calculated using McCance and Widdowson's *The Composition of Foods* (5th edition) and supplements⁽¹⁰⁾. All meals on the LC diet comprised 30% protein, 4% carbohydrate and 66% fat by energy; while the MC meals contained 30% protein, 35% carbohydrate and 35% fat. All meals within both diets had a fixed energy density of 5.75 MJ/kg. Lunch and dinner meals were fixed intake, with the LC meals containing 47.1 g (800 kJ) protein, 47.7 g (1765 kJ) fat, 6.63 g (106 kJ) carbohydrate and 2667 kJ energy. The MC lunch and dinner meals contained 47.2 g (802 kJ) protein, 25.3 g (936 kJ) fat, 58.5 g (936 kJ) carbohydrate and 2667 kJ energy. Subjects had a choice of seven breakfast meals, cooked on request, which varied from, per meal, 2.46 to 3.00 MJ and 2.31 to 2.92 MJ for the LC and the

MC diets, respectively, with fixed macronutrient composition (%). See Appendices 1 and 2 for menus.

Presentation of the diets and measurement of food intake

While resident at the Human Nutrition Unit, each subject was allocated a refrigerator that was stocked daily with food. Kitchen research staff prepared and weighed all meals daily, with any leftovers weighed to the nearest gram. Breakfasts were eaten in the Unit. Subjects completed food diaries to record the time of eating. Subjects had free access to water and decaffeinated drinks. Energy and nutrient intakes were calculated using the Windiets program (Univation Limited; The Robert Gordon University, Aberdeen, Scotland).

Measurement of anthropometry, blood pressure and body composition

Measurements of body composition and metabolic rate were conducted under standardised conditions, with subjects instructed to fast overnight (10 h). Height was measured at the beginning of the study to the nearest 0.5 cm using a stadiometer (Holtain Limited, Crymych, Dyfed, Wales, UK). Subjects were weighed daily, after voiding, wearing only a pre-weighed dressing gown, to the nearest 50 g on a digital scale (DIGI DS-410; CMS Weighing Equipment Limited, London, UK). Abdominal and gluteal (hip) circumference was measured at the beginning and end of each dietary intervention period, as described previously⁽¹¹⁾, following the guidelines of the International Standards for Anthropometric Assessment. Blood pressure was monitored at the beginning and end of each dietary intervention period, using an automated system (Omron M5-1; Omron Healthcare Incorporation, Bannockburn, IL, USA). Subjects were supine for 10 min before the measurement, and the average of three measures, taken 5 min apart, was recorded. Body composition was calculated by using a four-compartment model⁽¹²⁾ that involved dual-energy X-ray absorptiometry scanning (Norland XR-36, Mark II high-speed pencil beam scanner; Norland Corporation, Fort Atkinson, WI, USA, equipped with dynamic filtration, with version 2.5.2 of Norland software), body density by air displacement whole-body plethysmography (BodPod[®] Body Composition System; Life Measurement Instruments, Concord, CT, USA) and total body water by ^2H dilution⁽¹³⁾.

Blood sampling

Whole blood was sampled from a large antecubital vein in the morning after an overnight fast, before breakfast, using an 18G butterfly needle (Sarstedt, Nuernbrecht, Germany) and an adaptor and collected into separate EDTA and Li heparin tubes. The samples were immediately centrifuged (1000 g at 4°C for 10 min) and divided into aliquots before storage at -80°C . Plasma for vitamin C analysis was diluted with an equal volume of chilled 10% (v/v) metaphosphoric acid before freezing.

Carotenoids and antioxidant vitamins

Plasma vitamin C concentrations were determined by HPLC using an ion pairing technique, and reverse-phase HPLC was used to quantify vitamins A and E (α - and γ -tocopherols) and several carotenoids⁽¹⁴⁾. Analyses were conducted under the round-robin scheme of the National Institute of Standards Micronutrient Measurement Quality Assurance Programme and ISO9001 quality assurance status. Echinone was used as an internal standard, and intra- and inter-specific CV were 2 and 4%, respectively. Analyses were conducted on weekly samples, so data are presented from weeks 1 to 4 for both WL (LC and MC) diets.

Hormone and metabolites

Analysis was conducted at the end of each dietary period, thus data are presented for the three diet periods (M, LC and MC). Insulin was measured on duplicate samples with an ELISA kit (Merckodia, Uppsala, Sweden), with a within-assay CV of 5% and between-assay of 3%. A Kone discrete automated clinical analyser (Kone Oyj, Espoo, Finland) was used for the analysis of urea, glucose, TAG, total cholesterol, LDL and HDL using commercial kits (Labmedics, Salford, Manchester, UK). The homeostasis model assessment of insulin resistance was applied using the fasting glucose and insulin values⁽¹⁵⁾. The bead array Luminex system was employed for the determination of a range of inflammatory markers (IL-1 β , IL-2, IL-6, IL-8, IL-10 and TNF- α), leptin, adhesion molecules (soluble intercellular adhesion molecule-1 and vascular cell adhesion molecule-1) and adiponectin (Millipore, Billerica, MA, USA).

Statistics

Data on energy intake, body weight and composition and blood metabolites for the two WL diets were analysed by ANOVA, with subject and period within subject as random effects, and diet and order as treatment terms (fixed effects). Data on plasma antioxidants were analysed by ANOVA, with subject, period within subject and the week within period as random effects, and diet, order, week and the week \times diet and week \times order interactions as fixed effects. Analyses of all three diets (i.e. including M) were also done, with order terms omitted (as diet M is always first), and antioxidant data for the WL diets averaged across weeks (as diet M was given 1 week only). All analyses were performed using Genstat 8.1 (Lawes Agricultural Trust, VSN International Limited, Hemel Hempstead, Herts, UK).

Results

The baseline characteristics of the sixteen participants are shown in Table 1; two subjects from the original eighteen withdrew for personal reasons.

Body weight and composition

During the M diet, subjects lost, on average, 1.0 kg body weight over the week, so may have been in slight negative

energy balance. During the WL diets, on average, subjects lost 6.75 and 4.32 kg (6 and 4%) body weight, on the LC and MC diets, respectively ($P < 0.001$, SED 0.350). This difference between the diets was explained, in part, by the larger total body water loss on the LC diet (-2.55 v. -0.29 kg; $P = 0.075$; SED 1.175), probably linked to the mobilisation of hepatic glycogen stores. This was reflected in the change in fat-free mass stores with an apparent loss of -2.72 and -0.61 kg on the LC and MC diets, respectively ($P = 0.042$; SED 0.941). There was a gain in protein mass on the LC diet and a loss on the MC diet, but this did not reach significance. There were similar losses in fat mass between the diets with -4.03 and -3.71 kg on the LC and MC diets, respectively ($P = 0.733$; SED 0.919) (Table 2).

Energy and nutrient intake

On average, subjects consumed slightly less food and energy (0.3 MJ) on the LC diet compared with the MC diet ($P < 0.001$), when leftovers were accounted for. They consumed similar amounts of protein (30% of energy) but varying amounts of fat ($P < 0.001$) and carbohydrate ($P < 0.001$). In terms of micronutrient intake, the LC diet contained twice as much retinol ($P < 0.001$) and vitamin E ($P < 0.001$), in comparison with the MC diet, possibly due to the presence of these lipid-soluble vitamins within the higher fat content of the diet. Interestingly, the LC diet also provided more vitamin C ($P < 0.001$) from the high citrus (LC) fruit intake and also vitamin D ($P < 0.001$). The MC diet provided more carotene, in the form of vegetables and tomatoes ($P < 0.001$). All the diets achieved the UK reference nutrient intake for vitamin E (4.0 mg/d), vitamin C (40 mg/d) and vitamin D (2.50 μ g/d) but provided only 50–75% of the reference nutrient intake for vitamin A (700 μ g retinol equivalents for males)⁽¹⁶⁾. Currently, there is no dietary recommendation for carotene intake (as a precursor of vitamin A), but the diets provided 3.7–4.9 mg carotenoids/d, within the range of 3–6 mg β -carotene/d suggested by the Food and Nutrition Board⁽¹⁷⁾ as prudent (Table 3).

Plasma vitamins and carotenoids

Data for values for selected plasma antioxidant vitamins and carotenoids (α -tocopherol, γ -tocopherol, xanthophyll, β -cryptoxanthin, lycopene, α -carotene, β -carotene, vitamin C and retinol) were compared across all three diets and for the

Table 1. Baseline characteristics of the study participants (Mean values, standard deviations and ranges, n 16)

	Mean	SD	Range
Age (years)	55	14	31–74
Height (m)	1.76	0.05	1.68–1.80
Body weight (kg)*	111.9	19.3	86.3–154.8
BMI (kg/m ²)	35.8	5.5	30.0–48.5

* Body weight at the end of maintenance with all subject data pooled, before randomisation to diet treatment.

Table 2. Average measured change in body weight (kg) and composition (kg) on the study diets, where the low carbohydrate (LC) diet is a high-protein, low-carbohydrate diet and the moderate carbohydrate (MC) diet is a high-protein, medium-carbohydrate diet, for the 4-week intervention period*

Body composition (kg)†	LC			MC			P	SED
	Pre	Post	Δ	Pre	Post	Δ		
Body weight	108.8	102.0	-6.75	108.9	104.6	-4.32	<0.001	0.350
Fat mass	41.6	37.6	-4.03	42.4	38.7	-3.71	0.733	0.919
Fat-free mass	67.2	64.4	-2.72	66.5	65.9	-0.61	0.042	0.941
Body fat (%)	37.4	36.0	-1.45	38.0	36.0	-1.95	0.581	0.871
Protein mass	20.4	20.9	0.57	21.3	21.0	-0.26	0.325	0.816
Total body water	48.9	46.4	-2.55	47.8	47.5	-0.29	0.075	1.175

* The data are for the sixteen subjects analysed by ANOVA, where the *P* value refers to the analysis between the weight-loss diets (ΔLC and ΔMC).

† Measured with a four-compartment model⁽¹²⁾.

two WL diets, with the effect of time (week on diet) also considered in Table 4. There were treatment effects of the WL diets compared with baseline (M) for all the antioxidant vitamins ($P < 0.05$). Values were lower on both the WL diets for α -tocopherol ($P < 0.001$), β -cryptoxanthin ($P < 0.001$) and retinol ($P < 0.001$). Relative to the M diet, γ -tocopherol ($P < 0.001$), xanthophyll ($P < 0.001$) and vitamin C ($P < 0.001$) were all greater on the LC diet ($P < 0.05$) but lower on the MC diet ($P < 0.05$). Relative to the M diet, lycopene ($P < 0.001$), α -carotene ($P = 0.002$) and β -carotene ($P = 0.004$) decreased on the LC diet and increased on the MC diet ($P < 0.001$).

There were differences between the two WL diets (LC and MC) for seven of the nine markers measured. For example, values were greater in the LC diet than in the MC diet for γ -tocopherol ($P < 0.001$), xanthophyll ($P = 0.002$) and vitamin C ($P = 0.039$). In contrast, concentrations were higher on the MC diet for lycopene ($P < 0.001$), α -carotene ($P = 0.004$), β -carotene ($P < 0.001$) and retinol ($P < 0.001$). When the effect of week within each diet was compared, there were WL diet \times time interactions for seven of the nine markers. Only vitamin

Table 3. Mean energy and micronutrient intake over the 4 weeks for the sixteen participants, with *P* value and SED for diet differences between the weight loss diets (low carbohydrate (LC) and moderate carbohydrate (MC))

	M*	LC	MC	SED	P
Energy (kJ)	11.49	7.98	8.31	0.007	<0.001
Protein (g)	85.2	137.4	138.9	1.07	0.188
Fat (g)	115.6	143.0	82.2	3.34	<0.001
SFA	39.9	43.7	26.0	0.76	<0.001
PUFA	35.8	24.9	12.0	0.43	<0.001
MUFA	17.3	60.7	34.0	0.67	<0.001
PUFA:SFA	0.89	0.57	0.46	0.57	<0.001
CHO (g)	359.5	22.3	181.3	0.59	<0.001
Starch	177.6	1.95	117.7	2.02	<0.001
Sugar	135.2	19.2	57.1	1.96	<0.001
NSP (g)	21.9	8.72	12.68	0.24	<0.001
Retinol (μ g)	533.4	624.9	325.2	27.8	<0.001
Vitamin E (mg)	7.2	11.9	6.2	0.34	<0.001
Vitamin C (mg)	153.1	148.0	88.4	3.08	<0.001
Carotene (μ g)	3705	3778	4944	98.1	<0.001
Vitamin D (μ g)	3.66	6.43	3.73	0.16	<0.001

M, maintenance; CHO, carbohydrate.

* M diet values provided for reference only.

C and retinol did not show a change in concentration over the 4 weeks, on each WL diet. Notably, for α -tocopherol, γ -tocopherol, β -cryptoxanthin, lycopene and β -carotene, concentrations declined with time on the LC diet but remained constant or increased with the MC diet. In contrast, xanthophyll increased over the 4 weeks with the LC diet but decreased with the MC diet.

Plasma inflammatory markers

Of the cytokines measured (Table 5), only IL-10 ($P = 0.049$) and TNF- α (0.006) showed diet-related effects, with both reduced by WL therapy. The MC WL diet reduced IL-10 to a greater degree than the LC WL diet ($P < 0.05$), but the WL diet composition had no effect on TNF- α .

Plasma cardiovascular risk markers

Mean values for the CVD risk markers adiponectin, soluble intercellular adhesion molecule-1 and soluble vascular cell adhesion molecule-1 in the plasma are presented in Table 6. There was no significant effect of WL or diet composition on any of these markers.

Plasma metabolic health markers

Data on the homeostasis model assessment of insulin resistance, the plasma lipid profile, urea and β -hydroxybutyrate are presented in Table 7, and all parameters changed in response to WL ($P < 0.05$). Glucose was decreased by the WL diet ($P < 0.001$), and more so on the LC diet ($P < 0.001$). Homeostasis model assessment of insulin resistance was reduced during the WL diet, relative to the M diet ($P < 0.001$), but to a similar degree on both the WL diets. This was also reflected in insulin concentration, with a reduction relative to the M diet ($P < 0.001$), but there was no significant WL diet composition effect. As anticipated, the ketone body β -hydroxybutyrate was elevated in the LC (ketogenic) WL diet ($P < 0.001$) but remained similar between the M and MC WL diets. Urea concentration was elevated on both (high-protein) WL diets ($P < 0.001$) relative to the M diet, more so on the LC diet ($P < 0.05$). Lipid profiles were improved with the WL diet, with total cholesterol ($P = 0.02$)

Table 4. Mean values ($\mu\text{g/ml}$) for plasma antioxidants for maintenance (M), low-carbohydrate (LC) and moderate-carbohydrate (MC) weight-loss (WL) diets throughout the trial (weeks 1–4) for the sixteen participants

	M 0	LC				MC				WL diet			
		1	2	3	4	1	2	3	4	Treatment* P	WL diet† P	Week‡ P	SED§
α -Tocopherol ($\mu\text{g/ml}$)	11.46	9.89	9.70	8.91	8.70	9.38	8.75	9.63	8.72	< 0.001	0.396	0.039	0.207
γ -Tocopherol ($\mu\text{g/ml}$)	0.754	0.834	0.890	0.744	0.741	0.563	0.477	0.575	0.460	< 0.001	< 0.001	0.014	0.055
Xanthophyll ($\mu\text{g/ml}$)	0.186	0.187	0.199	0.198	0.207	0.175	0.155	0.165	0.142	< 0.001	0.002	0.020	0.010
β -Cryptoxanthin ($\mu\text{g/ml}$)	0.065	0.048	0.044	0.035	0.030	0.043	0.042	0.042	0.038	< 0.001	0.339	0.010	0.002
Lycopene ($\mu\text{g/ml}$)	0.392	0.321	0.283	0.194	0.176	0.443	0.468	0.526	0.496	< 0.001	< 0.001	< 0.001	0.024
α -Carotene ($\mu\text{g/ml}$)	0.039	0.038	0.036	0.032	0.035	0.041	0.043	0.049	0.047	0.002	0.003	0.020	0.003
β -Carotene ($\mu\text{g/ml}$)	0.135	0.143	0.135	0.123	0.128	0.148	0.149	0.169	0.161	0.004	< 0.001	0.003	0.006
Vitamin C ($\mu\text{mol/l}$)	46.5	50.2	50.9	53.4	52.3	46.8	44.6	43.6	41.7	0.039	0.022	0.177	2.912
Retinol ($\mu\text{g/ml}$)	0.623	0.508	0.461	0.428	0.426	0.553	0.529	0.528	0.489	< 0.001	< 0.001	0.280	0.015

* Treatment effect relates to the P value from comparison of the M and LC and MC diets (average of weeks 1–4).
 † WL diet effect relates to the P value from comparison of the LC and MC diets (average of weeks 1–4).
 ‡ WL diet \times week effect relates to the interaction between the diet (LC and MC) and time course (weeks 1–4).
 § SED is for the WL diet effect.

and LDL ($P=0.009$) reduced, particularly with the MC diet ($P<0.05$), whereas the improvement in HDL-cholesterol ($P<0.001$) was better on the LC diet ($P<0.05$). TAG levels were reduced to similar extents ($P<0.001$) with both the WL diets. Leptin concentration was halved by WL treatment ($P<0.001$), with both diets having a similar effect.

Metabolic health – blood pressure and waist circumference (data not tabulated)

As anticipated, there was a significant effect of the WL diet on waist circumference (cm), relative to the baseline (M) diet ($P<0.001$), with girth reduced from 117.7 to 111.1 and 114.2 cm at the end of the LC and MC WL diets, respectively. The LC diet promoted a greater reduction in girth than the MC diet ($P=0.011$). As a group, the subjects were hypertensive, 144/90 mmHg at baseline (diastolic/systolic), which significantly dropped to a similar degree, ($P<0.001$ for both) by the end of each WL diet regimen (127/79 and 128/79 mmHg) for the LC and MC diets, respectively.

Discussion

The present study compared the impacts of both a high-protein LC WL diet and a high-protein MC WL diet (in obese men) on plasma hormones (insulin), metabolites (glucose, antioxidant vitamins and lipid profile), inflammatory markers (cytokines) and endothelial markers (intercellular adhesion molecule, vascular cell adhesion molecule, adiponectin) as indicators of the metabolic and cardiovascular risk profile. It was hypothesised that the LC diet, which contained only 4% carbohydrate, would have a deleterious impact on vitamin and antioxidant status, which would increase CVD risk. The present data suggest that the LC WL diet used in the present study does not impair plasma indices of cardiometabolic health, at least within a short time period (4 weeks) in otherwise healthy obese subjects. In general, improvements in metabolic health were similar between the LC and MC diets.

Body weight and composition changes

Although subjects consumed a similar amount of food energy, there was a 2.43 kg greater WL on the LC diet. This is similar to a previous study on subjects fed *ad libitum*⁽¹⁾, though, in part, due to carbohydrate restriction depleting hepatic glycogen

Table 5. Effect of the maintenance (M), low-carbohydrate (LC) and medium-carbohydrate (MC) diets on plasma inflammatory markers (pg/ml) in twelve volunteers

	M	LC	MC	SED	P
IL-1 β	6.81	5.23	5.14	0.807	0.089
IL-2	8.41	6.96	6.40	1.185	0.239
IL-6	19.60	19.19	19.17	0.986	0.886
IL-8	5.19	5.55	4.73	0.517	0.299
IL-10	45.7 ^a	36.9 ^{a,b}	28.7 ^b	6.42	0.049
TNF- α	3.29 ^a	2.76 ^b	2.83 ^b	0.161	0.006

^{a,b} Values within rows with unlike superscript letters were significantly different ($P<0.05$).

Table 6. Effect of the maintenance (M), low-carbohydrate (LC) and medium-carbohydrate (MC) diets on plasma cardiovascular risk markers (ng/ml) in twelve volunteers

	M	LC	MC	SED	P
Adiponectin (ng/ml)n	15 683	16 726	14 662	1023.2	0.153
sICAM-1 (ng/ml)	356	280	293	37.2	0.112
sVCAM-1 (ng/ml)	1308	1393	1273	68.2	0.214

sICAM-1, soluble intercellular adhesion molecule-1; sVCAM-1, soluble vascular cell adhesion molecule-1.

stores, with loss of associated water⁽¹⁸⁾. In the four-compartment model of body composition used, glycogen is considered to be part of fat-free mass and is not directly measured. This could account for the greater apparent loss in fat-free mass on the LC diet. Volek *et al.*⁽¹⁹⁾ concluded that LC diets favour the preservation of lean body mass, a finding supported by the present study.

Antioxidant vitamins and carotenoids

To our knowledge, this is the first dietary WL study to report plasma antioxidant values, with previous emphasis on the role of supplementation with the obesity drug Orlistat⁽²⁰⁾ and gastric surgery⁽²¹⁾. The present study found that plasma concentrations of vitamin E, vitamin C and carotenoids on the M diet were similar to that previously reported in a Scottish population⁽²²⁾. The LC diet used in the present study used a wide range of easily obtained fruit and vegetable sources to achieve the 4% carbohydrate (of energy) intake. This wide variety would help provide a number of the vitamin and antioxidant metabolites considered important for metabolic and cardiovascular health⁽²³⁾. These diets were adequate in terms of recommended daily intake for vitamins C, D and E, but deficient in vitamin A. Indeed, plasma concentrations of retinol were lower for the LC diet than for either the MC or the M diet. Both the LC and MC diets produced a similar pattern of decline in plasma retinol over the 4 weeks of ingestion. This may have implications for longer-term WL on such diets. Interestingly, although vitamin E recommended daily

Table 7. Effect of the maintenance (M), low-carbohydrate (LC) and medium-carbohydrate (MC) diets on the homeostasis model assessment of insulin resistance (HOMA-IR) and plasma lipaemia* in sixteen volunteers

	M	LC	MC	SED	P
HOMA-IR	3.94 ^a	2.15 ^b	2.47 ^b	0.430	<0.001
Glucose (mmol/l)	5.82 ^a	5.21 ^b	5.66 ^c	0.190	0.01
Insulin (pm)	14.7 ^a	8.8 ^b	9.7 ^b	1.33	<0.001
3-OHB (mm)	0.05 ^a	1.96 ^b	0.44 ^a	0.236	<0.001
Cholesterol (mm)	4.81 ^a	4.43 ^{a,b}	4.00 ^b	0.273	0.022
LDL (mm)	2.89 ^a	2.53 ^{a,b}	2.45 ^b	0.139	0.009
HDL (mm)	1.04 ^a	1.28 ^c	1.15 ^b	0.055	<0.001
TAG (mm)	1.66 ^b	0.95 ^a	1.01 ^a	0.133	<0.001
Leptin (µg/l)	21.20 ^a	8.21 ^b	10.78 ^b	1.98	<0.001
Urea (mm)	5.3 ^a	7.6 ^c	6.9 ^b	0.33	<0.001

3-OHB, β-hydroxybutyrate.

^{a,b,c} Values within rows with unlike superscript letters were significantly different ($P < 0.05$).

* HOMA-IR (from fasting insulin (pm) × fasting glucose (mm)/22.5).

intake was met on all diets, plasma concentrations of both α-tocopherol and γ-tocopherol declined over time on the LC diet. Deficiencies of α-tocopherol are implicated in several clinical conditions including heart disease, strokes and cancer⁽²⁴⁾. This decrease over a period of WL is in contrast to studies which indicate that obesity is associated with lower plasma vitamin E concentrations^(21,25). Smoking-related effects can be excluded, as all volunteers were non-smokers. Consequently, changes in plasma α-tocopherol concentrations probably reflect the tocopherol homologues present in the plant oil components of the WL diets. Nevertheless, concentrations were not indicative of an overt deficiency as classified by the UK guidelines of below 6.5 µg/ml⁽²⁶⁾. Whether such concentrations are attained with the prolonged use of WL diets is unclear, but supplementation with α-tocopherol may be warranted in the longer term. In contrast, vitamin C values were stable and greater on the LC diet compared with the MC and M diets. This was probably due to the use of unsweetened orange juice as part of the LC menu, but illustrates that dietary manipulation can be used to achieve a good nutritional status, even when food sources containing carbohydrate are severely restricted. Indeed, the LC diet contained twice as much fruit and vegetables (g) as the MC diet (Appendix 2) to match the weight of the meals. The duration of the study is relatively short, when considering the storage and turnover of fat-soluble vitamins *v.* water-soluble vitamins. For example, symptoms of scurvy (vitamin C deficiency) are apparent after a month on a deficient diet, but for vitamin E with tissue-specific stores, it could take up to 2 years for subjects to become deficient on the diets used. However, much of nutrition interest in this area is around defining 'optimal levels for health', and little is known about how diet and phenotype interact to determine the optimal nutrient profile at a population level or individual level during WL or weight maintenance.

Plasma cardiovascular risk markers

Markers of inflammation and endothelial dysfunction are elevated in obesity and type 2 diabetes mellitus and allegedly predict systemic atherogenesis risk⁽²⁷⁾. Interventions that lower the glycaemic response to foods, by lowering either the glycaemic index or the glycaemic load, have demonstrated improvement in risk markers for both type 2 diabetes and CHD⁽²⁸⁾. WL by lifestyle modification also improves the cardiovascular risk profile, indicated by large-scale efficacy trials, as reviewed by Horton⁽²⁹⁾. Much work on these adhesion molecules have focused on surgical patients undergoing gastric procedures for weight control and known to improve metabolic health⁽³⁰⁾. Although the present study reports no statistical improvement in the markers with WL, larger studies have. The numerical values for intercellular adhesion molecules are similar to previous reports before (333 ng/ml) and after (274 ng/ml) WL, in patients undergoing laparoscopic gastric banding⁽²⁷⁾. In terms of the role of diet composition in WL, Keogh *et al.*^(31,32) compared the LC with MC WL diet in two groups of subjects and reported no specific diet effects on short-term (8-week) improvement in CVD risk factors, but

observed a decrease in flow-mediated dilation in the LC group at 1 year⁽³³⁾.

Adiponectin is associated with markers of chronic inflammation, with a rise following WL in morbidly obese patients after gastropasty⁽³⁴⁾ or modest WL⁽³⁵⁾. The lack of change in the present study suggests that this marker to be less responsive to more modest WL. Indeed, Keogh *et al.*⁽³²⁾ reported no change in adiponectin associated with short-term (8-week) WL on a LC or a MC diet, but there was an improvement after 1 year of maintenance following a LC diet⁽³¹⁾. Furthermore, while adiponectin is more responsive to modest WL of between 5 and 10%⁽³⁶⁾, smaller losses (up to 5% body weight) result in no change in obese women⁽³⁷⁾.

Plasma inflammatory markers – cytokines

Plasma adiponectin concentrations are thought to be associated with the inhibition of vascular inflammation, being capable of lowering TNF- α levels in knockout mice⁽³⁸⁾. Obesity is associated with reduced insulin sensitivity linked to inflammatory status⁽³⁹⁾. Many of the cytokines, including TNF- α , IL-1, IL-6 and IL-18, have shown elevated plasma concentration in the insulin-resistant state⁽³⁹⁾ and an increase with obesity⁽⁴⁰⁾, but a decrease with WL⁽⁴¹⁾. As with many other WL studies, the volunteers in the present study showed an improved (lowered) homeostasis model assessment of insulin resistance in response to WL^(42,43). This substantial improvement was not associated, however, with altered cytokine status, except for decreased TNF- α and IL-10. Responses in cytokines with WL have been inconsistent, however. For example, although IL-6 has been shown to decrease⁽⁴⁴⁾, in other studies, no change has been observed in others⁽⁴⁵⁾. Similarly, TNF- α either decreases⁽⁴⁶⁾ or is unaltered⁽⁴⁴⁾ by fat loss. Clearly, links between obesity, inflammatory status and response to WL need to be better understood.

Plasma metabolic markers

Despite the LC WL diet being high in fat (66% energy), there was no detrimental effect on lipaemia. Indeed, the present study indicates a significant improvement in the lipid profile. This is in accordance with several reviews^(47–49), reporting that LC diets, when accompanied with WL, positively improve the lipid profile. Nonetheless, greater improvements in total cholesterol and LDL were observed with the MC diet, which probably reflects the lower fat (35% fat) content of this diet. Furthermore, it is well recognised that LC diets promote a reduction in fasting glucose and insulin⁽⁴⁸⁾ and improve insulin sensitivity⁽⁵⁰⁾, although in the present study, the LC diet did not offer any metabolic advantage over the MC diet. The measurement of metabolic markers was conducted during active WL, and it may be that the data reflect negative energy balance and not the lower body-weight status or diet composition, *per se*. It may be that these markers change accordingly once subjects return to energy balance status. We applied the Mensink & Katan⁽⁵¹⁾ predictive equation for estimating change in plasma cholesterol from diet composition. This regression equation assumes substituting

carbohydrate with fatty acids for an isoenergetic diet. This predicted a decrease of 0.2 mmol/l from the MC to LC diets, reflecting the decrease in saturated fat consumption. There was a decrease of 0.43 mmol/l. There are a number of limitations of the equations, and in this scenario, they are being applied in negative energy balance, which may alter the predictive power. Although leptin concentrations are usually correlated with BMI and body fat, there is usually a rapid decrease associated with negative energy balance and WL⁽⁵²⁾, a finding confirmed in the present study where values halved from baseline, with a fat loss of only 3.7–4.0 kg or 1.5–2.0% initial body fat.

Although the study has small subject numbers, it benefits from a within-subject design and complete control over dietary input, allowing precise metabolic measurements. The assumption that LC diets have a negative impact on vitamin and antioxidant intake is not supported. In conclusion, the present data would suggest that both high-protein LC and high-protein MC WL diets are safe within this relatively short period of time (4 weeks), as assessed by the reported biomarkers, and under medically supervised conditions, they could be used to achieve considerable WL in order to improve mortality and morbidity in obese patients. Antioxidant supplements may be warranted if LC WL diets are consumed for a prolonged period.

Acknowledgements

We are grateful for the assistance from Glenn Harden in processing the plasma samples, and Marion Scott, Jean Bryce, Nina Lamza and Kim Giles for the preparation of the study diets. We also thank Sylvia Hay and Linda Dewar for support in the Human Nutrition Unit. A. M. J., G. E. L. and G. G. D. were responsible for the study concept and design. A. M. J., D. M. B., P. C. M., C. L. F. and G. G. D. were responsible for the sample collection and analysis. A. M. J., G. E. L., G. G. D. and G. W. H. were responsible for the data analysis. A. M. J., G. G. D., G. W. H. and G. E. L. were responsible for the first draft and critical revision of the manuscript for important intellectual content. None of the authors had a personal or financial conflict of interest. The Rowett Institute and Biomathematics and Statistics Scotland are grateful for funding from the Scottish Government who provided funding for the present study.

References

1. Johnstone AM, Horgan GW, Murison SD, *et al.* (2008) Effects of a high-protein ketogenic diet on hunger, appetite, and weight loss in obese men feeding *ad libitum*. *Am J Clin Nutr* **87**, 44–55.
2. Skov AR, Toubro S, Ronn B, *et al.* (1999) Randomized trial on protein *vs.* carbohydrate in *ad libitum* fat reduced diet for the treatment of obesity. *Int J Obes Relat Metab Disord* **23**, 528–536.
3. Mayor S (2003) Researcher criticised for comments on the Atkins diet. *BMJ* **327**, 414.
4. Nöthlings U, Schulze MB, Weikert C, *et al.* (2008) Intake of vegetables, legumes, and fruit, and risk for all-cause,

- cardiovascular, and cancer mortality in a European diabetic population. *J Nutr* **138**, 775–781.
5. Nomura AM, Wilkens LR, Murphy SP, *et al.* (2008) Association of vegetable, fruit, and grain intakes with colorectal cancer: the Multiethnic Cohort Study. *Am J Clin Nutr* **88**, 730–737.
 6. Duncan SH, Belenguer A, Holtrop G, *et al.* (2007) Reduced dietary intake of carbohydrate by obese subjects, undergoing weight loss, results in decreased butyrate and the populations of butyrate-producing bacteria in faeces. *Appl Environ Microbiol* **73**, 1073–1078.
 7. Bravata DM, Sanders L, Huang J, *et al.* (2003) Efficacy and safety of low-carbohydrate diets: a systematic review. *JAMA* **289**, 1837–1850.
 8. Skov AR, Toubro S, Bülow J, *et al.* (1999) Changes in renal function during weight loss induced by high vs. low-protein low-fat diets in overweight subjects. *Int J Obes Relat Metab Disord* **23**, 1170–1177.
 9. Skov AR, Haulrik N, Toubro S, *et al.* (2002) Effect of protein intake on bone mineralization during weight loss: a 6-month trial. *Obes Res* **10**, 432–438.
 10. Holland B, Welch AA, Unwin ID, *et al.* (1991) *McCance and Widdowson's The Composition of Foods*. Cambridge: The Royal Society of Chemistry.
 11. Johnstone AM, Murison SD, Duncan JS, *et al.* (2005) Factors influencing variation in basal metabolic rate include fat-free mass, fat mass, age, and circulating thyroxine but not sex, circulating leptin, or triiodothyronine. *Am J Clin Nutr* **82**, 941–948.
 12. Fuller NJ, Jebb SA, Laskey MA, *et al.* (1992) Four-compartment model for the assessment of body composition in humans: comparison with alternative methods and evaluation of the density and hydration of fat-free mass. *Clin Sci* **82**, 687–693.
 13. Pullicino E, Coward WA, Stubbs RJ, *et al.* (1990) Bedside and field methods for assessing body composition: comparison with the deuterium dilution technique. *Eur J Clin Invest* **40**, 753–762.
 14. Duthie GG (1999) Determination of activity of antioxidants in human subjects. *Proc Nutr Soc* **58**, 1015–1024.
 15. Matthews DR, Hosker JP, Rudenski AS, *et al.* (1985) Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* **28**, 412–419.
 16. Department of Health (1991) *Dietary Reference Values for Food Energy and Nutrients for the United Kingdom. Report of the Panel on Dietary Reference Values of the Committee on Medical Aspects of Food Policy*. London: HMSO.
 17. Food and Nutrition Board (2000) *A Report of the Panel on Dietary Antioxidants and Related Compounds Subcommittee on Upper Reference Levels of Nutrients and Interpretation and Uses of Dietary Reference Intakes, and the Standing Committee on the Scientific Evaluation of Dietary Reference Intakes*. New York, NY: Academic Press.
 18. Astrup A, Meinert Larsen T, Harper A, *et al.* (2004) Atkins and other low-carbohydrate diets: hoax or an effective tool for weight loss? *Lancet* **364**, 897–899.
 19. Volek J, Sharman M, Gómez A, *et al.* (2004) Comparison of energy-restricted very low-carbohydrate and low-fat diets on weight loss and body composition in overweight men and women. *Nutr Metab (Lond)* **8**, 13.
 20. Melia AT, Koss-Twardy SG & Zhi J (1996) The effect of orlistat, an inhibitor of dietary fat absorption, on the absorption of vitamins A and E in healthy volunteers. *J Clin Pharmacol* **36**, 647–653.
 21. Gasteyer C, Suter M, Gaillard RC, *et al.* (2008) Nutritional deficiencies after Roux-en-Y gastric bypass for morbid obesity often cannot be prevented by standard multivitamin supplementation. *Am J Clin Nutr* **87**, 1128–1133.
 22. Ross M, Crosley KM, Brown KM, *et al.* (1995) Plasma concentrations of carotenoids and antioxidant vitamins in Scottish males: influences of smoking. *Eur J Clin Nutr* **49**, 861–865.
 23. Moats C & Rimm EB (2007) Vitamin intake and risk of coronary disease: observation versus intervention. *Curr Atheroscler Rep* **9**, 508–514.
 24. Dietrich M, Block G, Norkus EP, *et al.* (2003) Smoking and exposure to environmental tobacco smoke decrease some plasma antioxidants and increase gamma-tocopherol *in vivo* after adjustment for dietary antioxidant intakes. *Am J Clin Nutr* **77**, 160–166.
 25. Aasheim ET, Hofso D, Hjelmestaeth J, *et al.* (2008) Vitamin status in morbidly obese patients: a cross-sectional study. *Am J Clin Nutr* **87**, 362–369.
 26. COMA (1991) *Report 41: Dietary Reference Values for Food Energy and Nutrients for the United Kingdom*. London: HMSO.
 27. Pontiroli AE, Pizzocri P, Koprivec D, *et al.* (2004) Body weight and glucose metabolism have a different effect on circulating levels of ICAM-1, E-selectin, and endothelin-1 in humans. *Eur J Endocrinol* **150**, 195–200.
 28. Kelly S, Frost G, Whittaker V, *et al.* (2004) Low glycaemic index diets for coronary heart disease. *The Cochrane Database of Systematic Reviews* 2004, issue 4, CD004467. Chichester: John Wiley & Sons Ltd.
 29. Horton ES (2009) Effects of lifestyle changes to reduce risks of diabetes and associated cardiovascular risks: results from large scale efficacy trials. *Obesity* **17**, 43–48.
 30. Pontiroli AE, Frigè F, Paganelli M, *et al.* (2009) In morbid obesity, metabolic abnormalities and adhesion molecules correlate with visceral fat, not with subcutaneous fat: effect of weight loss through surgery. *Obes Surg* **19**, 745–750.
 31. Keogh JB, Brinkworth GD & Clifton PM (2007) Effects of weight loss on a low-carbohydrate diet on flow-mediated dilatation, adhesion molecules and adiponectin. *Br J Nutr* **98**, 852–859.
 32. Keogh JB, Brinkworth GD, Noakes M, *et al.* (2008) Effects of weight loss from a very-low-carbohydrate diet on endothelial function and markers of cardiovascular disease risk in subjects with abdominal obesity. *Am J Clin Nutr* **87**, 567–576.
 33. Wycherley TP, Brinkworth GD, Keogh JB, *et al.* (2010) Long-term effects of weight loss with a very low carbohydrate and low fat diet on vascular function in overweight and obese patients. *J Intern Med* **267**, 452–461.
 34. Kopp HP, Krzyzanowska K, Möhlig M, *et al.* (2005) Effects of marked weight loss on plasma levels of adiponectin, markers of chronic subclinical inflammation and insulin resistance in morbidly obese women. *Int J Obes (Lond)* **29**, 766–771.
 35. Forsythe LK, Wallace JM & Livingstone MB (2008) Obesity and inflammation: the effects of weight loss. *Nutr Res Rev* **21**, 117–1133.
 36. Madsen EL, Rissanen A, Bruun JM, *et al.* (2008) Weight loss larger than 10% is needed for general improvement of levels of circulating adiponectin and markers of inflammation in obese subjects: a 3-year weight loss study. *Eur J Endocrinol* **158**, 179–187.
 37. Varady KA, Tussing L, Bhutani S, *et al.* (2009) Degree of weight loss required to improve adipokine concentrations and decrease fat cell size in severely obese women. *Metabolism* **58**, 1096–1101.

38. Manigrasso MR, Ferroni P, Santilli F, *et al.* (2005) Association between circulating adiponectin and interleukin-10 levels in android obesity: effects of weight loss. *J Clin Endocrinol Metab* **90**, 5876–5879.
39. Eckel RH (2007) Mechanisms of the components of the metabolic syndrome that predispose to diabetes and atherosclerotic CVD. *Proc Nutr Soc* **66**, 82–95.
40. Ruan H & Lodish HF (2003) Insulin resistance in adipose tissue: direct and indirect effects of tumor necrosis factor- α . *Cytokine Growth Factor Rev* **14**, 447–455.
41. Barinas-Mitchell E, Kuller LH, Sutton-Tyrrell K, *et al.* (2006) Effect of weight loss and nutritional intervention on arterial stiffness in type 2 diabetes. *Diabetes Care* **29**, 2218–2222.
42. Belobrajdic DP, Frystyk J, Jeyaratnaganthan N, *et al.* (2010) Moderate energy restriction-induced weight loss affects circulating IGF levels independent of dietary composition. *Eur J Endocrinol* **162**, 1075–1082.
43. Sheu WH, Chang TM, Lee WJ, *et al.* (2008) Effect of weight loss on proinflammatory state of mononuclear cells in obese women. *Obesity (Silver Spring)* **16**, 1033–1038.
44. Bastard JP, Jardel C, Bruckert E, *et al.* (2000) Elevated levels of interleukin 6 are reduced in serum and subcutaneous adipose tissue of obese women after weight loss. *J Clin Endocrinol Metab* **85**, 3338–3342.
45. Shadid S, Stehouwer CD & Jensen MD (2006) Diet/exercise versus pioglitazone: effects of insulin sensitization with decreasing or increasing fat mass on adipokines and inflammatory markers. *J Clin Endocrinol Metab* **91**, 3418–3425.
46. Bruun JM, Verdich C, Toubro S, *et al.* (2003) Association between measures of insulin sensitivity and circulating levels of interleukin-8, interleukin-6 and tumor necrosis factor- α . Effect of weight loss in obese men. *Eur J Endocrinol* **148**, 535–542.
47. Westman EC, Mavropoulos J, Yancy WS, *et al.* (2003) A review of low-carbohydrate ketogenic diets. *Curr Atheroscler Rep* **5**, 476–483.
48. Volek JS & Sharman MJ (2004) Cardiovascular and hormonal aspects of very-low-carbohydrate ketogenic diets. *Obes Res* **12**, 115–1123.
49. Noble CA & Kushner RF (2006) An update on low-carbohydrate, high-protein diets. *Curr Opin Gastroenterol* **22**, 153–159.
50. Bisschop PH, de Metz J, Ackermans MT, *et al.* (2001) Dietary fat content alters insulin-mediated glucose metabolism in healthy men. *Am J Clin Nutr* **73**, 554–559.
51. Mensink RP & Katan MB (1992) Effect of dietary fatty acids on serum lipids and lipoproteins. A meta-analysis of 27 trials. *Arterioscler Thromb* **12**, 911–919.
52. Kerkisick C, Thomas A, Campbell B, *et al.* (2009) Effects of a popular exercise and weight loss program on weight loss, body composition, energy expenditure and health in obese women. *Nutr Metab (Lond)* **6**, 23.

Appendix 1. Menu of the maintenance meals provided

Meals (3 d rotation)	
Breakfast	
1	Cornflake cereal, croissant, scrambled eggs and orange juice
2	Alpen cereal, toast and yogurt
3	Sultana bran cereal, cheesy scrambled eggs and toast
Lunch	
1	Cheese and pickle roll with salad
2	Ham and coleslaw roll with salad
3	Tuna mayonnaise roll with salad
Dinner	
1	Soya mince stir fry and apple crumble
2	Tuna and tomato pasta and apple pie
3	Chicken and herb pasta and apricots in custard
Average fruit and vegetable intake	
Days 1–3	479 g

Appendix 2. Menu of high-protein weight-loss meals provided

Meals	LC	MC
Breakfast (<i>ad libitum</i> choice)		
1	Scrambled eggs and turkey rashers	Mixed grill (1)
2	Mixed grill and grapefruit	Crumpet, ham and yogurt
3	Bacon with cheesy scrambled eggs	Mixed grill (2) with toast
4	Mixed grill	Mixed grill (3) with toast
5	Spanish omelette and yogurt	Red fruit dairy smoothies
6	Turkey rashers, poached eggs and salad	Continental
7	Bacon and poached eggs with raspberry yogurt	Bacon, beans, potato waffle and milk
8	Omelette, grilled tomatoes and mushrooms	Scrambled eggs, turkey rashers and croissant
Lunch (7 d rotation)		
1	Chicken breast salad	Chicken and macaroni salad
2	Cheese and ham salad	Cheese and ham salad
3	Cheese and chicken Caesar salad	Cheese and chicken Caesar salad
4	Tuna mayonnaise salad	Tuna mayonnaise salad
5	Ham, eggs and cheese salad	Chicken and sweetcorn salad
6	Pork salad	Pork salad
7	Tuna and egg salad	Tuna and bacon salad
Dinner (7 d rotation)		
1	Chilli beef	Chilli beef risotto
2	Chicken creole	Chicken creole
3	Chicken curry	Chicken curry
4	Pork loin and ratatouille	Pork grill
5	Gammon and cauliflower bake	Spaghetti carbonara
6	Chicken and bacon stir fry	Chicken stir fry
7	Steak and mushrooms	Meat stew
Average fruit and vegetable intake		
Days 1–7	627 g	363 g

LC, low carbohydrate; MC, moderate carbohydrate.