
INSTRUCTION TO CONTRIBUTORS

The *Journal of Helminthology* publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance, but only in exceptional circumstances will systematic or taxonomic studies be acceptable.

Manuscripts, which must be in English or French (with an English summary), should be addressed to:

The Editor, Journal of Helminthology,
London School of Hygiene and Tropical Medicine,
Winches Farm Field Station,
395 Hatfield Road,
St Albans, Herts AL4 0XQ,
England.

Two copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief Introduction; Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Research Notes should also be preceded by a brief abstract. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration will be printed is 12.0 × 20.0 cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space, should be indicated. Some indication of scale (preferably a scale bar) should normally be given on the figure. Photocopies of illustrations should be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints of the same size as they are to appear in the Journal (maximum size 12.0 × 20.0 cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

DUKE, B. O. L. (1971) The ecology of onchocerciasis in man and animals. In: *Ecology and Physiology of parasites* (editor, A. M. Fallis) pp. 213–222. Adam Hilger Ltd.: London.

JAMES, C. & WEBBE, G. (1973) A comparison of Egyptian and East African strains of *Schistosoma haematobium*. *Journal of Helminthology*, **47**, 49–59.

25 offprints are provided free of charge; additional copies may be ordered at the proof stage.

Contents

	Pages
A comparison of a modified quick-Kato technique and the Stoll dilution method for field examination for <i>Opisthorchis viverrini</i> eggs. V. VIYANANT, W. Y. BROCKELMAN, P. LEE, S. ARDSUNGNOEN and E. S. UPATHAM	191-195
Factors influencing the establishment of <i>Mesocestoides corti</i> in mice following oral inoculation of tetrathyridia. T. R. WHITE, R. C. A. THOMPSON and W. J. PENHALE	197-203
<i>Anisakis simplex</i> (Rudolphi, 1809, det. Krabbe, 1878) (Nematoda: Ascaridoidea): morphology and morphometry of larvae from euphausiids and fish, and a review of the life-history and ecology. JOHN W. SMITH	205-224
Estimation of the numbers and infectivity of <i>Nippostrongylus</i> larvae. ANNE KEYMER, JEAN MARTIN and S. M. WAINWRIGHT	225-230
<i>In vivo</i> and <i>in vitro</i> development of <i>Schistosoma margrebowiei</i> . MARTIN G. OGBE	231-235
Pulmonary superoxide dismutase activity in the mouse infected with <i>Ancylostoma caninum</i> larvae. (Research Note). A. B. KULKARNI, M. K. BHOPALE and K. D. SHARMA	237-239
Demonstration of collagenase activity in adult <i>Strongyloides ratti</i> (Nematoda: Strongyloididae) and its absence in the infective larvae. GUTA WERTHEIM, SARAH LUSTIGMAN, HERMINA SILBERMAN and SHMUEL SHOSHAN	241-246
Schistosomiasis in Swaziland—a comparative study of three irrigated estates. JAMES W. M. LOGAN	247-253
Cement glands in <i>Acanthocephalus serendibensis</i> Cruz & Mills, 1970 (Acanthocephala: Palaeacanthocephala: Echinorhynchidae). W. R. BRECKENRIDGE and SHIRANI NATHANAEL	255-258
Biology of <i>Monanema globulosa</i> , a rodent filaria with skin-dwelling microfilariae. A. E. BIANCO, R. MULLER and G. S. NELSON	259-278
Absence of shock-like reactions to ivermectin in dogs infected with <i>Dirofilaria immitis</i> . P. F. L. BOREHAM and R. B. ATWELL	279-281
SUPPLEMENTARY REVIEW ARTICLE	
Ascaridoid nematodes of amphibians and reptiles: <i>Freitasascaris</i> n.g. J. F. A. SPRENT	283-284
Corrigendum	196