- 10 Saunders JB, Aasland OG, Babor TF, de la Fuente JR, Grant M. Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO collaborative project on early detection of persons with harmful alcohol consumption. Addiction 1993; 88: 791–804.
- 11 Zigmund AS, Snaith RP. The hospital anxiety and depression scale. Acta Psychiatr Scand 1983; 67: 361–70.
- 12 Kapur N, Cooper J, King-Hele S, Webb R, Lawlor M, Rodway C, et al. The repetition of suicidal behavior: a multicenter cohort study. J Clin Psychiatry 2006; 67: 1599–609.
- 13 Colman I, Newman SC, Schopflocher D, Bland RC, Dyck RJ. A multivariate study of predictors of repeat parasuicide. Acta Psychiatr Scand 2004; 109: 306–12.
- 14 Hawton K, Rodham K, Evans E, Weatherall R. Deliberate self harm in adolescents: self report survey in schools in England. BMJ 2002; 325: 1207–11
- 15 Horvath AO, Greenberg LS. Development and validation of the Working Alliance Inventory. *J Couns Psychol* 1989; 36: 223–33.
- 16 Larsen DL, Attkisson CC, Hargreaves WA, Nguyen TD. Assessment of client/patient satisfaction: development of a general scale. Eval Program Plan 979; 2: 197–207
- 17 Tait L, Birchwood M, Trower P. A new scale (SES) to measure engagement with community mental health services. *J Ment Health* 2002; 11: 191–8.
- 18 Tennant R, Hiller L, Fishwick R, Platt S, Joseph S, Weich S, et al. The Warwick-Edinburgh Mental Well-being Scale (WEMWBS): development and UK validation. *Health Qual Life Outcomes* 2007; 5: 63–75.
- 19 Mundt JC, Marks IM, Shear MK, Greist JM. The Work and Social Adjustment Scale: a simple measure of impairment in functioning. Br J Psychiatry 2002; 180: 461–4.
- 20 Gardner W, Hoge SK, Bennett N, Roth LH, Lidz C, Monahan J, et al. Two scales for measuring patients' perceptions for coercion during mental hospital admission. Behav Sci Law 1993; 11: 307–21.
- 21 Kind P. The EuroQol instrument: an index of health related quality of life. In Quality of Life and Pharmacoeconomics in Clinical Trials (2nd edn) (ed. B Spiker). Lippincott-Raven, 1996.
- 22 Byford S, Knapp M, Greenshields J, Ukoumunne OC, Jones V, Thompson SG, et al. Cost-effectiveness of brief cognitive behaviour therapy versus treatment as usual in recurrent deliberate self-harm: a decision-making approach. *Psychol Med* 2003; 33: 977–86.

- 23 National Institute for Health and Clinical Excellence. Guide to the Methods of Technology Appraisal. NICE, 2008.
- 24 Tyrer P, Thompson G, Schmidt U, Jones V, Knapp M, Davidson K, et al. Randomized controlled trial of a brief cognitive behaviour therapy versus treatment as usual in recurrent deliberate self-harm: the POPMACT study. Psychol Med 2003; 33: 969–76.
- 25 Barber JA, Thompson SG. Analysis of cost data in randomised trials: an application of the non-parametric bootstrap. Stat Med 2000; 19: 3219–36.
- 26 Efron B, Tibshirani RJ. An Introduction to the Bootstrap. Chapman Hall, 1993.
- 27 Bateman A, Fonagy P. Randomized controlled trial of outpatient mentalization-based treatment versus structured clinical management for borderline personality disorder. Am J Psychiatry 2009; 166: 1355–64.
- 28 Linehan MM, Armstrong HE, Suarez A, Allmon D, Heard HL. Cognitive-behavioral treatment of chronically parasuicidal borderline patients. Arch Gen Psychiatry 1991; 48: 1060–4.
- 29 Bateman AW, Fonagy P. Effectiveness of psychotherapeutic treatment of personality disorder. *Br J Psychiatry* 2000; 177: 138–43.
- 30 Campbell MK, Snowdon C, Francis D, Elbourne D, McDonald AM, Knight R, et al. Recruitment to randomised trials: strategies for trial enrolment and participation study. The STEPS study. *Health Technol Assess* 2007; 11: iii, ix–105.
- 31 Bateman A, Fonagy P. Effectiveness of partial hospitalization in the treatment of borderline personality disorder: a randomized controlled trial. Am J Psychiatry 1999; 156: 1563–9.
- 32 Borschmann R, Hogg J, Phillips R, Moran P. Measuring self-harm in adults: a systematic review. *Eur Psychiatry* 2011; 27: 176–80.
- 33 Evans C, Crawford B. Patient self-reports in pharmacoeconomic studies: their use and impact on study validity. Pharmacoecon 1999; 15: 241–56.
- 34 Giesen-Bloo J, Van Dyck R, Spinhoven P, Van Tilburg W, Dirkson C, Van Asselt T, et al. Outpatient psychotherapy for borderline personality disorder: randomized trial of schema-focused therapy vs transferencefocused psychotherapy. Arch Gen Psychiatry 2006; 63: 649–58.
- 35 Henderson C, Flood C, Leese M, Thornicroft G, Sutherby K. Views of service users and providers on joint crisis plans. Soc Psychiatry Psychiatr Epidemiol 2008; 44: 369–76.
- **36** Kim MM, Van Dorn RA, Scheyett AM, Elbogen EE, Swanson JW, Swartz MS, et al. Understanding the personal and clinical utility of psychiatric advance directives: a qualitative perspective. *Psychiatry* 2007; **70**:





## A Message

Olive M. Ritch

One of the best of minds destroyed by dementia

does not howl on her knees in the street, does not masturbate

in the magnolia living-room, is not dragged off the roof-top,

naked; no, she leaves a message on her daughter's answer-phone

saying: there's an echo, an echo in my head.

Olive M. Ritch is currently completing a PhD in creative writing (on the theme of memory) at the University of Aberdeen. This poem is from *The Hippocrates Prize 2011*, published by The Hippocrates Prize in association with Top Edge Press. © Olive M. Ritch.

Chosen by Femi Oyebode.

The British Journal of Psychiatry (2013) 202, 364. doi: 10.1192/bjp.bp.111.103150