



Review Article

Cite this article: Licá ICL *et al.* (2023). Immunological mechanisms involved in macrophage activation and polarization in schistosomiasis. *Parasitology* **150**, 401–415. <https://doi.org/10.1017/S0031182023000021>

Received: 18 May 2022

Revised: 21 December 2022

Accepted: 24 December 2022

First published online: 5 January 2023

Keywords:

Granuloma; M1; M2; phagocytes; schistosomiasis

Author for correspondence:

Flávia Raquel Fernandes Nascimento,
E-mail: flavia.nascimento@ufma.br

Immunological mechanisms involved in macrophage activation and polarization in schistosomiasis

Irla Correia Lima Licá^{1,2} , Gleycka Cristine Carvalho Gomes Frazão^{1,2} ,
Ranielly Araujo Nogueira^{1,2} , Maria Gabriela Sampaio Lira^{1,2} ,
Vitor Augusto Ferreira dos Santos^{1,2}, João Gustavo Mendes Rodrigues³ ,
Guilherme Silva Miranda⁴ , Rafael Cardoso Carvalho¹ , Lucilene Amorim Silva^{1,2,5},
Rosane Nassar Meireles Guerra^{1,2,5} and Flávia Raquel
Fernandes Nascimento^{1,2,5}

¹Graduate Program in Health Sciences, Center for Biological and Health Sciences, Federal University of Maranhão, São Luís, MA, Brazil; ²Laboratory of Immunophysiology, Center for Biological and Health Sciences, Federal University of Maranhão, São Luís, MA, Brazil; ³Department of Parasitology, Institute of Biological Sciences, Federal University of Minas Gerais, Belo Horizonte, Minas Gerais, Brazil; ⁴Department of Biology, Federal Institute of Education, Science and Technology of Maranhão, São Raimundo das Mangabeiras, Brazil and ⁵Department of Pathology, Center for Biological and Health Sciences, Federal University of Maranhão, São Luís, MA, Brazil

Abstract

Human schistosomiasis is caused by helminths of the genus *Schistosoma*. Macrophages play a crucial role in the immune regulation of this disease. These cells acquire different phenotypes depending on the type of stimulus they receive. M1 macrophages can be ‘classically activated’ and can display a proinflammatory phenotype. M2 or ‘alternatively activated’ macrophages are considered anti-inflammatory cells. Despite the relevance of macrophages in controlling infections, the role of the functional types of these cells in schistosomiasis is unclear. This review highlights different molecules and/or macrophage activation and polarization pathways during *Schistosoma mansoni* and *Schistosoma japonicum* infection. This review is based on original and review articles obtained through searches in major databases, including Scopus, Google Scholar, ACS, PubMed, Wiley, Scielo, Web of Science, LILACS and ScienceDirect. Our findings emphasize the importance of *S. mansoni* and *S. japonicum* antigens in macrophage polarization, as they exert immunomodulatory effects in different stages of the disease and are therefore important as therapeutic targets for schistosomiasis and in vaccine development. A combination of different antigens can provide greater protection, as it possibly stimulates an adequate immune response for an M1 or M2 profile and leads to host resistance; however, this warrants *in vitro* and *in vivo* studies.

Introduction

Human schistosomiasis is a neglected parasitic disease with great relevance to public health. Worldwide, it is estimated that approximately 230–250 people are infected and 700–800 million live in areas that are at risk of infection, mainly in countries located in South America, Asia and Africa (Steinmann *et al.*, 2006; Colley *et al.*, 2014; McManus *et al.*, 2018; Wei *et al.*, 2018; WHO, 2020). In addition, approximately 200 000–280 000 deaths occur each year due to schistosomiasis and its complications (LoVerde, 2019). The high prevalence of schistosomiasis is mainly related to people living in extreme poverty and poor sanitation, which represent a serious risk to human health (Ismail *et al.*, 2014; Bajiro *et al.*, 2017; Verjee, 2019).

The infection is caused by helminths of the genus *Schistosoma* (Colley *et al.*, 2014; Stingl and Stingl, 2017; WHO, 2020), belonging to the class Trematoda and phylum Platyhelminthes. The main aetiologic agents of this disease, in terms of clinical relevance, are *Schistosoma japonicum*, *Schistosoma mansoni* and *Schistosoma haematobium* (WHO, 2020). In this review, we focus only on *S. mansoni* and *S. japonicum*, as they are the main species associated with hepatic and intestinal schistosomiasis (Wilson *et al.*, 2007; Chen *et al.*, 2013; McManus *et al.*, 2018).

There are 2 distinct phases of clinical progression of intestinal schistosomiasis: the acute and the chronic phases (Gobbi *et al.*, 2020). During the early stages of acute phase of schistosomiasis (before parasite oviposition), there is a predominance of the T helper type 1 (Th1) immune response (Pearce *et al.*, 1991; Hesse *et al.*, 2001; Pearce and MacDonald, 2002; Colley and Secor, 2014). After schistosome oviposition, the immune response becomes strongly polarized to the Th2 profile, which is related to increasing production of interleukin-4 (IL-4), IL-5, IL-9 and IL-13 (Pearce and MacDonald, 2002; Burke *et al.*, 2009). This immune environment is responsible for the formation of granulomas in tissues (Grzych *et al.*, 1991; Brunet *et al.*, 1997; Hoffmann *et al.*, 2000). The granuloma has an important role for the host, because it contains the tissue damage caused by antigens secreted by the schistosome

© The Author(s), 2023. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

eggs (Hams *et al.*, 2013; Schwartz and Fallon, 2018). In the chronic phase of schistosomiasis, there is an increase in the production of regulatory cells in the granuloma, which can modulate granulomatous inflammation, promoting a minimization of the disease severity (Hesse *et al.*, 2004; Lundy and Lukacs, 2013). However, if this inflammatory reaction does not have an adequate modulation, the granulomas progressively may evolve into large areas of fibrosis, responsible for the main pathology of schistosomiasis (Hams *et al.*, 2013; Schwartz and Fallon, 2018), including hepatosplenomegaly (Masi *et al.*, 2020), portal hypertension (Grieco *et al.*, 2016) and ascites (Fei-Yue *et al.*, 2017).

Macrophages are cells of the innate immune system that play important roles in controlling infections (Shapouri-Moghaddam *et al.*, 2018), as well as in tissue remodelling processes, both in ontogenesis and wound healing (Kloc *et al.*, 2019). In the course of *S. mansoni* and *S. japonicum* infection, either at its initial stage or during the evolution to the chronic phase, macrophages participate in the immune regulation of the disease (Cortes-Selva *et al.*, 2018; Ho *et al.*, 2022).

Macrophages can acquire different phenotypes depending on the stimuli to which they are subjected to (Atri *et al.*, 2018). These cells can be classified into M1 or 'classically activated' cells, with pro-inflammatory action, and M2 or 'alternatively activated' macrophages, which are considered anti-inflammatory cells (Mills, 2015; Ley, 2017; Locati *et al.*, 2020). However, despite the relevance of macrophages in controlling infections, the participation of the functional types of these cells in acute and chronic schistosomiasis is still not well defined. Thus, this review discusses the different molecules and/or pathways of activation and polarization of macrophages during infection by *S. mansoni* and *S. japonicum*, leading to a better understanding of the role of these cells in the immunopathology of schistosomiasis. Based on this knowledge, we may help identify potential targets for the development of better treatment strategies to reduce the morbidity of this disease.

Methods and criteria for literature selection

This literature review was performed using recognized databases including Scopus, Google Scholar, ACS, PubMed, Wiley, Scielo, Web of Science, LILACS and ScienceDirect and covered original and review articles published in English from 1966 to 2022. Articles involving *in vitro* and/or *in vivo* experiments were included and addressed the main immunological aspects of *S. mansoni* and *S. japonicum* infection related to macrophage polarization, activation and effector functions. To search for these articles, combinations of keywords were used, such as 'Macrophage', 'Schistosoma', 'macrophage polarization and Schistosoma', 'Macrophage and Schistosoma', 'Macrophage and Schistosoma mansoni', 'Macrophage and Schistosoma japonicum'. Research involving coinfections was not included in this study.

Immunopathology of the definitive host against infection by *S. mansoni* and *S. japonicum*

Parasites of the genus *Schistosoma* have complex life cycles (Fig. 1), with generations of asexual reproducing larvae living in freshwater snails, the intermediate hosts (some species of the genus *Biomphalaria* for *S. mansoni* and the genus *Oncomelania* for *S. japonicum*) and another stage of sexual reproduction of adult worms in vertebrate hosts (definitive), including humans (McManus *et al.*, 2018; Nelwan, 2019). Each stage of the parasite's life cycle (cercariae, schistosomulae, adult worms and eggs) within the definitive host triggers a series of immune responses, and consequently, clinical signs that can be harmful to humans (Molehin, 2020; Hambrook and Hanington, 2021; Masamba and Kappo,

2021). The interactions between the host immune system and the parasite can be divided into 2 phases (Fig. 2): acute phase (after and before parasite oviposition) and chronic phase (Gobbi *et al.*, 2020).

The first clinical manifestations of the acute phase (cercarial dermatitis, oedema and pruritus) begin 48–72 h after cercariae penetrate the host's skin, and occurs mainly in individuals from endemic areas (frequently exposed to infection) (He *et al.*, 1990, 2005; Khammo *et al.*, 2002; Ingram *et al.*, 2003; Lambertucci, 2010). The first innate immune barrier encountered by cercariae is the skin (Bartlett *et al.*, 2000; Whitfield *et al.*, 2003; He *et al.*, 2005). This tissue is composed of keratinocytes, whose function is to secrete cytokines with antimicrobial functions (Roupé *et al.*, 2010; Piipponen *et al.*, 2020). Indeed, the keratinocytes are considered the first active cells in response to cercariae infection (Bourke *et al.*, 2015). These cells rapidly respond to infections by secreting inflammatory cytokines [IL-6, IL-12, tumour necrosis factor-alpha (TNF- α) and IL-1 β] to repair damaged tissue (Hogg *et al.*, 2003a, 2003b). When penetrating the host's skin, cercariae also cause an increase in antigen-presenting cells in the innate immune system, such as Langerhans cells and dendritic cells (DCs), as shown in Fig. 2 (Angeli *et al.*, 2001; Kumkate *et al.*, 2007; Hambrook and Hanington, 2021), which contribute to a type 1 cellular immune response (He *et al.*, 2005; Perona-Wright *et al.*, 2006).

Initial immune responses are activated as a result of excretory/secretory (E/S) products released by the cercariae penetrating glands at the time of penetration into the host's skin (Salter *et al.*, 2000; Jenkins *et al.*, 2005a, 2005b; Curwen *et al.*, 2006; Paveley *et al.*, 2009). E/S products assist in the immunomodulatory function exerted by cercariae, as well as condition the remodelling of the extracellular matrix, facilitating its penetration into the skin (Janssen *et al.*, 2016; Leontovych *et al.*, 2020). Liu *et al.* (2015) performed a proteomic analysis of products excreted by *S. japonicum* cercariae at the time of skin entry and identified a variety of E/S proteins, mainly proteases. Among the enzymes that allow this remodelling, the cercarial elastase of *S. mansoni* stands out, which is of great importance in the penetration of cercariae into the skin and can degrade a wide variety of macromolecules present in the human integument (Ingram *et al.*, 2012; El-Faham *et al.*, 2017).

Parasitic E/S products also promote the activation of prostaglandin E2 (PGE2) and prostaglandin D2-producing keratinocytes (Kaisar *et al.*, 2018; Oyesola *et al.*, 2021), which are molecules that induce the production of IL-10 via a cyclooxygenase 2-dependent pathway (Ramaswamy *et al.*, 2000; Harizi *et al.*, 2002; Xue *et al.*, 2005). This type of response is responsible for modulating the immune response that favours parasite survival (Angeli *et al.*, 2001; Hervé *et al.*, 2003; De Oliveira Fraga *et al.*, 2010). Abdel-Ghany *et al.* (2015) suggested that blocking PGE2 might provide partial protection in *S. mansoni*-infected mice. In addition, during the period when cercariae transform into schistosomules and migrate through the skin, PGE2 acts as a potent vasodilator, helping the passage of these larval forms into circulation (Ruzicka and Printz, 1984).

After penetrating the host's skin, cercariae undergo morphological and biochemical changes, transforming into juvenile forms, known as schistosomula, that reach blood vessels (Brink *et al.*, 1977; Wilson, 1987; Curwen and Wilson, 2003). In the bloodstream, the schistosomula is passively transported to the lungs and heart until they finally reach the hepatic portal system, where they develop into adult male or female worms (Miller and Wilson, 1978; Wheater and Wilson, 1979; Nation *et al.*, 2020) (Fig. 1). In this phase before the parasite's oviposition (early stages of acute phase), the host produces a predominantly type 1 immune response, which reaches greater activation between the




Fig. 1. Life cycle of *Schistosoma mansoni* and *Schistosoma japonicum*. (1) The eggs shed in the feces of the definitive host release the miracidia when they come in contact with water (2), which penetrate in soft tissue the intermediate host snail (*Biomphalaria* spp./*Oncomelania* spp.). Inside the snail, the miracidia transform into mother sporocysts, which in turn produce daughter sporocysts by asexual reproduction. After around 30 days post-infection, cercariae emerge from the daughter sporocysts and are shedding by the snails in response to the light and heat (4). The cercariae penetrate the skin of the definitive host (5) and later transform into schistosomula. These larvae enter venous blood vessels and are passively carried to the lungs and heart (6). Upon reaching the hepatic portal system, schistosomula mature, become adult worms (male or female) and mate (7). The mated worms migrate to the lower mesenteric veins of the intestine, where the female sheds the eggs. Part of these eggs pass through the intestinal wall and are eliminated in the feces, starting the cycle again. However, some eggs are not eliminated and get trapped in several organs (mainly the liver and intestines), inducing a potent granulomatous inflammatory response, responsible for schistosomiasis pathology. Source: Created with [BioRender.com](https://www.biorender.com).




Fig. 2. Different immune response profiles during *S. mansoni* and *S. japonicum* infection. Source: Created with [BioRender.com](https://www.biorender.com).

3rd and 5th weeks after exposure to cercariae (Dunne and Cooke, 2005; Gryseels *et al.*, 2006). This response is characterized by high production of pro-inflammatory cytokines, such as IL-1, IL-2,

IL-6, IL-12, interferon-gamma (IFN- γ) and TNF- α (Fig. 2) (Grzych *et al.*, 1991; Pearce *et al.*, 1991; Egesa *et al.*, 2018; Zheng *et al.*, 2020). Coinciding with the migration and sexual

maturation of adult worms, a systemic hypersensitivity reaction occurs in the host, called Katayama syndrome (for *S. japonicum*) or the toxæmic form (for *S. mansoni*), which is associated with an intense Th1 response (Neves, 1992; Ross *et al.*, 2007; Caldas *et al.*, 2008; Langenberg *et al.*, 2019). During primary infections in non-immune individuals, the main symptoms related to this systemic inflammation include a high fever accompanied by chills, profuse sweating, asthenia, myalgia, headache and a non-productive cough (Schwartz *et al.*, 2000; Bottieau *et al.*, 2006).

After parasite oviposition (between 5th and 6th weeks post-infection), there is a change in the profile of immune mediators produced by the host, and the immune response becomes predominantly Th2, which is associated with increasing production of IL-4, IL-5, IL-9 and IL-13 (MacDonald *et al.*, 2002; Pearce *et al.*, 2004; Bartley *et al.*, 2006; Burke *et al.*, 2009). Such changes are responses to soluble egg antigens (SEAs) (Hams *et al.*, 2013), that is composed of a complex mixture of immunostimulatory antigens that are known for their ability to condition DCs to initiate the induction of a Th2 profile (Mouser *et al.*, 2019).

DCs detect, capture and process antigens derived from eggs of *S. mansoni* (Cervi *et al.*, 2004; van Liempt *et al.*, 2007), resulting in their ability to lead to Th2 polarization both *in vitro* and *in vivo* (de Jong *et al.*, 2002; MacDonald *et al.*, 2002; Perona-Wright *et al.*, 2006). The main antigens responsible for this potent induction of a Th2 response are glycoproteins omega 1 (ω -1) and IPSE (IL-4-inducing principle of *S. mansoni* eggs)/alpha 1 (α -1) (Schramm *et al.*, 2006; Meevissen *et al.*, 2010). Glycoprotein ω -1 is present in both SEAs (Dunne *et al.*, 1991) and E/S products from live eggs (Cass *et al.*, 2007), and activates DCs (via C-type and Toll-type lectin receptors), which in turn promotes Th2 differentiation, the main source of type 2 cytokines such as IL-4, IL-5 and IL-13 (Everts *et al.*, 2009). On the other hand, a previous study (Schramm *et al.*, 2006) showed that the glycoprotein IPSE/ α -1 is exclusively released from mature eggs, but likely possesses the same potential to initiate a Th2 response during *S. mansoni* infection. IPSE/ α -1 binds to immunoglobulin and activates basophils, leading to the release of histamine and facilitating

the production of Th2-type cytokines, mainly IL-4 and IL-13 (Schramm *et al.*, 2007; Meyer *et al.*, 2015; Knuhr *et al.*, 2018). Thus, the Th2 response (Fig. 2) is related to low production of IFN- γ and high concentrations of anti-inflammatory cytokines (IL-4, IL-5, IL-10 and IL-13) (Grzych *et al.*, 1991; Pearce *et al.*, 2004; Zheng *et al.*, 2020).

Mechanisms associated with macrophage polarization

Macrophages are cells of the innate immune system that have phagocytic capacity and are involved in the elimination of foreign particles from the body (Gordon and Martinez-Pomares, 2017; Uribe-Querol and Rosales, 2020) and in the presentation of antigens, constituting an important link between innate and adaptive immunity. These cells are part of the mononuclear phagocytic system and are implicated in tissue homeostasis and various infectious and inflammatory processes (Rahman *et al.*, 2017; Shapouri-Moghaddam *et al.*, 2018).

Macrophages are activated during phagocytosis or by contact with molecular patterns associated with pathogenic microorganisms. This activation results in inflammatory responses and increased production of cytokines and/or physicochemical factors and, consequently, can differentiate into various phenotypes depending on the state and changes in the microenvironment (Schmall *et al.*, 2015; Murray, 2017). There are 2 main subtypes of macrophages classified according to the expression of their cell surface markers, production of specific factors and biological activities: classically activated or inflammatory M1 macrophages and alternatively activated or anti-inflammatory M2 macrophages (Parisi *et al.*, 2018) (Fig. 3). Macrophage subtypes play a role in the initiation and/or progression of many diseases. The M1/M2 paradigm emerged as homologous with the one previously described for Th response profiles, which also presents 2 subtypes: Th cell type 1 (Th1) and type 2 (Th2) (Mills, 2015).

M1 macrophage subtypes polarize in the presence of Th1 cytokines such as IFN- γ and TNF- α or when exposed to inflammatory molecules such as lipopolysaccharides (LPS) (Yunna *et al.*, 2020),




Fig. 3. Different macrophage phenotypes, specific stimuli and markers. Source: Created with BioRender.com.

through the following mechanisms: (1) JAK/STAT (Janus kinase/signal transducer and transcriptional activator) signalling pathway. IFN- γ activates JAK-inducing phosphorylation of STAT1, which in turn leads to macrophage polarization to M1 (Wang *et al.*, 2020); (2) Toll-like receptor (TLR) 4/nuclear factor κ B (NF- κ B) signalling pathway. LPS binds to TLR4 to activate NF- κ B and activator protein 1 (AP-1), promoting the expression of inflammatory factors (Chen *et al.*, 2017b; Ciesielska *et al.*, 2021) and (3) cytokine signalling through specific receptors that activate AP-1 (Liu *et al.*, 2014).

M1 macrophages are recruited soon after lesion formation and are mainly involved in the initial response to infectious processes (Vannella and Wynn, 2017). These increase local inflammation, producing large amounts of pro-inflammatory cytokines, including IL-1 β , IL-6, IL-12, IL-18, IL-23, TNF- α and IFN type 1 (Shapouri-Moghaddam *et al.*, 2018), as shown in Fig. 3. The M1 macrophage phenotype expresses high levels of inducible nitric oxide synthase (iNOS), major histocompatibility complex class II (MHC II), CD16/32, CD80 and CD86, as well as chemokines that attract Th1 cells, including CXCL9 and CXCL12 (Orecchioni *et al.*, 2019). Functionally, M1 macrophages are characterized by antimicrobial and antitumour activities and participate in the elimination of infectious agents through the activation of nicotinamide adenine dinucleotide phosphate (NADPH) oxidase and, consequently, the generation of reactive oxygen species (ROS) (Murray *et al.*, 2014).

On the other hand, M2 macrophages are induced by Th2 cytokines IL-4 and IL-13 (Fig. 3), mainly via STAT6 activation (Sica and Mantovani, 2012; Enderlin *et al.*, 2014). This pathway is extremely important, as IL-4 inhibits M1 and induces M2 polarization (He *et al.*, 2020). Gao *et al.* (2015) demonstrated that the expression of STAT6 was positively regulated by curcumin and by the secretion of IL-4 and IL-13, capable of inducing M0 and M1 macrophages to polarize into M2. IL-4 type I and type II receptors also activate STAT6 (Gong *et al.*, 2017), which in turn induces the transcription of typical M2 polarization genes, such as mannose receptor 1, type α resistin (Retnla) and chitinase 3-like 3 (Chi3l3, Ym1) (Martinez and Gordon, 2014). M2 polarization can also be induced by IL-10 through STAT3 activation (Yin *et al.*, 2018). However, the STAT6 pathway is considered to activate M2 macrophages (Murray, 2017).

The M2 macrophage phenotype has a profile of anti-inflammatory cytokines, characterized by low production of IL-1, IL-6 and TNF- α , and high production of IL-10 and transforming growth factor-beta (TGF- β) (Fig. 3), as well as chemokines CCL1, CCL17, CCL18, CCL22 and CCL24 (Yunna *et al.*, 2020). Additionally, this phenotype can be characterized by the expression of arginase 1 (Arg-1), CD163, CD209 and CD206. CD206 interacts with glycoproteins and glycolipids found on the surfaces of pathogens (Suzuki *et al.*, 2018; Xu *et al.*, 2019). Thus, CD206 plays a role in immunological recognition of pathogens after antigen internalization and presentation (Hussell and Bell, 2014). Functionally, M2 macrophages can inhibit inflammation, promote tissue repair and wound healing, prevent parasitic infection and have proangiogenic and profibrotic properties (Jetten *et al.*, 2014; Braga *et al.*, 2015). Furthermore, because M2 macrophages produce complex cytokines and are characterized by the functional expression of alternative activation markers, they can be divided into 4 subtypes: M2a, M2b, M2c and M2d (Yao *et al.*, 2019). These subtypes differ from each other based on their cell surface markers, secreted cytokines and biological functions, as is demonstrated in Fig. 3.

M2a macrophages are induced by the cytokine IL-4 or IL-13 and express high levels of CD86, CD200R and MHC II and low levels of CD14 and TLR4 (Yao *et al.*, 2019). In addition to being major producers of CCL24, CCL17 and CCL22, they use

CCR3 and CCR4 receptors, resulting in the recruitment of eosinophils, basophils and Th2 cells, promoting the upregulation of a type 2 immune response (Fraternale *et al.*, 2015). M2b-type macrophages are induced by immune complexes, LPS or IL-1 receptor antagonist and are characterized by increased expression of CD206 and CD86 (Viola *et al.*, 2019). Upon activation, this subtype secretes pro- and anti-inflammatory cytokines TNF- α , IL-1 β , IL-6 and IL-10 and functions in regulating the immune response and inflammation (Wang *et al.*, 2019). M2c macrophages are induced by IL-10, TGF- β or glucocorticoids, and express CD206 and CD163, in addition to secreting IL-10, TGF- β , CCL16 and CCL18, which play crucial roles in the phagocytosis of apoptotic cells (Ross *et al.*, 2021). Finally, induced by TLR antagonists, M2d macrophages express high levels of CD206, IL-10 and iNOS, secrete CCL5, CXCL10 and CXCL6 and express low levels of IL-12 and TNF- α (Viola *et al.*, 2019). This subtype also secretes the vascular endothelial growth factor and promotes angiogenesis and tumour progression (Ferrante *et al.*, 2013). Notably, all subtypes of M2 macrophages express IL-10.

M1 and M2 macrophages can also be differentiated by the way they metabolize arginine, as shown in Fig. 3. M1 macrophages metabolize arginine by the enzyme iNOS to produce nitric oxide (NO) and citrulline; on the contrary, M2 macrophages metabolize arginine by Arg-1 to produce L-ornithine and urea, a precursor molecule of polyamines involved in tissue repair and cell proliferation (Rath *et al.*, 2014; Yang and Ming, 2014).

The 2 macrophage populations must be balanced to maintain homeostasis and to protect the organism. Once an imbalance occurs, the exacerbated activity of M1 or M2 macrophages can lead to the development of inflammatory diseases or host immunosuppression (Sica *et al.*, 2015). However, the remarkable plasticity of macrophages confers significant benefits to the host, especially in the course of chronic helminth infections (Lechner *et al.*, 2021) since it limits excessive tissue damage when it is unable to overcome the initial injury. This feature has been well-documented in schistosomiasis.

Participation of M1 and M2 macrophages in the response to *Schistosoma* infection

Initially, blood monocytes differentiate into macrophages at inflammatory sites (Rückerl and Cook, 2019) and exhibit high plasticity as a result of exposure to various stimuli, signalling molecules, nutrients and metabolites in the context of schistosomiasis (Cortes-Selva and Fairfax, 2021). These phagocytes can exert pro-inflammatory or anti-inflammatory functions (Zhu *et al.*, 2014) in different clinical forms of schistosomiasis (acute and chronic phases) (Fig. 2). In the acute phase, macrophages secrete pro-inflammatory cytokines and consequently increase inflammation, recruit more immune cells and promote the formation of the initial granuloma. In the chronic phase, macrophages have an immunoregulatory activity to decrease the damage caused by granulomas (Wolde *et al.*, 2020).

During the life cycle of *S. mansoni* and *S. japonicum*, several antigens are excreted by their different evolutive forms (Curwen *et al.*, 2006; Jang-Lee *et al.*, 2007; Acharya *et al.*, 2021). For example, Sm16 – a low molecular weight protein that is secreted by *S. mansoni* cercariae, helps the parasite to enter the host's skin (Brännström *et al.*, 2009; Sanin and Mountford, 2015). Sm29, present in the tegument of schistosomula and adult *S. mansoni* worms, can induce the maturation and activation of human monocyte-derived DCs (Cardoso *et al.*, 2008; Lopes *et al.*, 2019). Sj-C is an example of a protein secreted from the tegument of *S. japonicum*, which may suppress the presentation of exogenous antigens by DCs (He *et al.*, 2011; Chen *et al.*, 2017a). IPSE/

Table 1. Molecules and/or antigens involved in macrophage polarization in *Schistosoma mansoni* infection

Macrophage profile	Molecule/antigen	Experimental model	Type of study	References
M1	Schistosomules	C57BL/6 mice	<i>In vivo</i>	Menson and Wilson (1990)
M2	SEA	BMDCs and DCs from C57BL/6 or TLR4 mice Macrophages derived from human monocytes	<i>In vitro</i>	Goh et al. (2009)
M2	p16 ^{INK4a}	Macrophages derived from mouse bone marrow Chimaeric mice	<i>In vitro</i> <i>In vivo</i>	Cudejko et al. (2011)
M2	Cercariae	Peritoneal macrophages Mice IL4Rα ^{flox/Δ} LysM ^{WT/Cre}	<i>In vitro</i> <i>In vivo</i>	Vanella et al. (2014)
	SEA	Mutant mice LysM ^{Cre/+} Shp2 ^{flox/flox} (control) and LysM ^{Cre/+} : Shp2 ^{flox/flox} (Shp2 ^{Δ/Δ})		
M2	CD14 TLR co-receptor	Mice <i>Cd14</i> ^{-/-} Wild-type mice (wt)	<i>In vivo</i>	Tundup et al. (2014)
M1 profile lock	Sm16 antigen	Macrophages derived from mouse bone marrow	<i>In vitro</i>	Sanin and Mountford (2015)
M2	LPC	Peritoneal macrophages and bone marrow derivatives of mice	<i>In vitro</i>	Assunção et al. (2017)
M2	IPSE/α-1	Human peripheral blood cells (monocytes and basophils)	<i>In vitro</i>	Knuhr et al. (2018)

α-1 and ω-1 are examples of proteins secreted by *S. mansoni* eggs, which help direct a Th2 response (Everts et al., 2009; Knuhr et al., 2018). These molecules can induce the activation and modulation of innate and adaptive immune responses and facilitate the evasion of the parasite from the host-defense mechanisms (Jenkins et al., 2005a, 2005b; Hai et al., 2014; Hambrook and Hanington, 2021). *Schistosoma* antigens can be proteins (such as enzymes), polysaccharides and the most commonly used are crude extracts prepared by breaking up worms, larvae or eggs (Doenhoff et al., 1993; Doenhoff, 1998). Thus, it is clear that antigen changes in the microenvironment during schistosomiasis are important for the polarization of macrophages to the M1 or M2 profile (Xu et al., 2014; Sanin and Mountford, 2015; Assunção et al., 2017).

According to Tables 1 and 2, we highlight some *in vitro* and *in vivo* studies that demonstrate the relationship between the stimulation of *S. japonicum* and *S. mansoni* antigens and macrophage polarization. In addition, we also highlight other molecules involved in macrophage polarization in schistosomiasis, providing molecular evidence of great relevance in the process of differentiation of these cells, which will be discussed in this article.

Cercariae and schistosomula antigens can induce an M1 profile

During the penetrating of the human skin, cercariae of *S. mansoni* and *S. japonicum* release E/S products, which have remodelling and immunoregulatory functions (Liu et al., 2015; Sanin and Mountford, 2015), that facilitate their penetration and subsequent establishment in the host's body, in the form of schistosomula (Janssen et al., 2016). This phase represents the first contact with innate immune responses in the skin, especially Langerhans cells, which are considered tissue-resident macrophages (West and Bennett, 2018). These cells phagocytize E/S and secrete pro-inflammatory (IL-6 and IL-12p40) and anti-inflammatory (IL-10) cytokines in a TLR-dependent manner (Jenkins et al., 2005a, 2005b).

One of the ways in which macrophages are activated is through the action of TLRs. These receptors are a family of pattern recognition receptors that are important for innate immune response (El-Zayat et al., 2019). These receptors recognize invading pathogens, trigger innate immune responses and subsequently initiate adaptive immunity against infections, including Gram-positive and Gram-negative bacteria, fungi, viruses and parasites (Lu et al., 2018). These receptors mediate macrophage recognition by microbial ligands, inducing the expression of microbicidal molecules and cytokines via the adapter protein MyD88 (Jin et al., 2019). Xu et al. (2014) showed that normal cercariae antigen (NCA) and attenuated cercariae antigen (ACA) from *S. japonicum* induced polarization to the M1 profile, with increased levels of IL-12, CD136/32 and iNOS (Table 2). However, these values decreased when the TLR4 pathway blockers were used. Thus, the authors suggested that the polarization of the M1 profile is dependent on the TLR4 pathway and this may play a protective role in *S. japonicum* infection (Tang et al., 2021).

In fact, the TLR4 pathway is extremely important for the polarization of macrophages to the M1 phenotype, as demonstrated in some studies (Freitas et al., 2016; Shi et al., 2020). Sanin and Mountford (2015) demonstrated that Sm16 (a molecule produced by *S. mansoni* cercariae) is able to block TLR4 and TLR3 pathways in human monocyte, which negatively affect the classic activation of macrophages (M1) in response to IFN-γ (Table 1). This is considered an important mechanism of immune evasion promoted by *S. mansoni* because it limits the production of NO, which is toxic to the parasite (Shiels et al., 2020).

After complete transformation from a cercariae into a schistosomula, the larva migrates into the bloodstream, travelling through the lungs until reaching maturation in the mesenteric veins. This stage of the cycle is also characterized as a key target for the elimination of infection through innate host immune responses (Houlder et al., 2021). Some histological studies conducted in the lungs of mice infected with *S. mansoni* and *S. japonicum* showed inflammatory foci consisting of neutrophils, eosinophils and macrophages (Crabtree and Wilson, 1986; Burke et al., 2011).

Table 2. Molecules and/or antigens involved in macrophage polarization in *Schistosoma japonicum* infection

Macrophage profile	Molecule/antigen	Experimental model	Type of study	References
M1/M2	NCA	Macrophages RAW264.7	<i>In vitro</i>	Xu <i>et al.</i> (2014)
	ACA			
	SWAP	Mice	<i>In vivo</i>	
	SEA	C57BL/6J		
M1/M2	SWA	Peritoneal macrophages of mice	<i>In vitro</i>	Zhu <i>et al.</i> (2014)
	SEA			
M1	EVs	Macrophages RAW264.7	<i>In vitro</i>	Wang <i>et al.</i> (2015)
M2	Corilagin	Ana-1 cell line	<i>In vitro</i>	Li <i>et al.</i> (2017)
		C57BL/6 mice	<i>In vivo</i>	
M2b	Egg-derived ES antigens	Macrophages derived from bone marrow of wild-type mice and TLR2 ^{-/-}	<i>In vitro</i>	Gong <i>et al.</i> (2018)
		C57BL/6 mice	<i>In vivo</i>	
M2	Sj16	Mouse peritoneal macrophages	<i>In vitro</i>	Shen <i>et al.</i> (2019)
		BALB/c mice	<i>In vivo</i>	
M1	EV miRNAs EVs	Macrophages RAW264.7	<i>In vitro</i>	Liu <i>et al.</i> (2019)
		BALB/c mice and rabbits	<i>In vivo</i>	
M2	Eggs	Mouse peritoneal macrophages	<i>In vitro</i>	Ye <i>et al.</i> (2020)
		Mice	<i>In vivo</i>	
		Kunming		
M1	SWA	Macrophages RAW264.7	<i>In vitro</i>	Shen <i>et al.</i> (2021)
		C57BL/6 mice	<i>In vivo</i>	
M2	SEA	Macrophages J774A.1	<i>In vitro</i>	Yu <i>et al.</i> (2021)

Macrophages function as cytotoxic cells, mainly in schistosomula (James and Glaven, 1989). Oswald *et al.* (1994) demonstrated that macrophages could produce NO, leading to schistosomula death in animal models independent of the production of pro-inflammatory cytokines. In contrast, Cardoso *et al.* (2008) determined that the antigen Sm29, present in the integument of *S. mansoni* schistosomula, induced a Th1-type immune response, with an increase in pro-inflammatory cytokines (IFN- γ , TNF and IL-12) in mice, leading to a reduction in worm burden and liver pathology. James *et al.* (1998) demonstrated that IFN- γ was a cytokine of great importance in the activation of macrophages in the lungs for the immunological killing of *S. mansoni* larvae and played a critical role in protective immunity.

In the early stages of schistosomiasis, lung macrophages may have an M1 phenotypic trait. Menson and Wilson (1990) characterized the expression of surface markers in alveolar macrophages associated with the immune response to *S. mansoni*. The authors demonstrated an increase in IFN- γ expression in the lungs of C57BL/6 mice and suggested that activated macrophages might be responsible for initiating and maintaining focal inflammation that blocks parasite migration (Table 1).

Worm antigens can induce an M1 or M2 profile

During the acute phase of *Schistosoma* infection, before parasite oviposition (approximately 5–7 weeks post-infection), immune responses are largely of the CD4⁺ Th1 type, associated with increased numbers of M1 macrophages that produce IL-12, IL-6, TNF- α and NO (Pearce *et al.*, 1991; Coulson *et al.*, 1998;

Gordon, 2003). These early pro-inflammatory responses are mainly related to the antigens from immature worms (schistosomula) during their migration (Wilson, 1998; Egesa *et al.*, 2018). Activation of these responses may be through binding to TLR and C-type lectin receptors on macrophages; however, further studies are needed to clarify this mechanism of macrophage activation *via* schistosomula antigens.

In contrast to the schistosomula antigens, the adult worm antigen preparations [soluble worm antigen (SWAP or SWA)] were better explored in experimental studies. Although the antigenic composition is not the same as the live worm, the use of SWAP or SWA constitutes a valuable experimental tool to evaluate many aspects of immune responses promoted by different host cells (Xu *et al.*, 2014; Zhu *et al.*, 2014). This antigen is the easiest to obtain and is essentially an extract based on Tris-HCl or phosphate-buffered saline from mixed male and female worms and prepared in various ways, either by homogenization, sonication or freeze/thaw (or a combination of these) (Grenfell *et al.*, 2012; Neves *et al.*, 2015). Some studies have demonstrated that SWAP could induce an M1-like profile (Xu *et al.*, 2014; Zhu *et al.*, 2014). Thus, Zhu *et al.* (2014), when performing a co-culture of peritoneal macrophages obtained from mice with *S. japonicum* SWA (Table 2), observed that there was an increase in the expression of specific markers related to M1 (TNF- α , IL-12, CXCL9, CXCL10, CXCL11 and iNOS).

Aiming to understand which mechanisms lead SWAP to induce polarization of the M1 profile, Shen *et al.* (2021) demonstrated that this antigen promoted the expression of a protein called lipocalin 2 (LCN2) and, consequently, induced the M1 profile of macrophages (Table 2) through the upregulation of the

NF- κ B signalling pathway. It has already been reported that this protein is increased in macrophages and can potentiate the M1 phenotype of microglia in the central nervous system (Jang *et al.*, 2013). The NF- κ B signalling pathway can activate macrophages to produce M1 polarization upon LPS induction (Liu *et al.*, 2017). In addition, some studies have shown that this pathway could regulate the expression of LCN2, thereby stimulating the inflammatory response in infectious processes (Zhao and Stephens, 2014; Ghosh *et al.*, 2017).

In addition to the antigens of adult worms that induce an M1 profile, studies have shown that adult worms of *S. mansoni* and *S. japonicum* also release extracellular vesicles (EVs), known as exosomes, which modulate the host immune response (Nowacki *et al.*, 2015; Wang *et al.*, 2015; Sotillo *et al.*, 2016; Zhu *et al.*, 2016). Exosomes are membrane-bound vesicles secreted by various types of mammalian cells in normal and diseased states (Avni and Avni, 2021). Exosomes play an important role in cell-cell communication and have been implicated in the regulation of cell development, immune regulation, angiogenesis and cell migration (Raposo and Stoorvogel, 2013; Zhu *et al.*, 2016). Wang *et al.* (2015) observed that RAW264.7 macrophages, when cultured with exosome-like vesicles isolated from *S. japonicum*, exhibited an M1 profile (Table 2), due to the increase in the surface markers CD16/32, iNOS and TNF- α . Liu *et al.* (2019) investigated miRNAs from *S. japonicum* EVs and found that they increased macrophage proliferation *in vitro* (RAW264.7) and *in vivo* (mice and rabbits) as well as TNF- α expression. miRNAs are involved in the regulation of the development, differentiation and activation of immune cells, including macrophages (Montagner *et al.*, 2014; Mehta and Baltimore, 2016). Thus, the polarization of M1 induced by schistosome EVs may represent an important mechanism for parasite survival in vertebrate hosts, *via* modulation of the immune response. However, there are still controversies about the possible role of the schistosome tegument as a source of EVs, because, to date, no study has been performed to prove the exact origin of these vesicles (Wilson and Jones, 2021).

On the other hand, adult worm products can also bias the M2 profile (Smith *et al.*, 2018). Indeed, Xu *et al.* (2014) showed that adult *S. japonicum* worms could induce an M2 macrophage profile. The authors, when stimulating RAW264.7 macrophages with SWAP from *S. japonicum*, observed an increase in the expression of surface markers (CD16/32 and CD206) and in the production of cytokines (IL-12 and IL-10), suggesting that this antigen could induce both M1 and M2 macrophage profiles. The potential explanation for this could be related to how the antigens of adult worms were obtained since some adult female worms possess eggs in the process of maturation into their uterus/ootype, and consequently, this antigen could have been contaminated with SEAs. However, further studies are needed to understand macrophage polarization by SWAP and its relationship with SEA contamination.

Besides the classical macrophage polarization (M1 and M2), products excreted by schistosomes, such as haemozoin, are also able to induce immunomodulation. Adult worms of *S. mansoni* acquire nutrients by haematophagy of the host's blood, and this process can form toxic haem for the parasite (Zussman *et al.*, 1970). However, the schistosomes are able to neutralize the free haem in their intestine through crystallization in haemozoin (Oliveira *et al.*, 2000). This haemozoin is regurgitated by the worms into the host bloodstream and can be accumulated in the liver (Kloetzel and Lewert, 1966), which may activate the immune response of the host. From this perspective, a previous study (Truscott *et al.*, 2013) highlighted that haemozoin formed from *S. mansoni* is able to maintain the M2 macrophage profile previously activated by IL-4 stimulation, but also exerts specific

modulatory effects on these cells (Table 1). These authors showed that haemozoin mediated the suppression of *Retnla* (resistin-like molecule- α or *Fizz1*) expression and *Retnla* protein secretion in the M2 macrophages. The role of *Retnla* during experimental schistosomiasis is associated with the limitation of Th2 inflammatory response (Pesce *et al.*, 2009). However, further studies are necessary to better explain the possible impact of haemozoin in the immunopathology of schistosomiasis.

SEAs can induce an M2 profile

After maturation of the adult worms and subsequent oviposition, the activation of a type 2 profile begins in response to the soluble antigens secreted by the eggs of *S. mansoni* and *S. japonicum* (Tables 1 and 2) (Cheever *et al.*, 2000; Pearce and MacDonald, 2002; Pearce *et al.*, 2004; Burke *et al.*, 2009; Costain *et al.*, 2018). The type 2 profile of schistosomiasis is characterized by the expansion of Th2 cells, eosinophils and basophils, and increased production of IL-4, IL-5 and IL-13 (Hams *et al.*, 2013; Schwartz *et al.*, 2014), as previously described. IL-4 and IL-13 protect hosts against various helminth parasites by signalling through the IL-4R α chain (Barron and Wynn, 2011; Jenkins *et al.*, 2011). The production of these cytokines reduces the inflammation levels produced by the type 1 profile of the initial stage of acute phase, preventing acute pathology, such as intestinal haemorrhage and liver damage; however, the Th2 immune response is responsible for the formation of hepatic and intestinal granulomas (Brunet *et al.*, 1997; Hams *et al.*, 2013; Zheng *et al.*, 2020).

Granulomas are essential for sequestering toxic antigens produced by eggs and preventing further tissue damage. However, if unregulated by the immune response of the host, granulomas grow excessively and progress to fibrotic stages, which are responsible for severe forms of the disease, such as cirrhosis, portal hypertension, liver failure and even host death (Lenzi *et al.*, 1998; Cheever *et al.*, 2000; Takaki *et al.*, 2021). Macrophages are one of the main cellular components of hepatic granulomas (Beljaars *et al.*, 2014; Schwartz and Fallon, 2018). Recent studies have demonstrated that M2 macrophages play a direct and critical role in fibrosis, granuloma maintenance, tissue repair and host survival (Cortes-Selva *et al.*, 2018; Song *et al.*, 2020). Ye *et al.* (2020) showed that M2 macrophage markers (CD200R, Arg-1 and Ym1) were increased in the liver, spleen, large intestine and peritoneal macrophages of *S. japonicum*-infected mice. Jenkins *et al.* (2011) observed that IL-4/IL-13 signalling *via* IL-4R α induces an alternative phenotype in resident macrophages. In this sense, a study performed with macrophages derived from the bone marrow of mice infected with *S. mansoni* demonstrated that the tumour suppressor gene p16^{INK4a} was an excellent modulator of the activation and polarization of macrophages induced by IL-4 through the JAK2-STAT1 pathway (Cudejko *et al.*, 2011).

Egg antigens induce granulomas, consisting mainly of M2 macrophages (Yu *et al.*, 2021). Zhu *et al.* (2014) showed that peritoneal macrophages obtained from healthy mice, when stimulated with *S. japonicum* SEAs, expressed high levels of chemokines (CCL2, CCL17 and CCL22), IL-10 and Arg-1. Similarly, Xu *et al.* (2014), after stimulating RAW264.7 macrophages with *S. japonicum* SEAs, also observed higher levels of IL-10. In chronic schistosomiasis infection, the main function of IL-10 is to control liver damage and regulate antifibrotic processes (Dewals *et al.*, 2010; Kamdem *et al.*, 2018). Previous studies have shown that low levels of IL-10 expression are related to liver fibrosis in *S. mansoni*-infected patients (Mutengo *et al.*, 2018). On the other hand, little is known about the mechanisms by which an SEA preferentially induces M2 macrophage differentiation.

Previous studies have demonstrated that an SEA from *S. mansoni* could induce the expression of the notch Jagged1 ligand in mice and human macrophages, suggesting that Jagged1 might have a specific role in the M2 polarization process of macrophages (Goh *et al.*, 2009) (Table 1). Macrophages found in liver tissues exhibit functional M2 polarization, which is dependent on the activation of notch1/Jagged1 signalling (Zheng *et al.*, 2016).

During *S. mansoni* infection, basophils detect egg IPSE/ α -1 glycoprotein and stimulate the production of IL-4 and IL-13, which trigger the alternative activation of human monocytes (Table 1), as observed by the increased expression of CD206 and CD209 (Knuhr *et al.*, 2018). IL-13 is a key cytokine that induces M2 macrophage polarization via the IL-13 α 1 signalling pathway (Chiaromonte *et al.*, 1999; Liu *et al.*, 2012). Li *et al.* (2017) performed a study with corilagin, an active component of many medicinal plants, and found that this component could suppress *Schistosoma* egg-induced liver fibrosis by inhibiting M2 macrophage polarization (Table 1) in the IL-13R α 1 signalling pathway. Corilagin has great potential to reduce liver fibrosis caused by egg antigens in *S. japonicum* infection by decreasing the expression of molecules associated with the IL-13/STAT6 signalling pathway in liver M2 macrophages (Du *et al.*, 2016).

Signaling via TLR2 may be another way egg antigens polarize M2 macrophages during schistosomiasis. Gong *et al.* (2018) showed that antigens derived from *S. japonicum* eggs could activate macrophages, which exhibit M2b polarization dependent on NF- κ B signalling, mediated by the MyD88/mitogen-activated protein kinase (MAPK) pathway in a TLR2-dependent manner (Table 1). In contrast, Tundup *et al.* (2014) showed that the CD14 TLR co-receptor was upregulated in hepatic macrophages after *S. mansoni* infection and acted as a crucial negative regulator of M2 polarization, possibly as part of a parasitic defense mechanism against granuloma formation (Table 1). Gao *et al.* (2013) observed that an SEA of *S. japonicum*, known as SJEA, upregulated programmed death ligand 2 (PD-L2) expression in mouse bone marrow-derived macrophages (BMDCs) via TLR2, which binds PD-1 primarily on CD4⁺ T cells. This mechanism can help inhibit the T cell response during *S. japonicum* infection.

Lysophosphatidylcholine (LPC) from *S. mansoni* eggs can also induce macrophage differentiation into the M2 phenotype (Assunção *et al.*, 2017), as shown in Table 1. The authors observed that LPC from *S. mansoni* activates peroxisome proliferator-activated receptor gamma (PPAR- γ), a transcription factor necessary for M2 polarization, leading to higher expression of Arg-1 and CD206, while increasing the production of IL-10, TGF- β and PGE2 in peritoneal macrophages *in vitro*. *Schistosoma mansoni* eggs induced a 7-fold increase in PPAR- γ expression in human liver cell cultures (Anthony *et al.*, 2010). PPAR- γ , in addition to being of great importance in M2 polarization, can regulate lipid uptake and metabolism (Ahmadian *et al.*, 2013; Abdalla *et al.*, 2020).

Fang *et al.* (2015) showed that BMDCs from C57BL/6 mice, when stimulated with a specific *S. japonicum* egg protein known as SJE16.7, promoted the production of pro- (IL-12, IL-6 and TNF- α) and anti-inflammatory (IL-10) cytokines through the phosphorylation of MAPKs and increased the expression of MHC II on the surface of macrophages. Previous studies have shown that *S. mansoni* and *S. japonicum* egg antigens could stimulate the MAPK pathway in macrophages (Wang *et al.*, 2006; de Andrade *et al.*, 2014). MAPKs are essential transmitters of extracellular signals that can mediate key cellular processes, including cell differentiation, division and death (Yang *et al.*, 2003). Thus, SJE16.7 is a potent macrophage activator. However, in another study, Shen *et al.* (2019), when using the Sjl6 antigen, noticed that it decreased hepatic granulomas in

mice infected with *S. japonicum* and associated this improvement with the suppression of cytokine production, such as IFN- γ , TNF- α , IL-4 and IL-6. The authors reported that the mechanisms of Sjl6 attenuation of hepatic granulomatous inflammation and fibrosis in these infected mice might be related to the induction of macrophages for M2 polarization (Table 2). These authors also demonstrated, by flow cytometry, the increase in the expression of CD206 after stimulation of Sjl6 in peritoneal macrophages and leucocytes from the livers of mice. Corroborating these results, Hu *et al.* (2009) showed that Sjl6 decreased the levels of pro-inflammatory cytokines, such as IL-6 and TNF- α , and increased the levels of IL-10 in RAW264.7 macrophages. Vannella *et al.* (2014) observed that mice infected with *S. mansoni* showed an increase in M2 macrophages that expressed Arg-1, which attenuated the progression of inflammation and fibrosis (Table 1). Stimulation of RAW264.7 macrophages with another *S. japonicum* egg protein (SjCP1412) also increased the expression of CD206, Arg-1 and IL-10, which are related to M2-type macrophage differentiation (Ke *et al.*, 2017). Overall, these findings emphasize that M2 macrophages are important in reducing the lesions caused by schistosomiasis through downregulation of the Th1 response and inflammation promoted by egg antigens. Additionally, the role of these cells was previously investigated in a mouse model of liver injury induced by acetaminophen (paracetamol) (Starkey Lewis *et al.*, 2020). The authors demonstrated that the injection of M2 macrophages in this experimental model was able to rapidly reduce liver damage and inflammation. These data indicate that M2 macrophages may constitute a new potential cell-based therapy for this disease. Based on this, it seems promissory also to apply these cells in the immunotherapy of schistosomiasis.

Interestingly, despite being produced by M1 macrophages, a recent study demonstrated that the production of ROS by egg antigens may be a potential mechanism for M2 macrophage differentiation (Table 2). ROS have several biological activities, such as participation in innate and adaptive immune responses, and can be cytotoxic against pathogens (Canton *et al.*, 2021). Yu *et al.* (2021) observed that a significant increase in ROS in the liver of mice infected with *S. japonicum* was related to fibrosis and the differentiation of M2 macrophages. The authors hypothesized that their findings were due to NADPH oxidase (NOX2) inhibiting SEA-stimulated ROS production in macrophages, suggesting that NOX might act as the main source of ROS production in SEA-stimulated macrophages. NADPH oxidase is the first source of ROS identified in macrophages (Nathan *et al.*, 2013). Macrophages produce large amounts of ROS, primarily through NOX2 activation (Paik *et al.*, 2014). Thus, the production of ROS induced by schistosome eggs may be a target for the treatment of schistosomiasis.

Future perspectives and final considerations

Findings about the mechanisms behind macrophage activation during different metabolic profiles in human diseases present an exciting prospect, as there are pathologies that have been associated with a particular macrophage phenotype. In this context, the polarization of macrophages in schistosomiasis and their consequent ability to promote an effective immune response seem to be an attractive therapeutic approach associated with conventional chemotherapy treatments.

Overall, the findings highlighted in this review demonstrate the relevance and complexity of understanding the mechanisms involved in macrophage polarization (M1/M2) in schistosomiasis. The *S. japonicum* and *S. mansoni* antigens in macrophage polarization are particularly important in this process. These products have been shown to have immunomodulatory effects in different

phases of schistosomiasis and are seen as potential therapeutic targets for this disease, especially in the chronic phase. Among the potential therapeutics, the combination of different schistosome antigens can result in higher levels of host protection, stimulating an adequate immune response for either an M1 or M2 profile; however, this can only be achieved after many *in vitro* and *in vivo* experiments.

Acknowledgements. We acknowledge the Graduate Program in Health Sciences at the Federal University of Maranhão for providing the support. We would like to acknowledge Editage (www.editage.com) for English-language editing.

Author's contributions. I. C. L. L.: investigation, conceptualization and writing of the original draft; I. C. L. L., G. C. C. G. F., R. A. N., M. G. S. L., V. A. F. S., J. G. M. R., G. S. M., R. C. C., L. A. S., R. N. M. G. and F. R. F. N.: review and editing of original draft. All the authors reviewed and approved the final version of this manuscript.

Financial support. I. C. L. L., G. C. C. G. F., R. A. N. and V. A. F. S. were supported by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior doctoral fellowship. R. N. M. G. and F. R. F. N. were supported by CNPq researcher fellowship. The research was supported by FAPEMA (Universal 01001/19) and by CAPES (Financial code 001).

Conflict of interest. The authors declare no conflict of interest.

References

- Abdalla HB, Napimoga MH, Lopes AH, de Macedo Maganin AG, Cunha TM, Van Dyke TE and Clemente Napimoga JT (2020) Activation of PPAR- γ induces macrophage polarization and reduces neutrophil migration mediated by heme oxygenase 1. *International Immunopharmacology* **84**, 106565.
- Abdel-Ghany R, Rabia I, El-Ahwany E, Saber S, Gamal R, Nagy F, Mahmoud O, Hamad RS and Barakat W (2015) Blockade of PGE2, PGD2 receptors confers protection against prepatent *Schistosomiasis mansoni* in mice. *Journal of the Egyptian Society of Parasitology* **45**, 511–520.
- Acharya S, Da'dara AA and Skelly PJ (2021) Schistosome immunomodulators. *PLoS Pathogens* **17**, e1010064.
- Ahmadian M, Suh JM, Hah N, Liddle C, Atkins AR, Downes M and Evans RM (2013) PPAR γ signaling and metabolism: the good, the bad and the future. *Nature Medicine* **19**, 557–566.
- Angeli V, Faveeuw C, Roye O, Fontaine J, Teissier E, Capron A, Wolowezuk I, Capron M and Trottein F (2001) Role of the parasite-derived prostaglandin D2 in the inhibition of epidermal Langerhans cell migration during schistosomiasis infection. *The Journal of Experimental Medicine* **193**, 1135–1147.
- Anthony B, Mathieson W, de Castro-Borges W and Allen J (2010) *Schistosoma mansoni*: egg-induced downregulation of hepatic stellate cell activation and fibrogenesis. *Experimental Parasitology* **124**, 409–420.
- Assunção LS, Magalhães KG, Carneiro AB, Molinaro R, Almeida PE, Atella GC, Castro-Faria-Neto HC and Bozza PT (2017) Schistosomal-derived lysophosphatidylcholine triggers M2 polarization of macrophages through PPAR γ dependent mechanisms. *Biochimica et Biophysica Acta. Molecular and Cell Biology of Lipids* **1862**, 246–254.
- Atri C, Guerfali FZ and Laouini D (2018) Role of human macrophage polarization in inflammation during infectious diseases. *International Journal of Molecular Sciences* **19**, 1801.
- Avni D and Avni O (2021) Extracellular vesicles: schistosomal long-range precise weapon to manipulate the immune response. *Frontiers in Cellular and Infection Microbiology* **11**, 196.
- Bajiro M, Dana D and Levecke B (2017) Prevalence and intensity of *Schistosoma mansoni* infections among schoolchildren attending primary schools in an urban setting in southwest, ethiopia. *BMC Research Notes* **10**, 1–6.
- Barron L and Wynn TA (2011) Macrophage activation governs schistosomiasis-induced inflammation and fibrosis. *European Journal of Immunology* **41**, 2509.
- Bartlett A, Brown M, Marriott C and Whitfield PJ (2000) The infection of human skin by schistosome cercariae: studies using Franz cells. *Parasitology* **121**, 49–54.
- Bartley PB, Ramm GA, Jones MK, Ruddell RG, Li Y and McManus DP (2006) A contributory role for activated hepatic stellate cells in the dynamics of *Schistosoma japonicum* egg-induced fibrosis. *International Journal for Parasitology* **36**, 993–1001.
- Beijaars I, Schippers M, Reker-Smit C, Martinez FO, Helming I, Poelstra K and Melgert BN (2014) Hepatic localization of macrophage phenotypes during fibrogenesis and resolution of fibrosis in mice and humans. *Frontiers in Immunology* **5**, 430.
- Bottiau E, Clerinx J, de Vega MR, Van den Enden E, Colebunders R, Van Esbroeck M, Vervoort T, Van Gompel A and Van den Ende J (2006) Imported Katayama fever: clinical and biological features at presentation and during treatment. *The Journal of Infection* **52**, 339–345.
- Bourke CD, Prendergast CT, Sanin DE, Oulton TE, Hall RJ and Mountford AP (2015) Epidermal keratinocytes initiate wound healing and pro-inflammatory immune responses following percutaneous schistosome infection. *International Journal for Parasitology* **45**, 215–224.
- Braga TT, Agudelo JSH and Camara NOS (2015) Macrophages during the fibrotic process: M2 as friend and foe. *Frontiers in Immunology* **6**, 602.
- Brännström K, Sellin ME, Holmfeldt P, Brattsand M and Gullberg M (2009) The *Schistosoma mansoni* protein Sm16/SmSLP/SmSPO-1 assembles into a nine-subunit oligomer with potential to inhibit Toll-like receptor signaling. *Infection and Immunity* **77**, 1144.
- Brink LH, McLaren DJ and Smithers SR (1977) *Schistosoma mansoni*: a comparative study of artificially transformed schistosomula and schistosomula recovered after cercarial penetration of isolated skin. *Parasitology* **74**, 73–86.
- Brunet LR, Finkelman FD, Cheever AW, Kopf MA and Pearce EJ (1997) IL-4 protects against TNF-alpha-mediated cachexia and death during acute schistosomiasis. *The Journal of Immunology* **159**, 777–785.
- Burke ML, Jones MK, Gobert GN, Li YS, Ellis MK and McManus DP (2009) Immunopathogenesis of human schistosomiasis. *Parasite Immunology* **31**, 163–176.
- Burke ML, McGarvey L, McSorley HJ, Bielefeldt-Ohmann H, McManus DP and Gobert GN (2011) Migrating *Schistosoma japonicum* schistosomula induce an innate immune response and wound healing in the murine lung. *Molecular Immunology* **49**, 191–200.
- Caldas IR, Campi-Azevedo AC, Oliveira LFA, Silveira AMS, Oliveira RC and Gazzinelli G (2008) Human *Schistosomiasis mansoni*: immune responses during acute and chronic phases of the infection. *Acta Tropica* **108**, 109–117.
- Canton M, Sánchez-Rodríguez R, Spera I, Venegas FC, Favia M, Viola A and Castegna A (2021) Reactive oxygen species in macrophages: sources and targets. *Frontiers in Immunology* **12**, 734229.
- Cardoso FC, Macedo GC, Gava E, Kitten GT, Mati VL, de Melo AL, Caliari MV, Almeida GT, Venancio TM, Verjovski-Almeida S and Oliveira SC (2008) *Schistosoma mansoni* tegument protein Sm29 is able to induce a Th1-type of immune response and protection against parasite infection. *PLoS Neglected Tropical Diseases* **2**, e-308.
- Cass CL, Johnson JR, Califf LL, Xu T, Hernandez HJ, Stadecker MJ, Yates JR and Williams DL (2007) Proteomic analysis of *Schistosoma mansoni* egg secretions. *Molecular and Biochemical Parasitology* **155**, 84.
- Cervi L, MacDonald AS, Kane C, Dzierszynski F and Pearce EJ (2004) Cutting edge: dendritic cells copulsed with microbial and helminth antigens undergo modified maturation, segregate the antigens to distinct intracellular compartments, and concurrently induce microbe-specific Th1 and helminth-specific Th2 responses. *The Journal of Immunology* **172**, 2016–2020.
- Cheever AW, Hoffmann KF and Wynn TA (2000) Immunopathology of *Schistosomiasis mansoni* in mice and men. *Immunology Today* **21**, 465–466.
- Chen BL, Peng J, Li QF, Yang M, Wang Y and Chen W (2013) Exogenous bone morphogenetic protein-7 reduces hepatic fibrosis in *Schistosoma japonicum*-infected mice via transforming growth factor- β /Smad signaling. *World Journal of Gastroenterology* **19**, 1405.
- Chen L, He B, Hou W and He L (2017a) Cysteine protease inhibitor of *Schistosoma japonicum* – a parasite-derived negative immunoregulatory factor. *Parasitology Research* **116**, 901–908.
- Chen XX, Tang L, Fu YM, Wang Y, Han ZH and Meng JG (2017b) Paralemmin-3 contributes to lipopolysaccharide-induced inflammatory response and is involved in lipopolysaccharide-Toll-like receptor-4 signaling in alveolar macrophages. *International Journal of Molecular Medicine* **40**, 1921–1931.
- Chiaromonte MG, Schopf LR, Neben TY, Cheever AW, Donaldson DD and Wynn TA (1999) IL-13 is a key regulatory cytokine for Th2 cell-mediated pulmonary granuloma formation and IgE responses induced by *Schistosoma mansoni* eggs. *Journal of Immunology* **162**, 920–930.

- Ciesielska A, Matyjek M and Kwiatkowska K (2021) TLR4 and CD14 trafficking and its influence on LPS-induced pro-inflammatory signaling. *Cellular and Molecular Life Sciences* **78**, 1233–1261.
- Colley DG and Secor WE (2014) Immunology of human schistosomiasis. *Parasite Immunology* **36**, 347–357.
- Colley DG, Bustinduy AL, Secor WE and King CH (2014) Human schistosomiasis. *The Lancet* **383**, 2253–2264.
- Cortes-Selva D and Fairfax K (2021) Schistosome and intestinal helminth modulation of macrophage immunometabolism. *Immunology* **162**, 123–134.
- Cortes-Selva D, Elvington AF, Ready A, Rajwa B, Pearce EJ, Randolph GJ and Fairfax KC (2018) *Schistosoma mansoni* infection-induced transcriptional changes in hepatic macrophage metabolism correlate with an athero-protective phenotype. *Frontiers in Immunology* **9**, 2580.
- Costain AH, MacDonald AS and Smits HH (2018) Schistosome egg migration: mechanisms, pathogenesis and host immune responses. *Frontiers in Immunology* **9**, 3042.
- Coulson PS, Smythies LE, Betts C, Mabbott NA, Sternberg JM, Wei XG, Liew FY and Wilson RA (1998) Nitric oxide produced in the lungs of mice immunized with the radiation-attenuated schistosome vaccine is not the major agent causing challenge parasite elimination. *Immunology* **93**, 55–63.
- Crabtree JE and Wilson RA (1986) The role of pulmonary cellular reactions in the resistance of vaccinated mice to *Schistosoma mansoni*. *Parasite Immunology* **8**, 265–285.
- Cudejko C, Wouters K, Fuentes L, Hannou SA, Paquet C, Bantubungi K, Bouchaert E, Vanhoutte J, Fleury S, Remy P, Tailleur A, Chinetti-Gbaguidi G, Dombrowicz D, Staels B and Paumelle R (2011) p16^{INK4a} deficiency promotes IL-4-induced polarization and inhibits proinflammatory signaling in macrophages. *Blood* **118**, 2556–2566.
- Curwen RS and Wilson RA (2003) Invasion of skin by schistosome cercariae: some neglected facts. *Trends in Parasitology* **19**, 63–66.
- Curwen RS, Ashton PD, Sundaralingam S and Wilson RA (2006) Identification of novel proteases and immunomodulators in the secretions of schistosome cercariae that facilitate host entry. *Molecular and Cellular Proteomics* **5**, 835–844.
- de Andrade LF, de Mourão MM, Geraldo JA, Coelho FS, Silva LL, Neves RH, Volpini A, Machado-Silva JR, Araujo N, Nacif-Pimenta R, Caffrey CR and Oliveira G (2014) Regulation of *Schistosoma mansoni* development and reproduction by the mitogen-activated protein kinase signaling pathway. *PLoS Neglected Tropical Diseases* **8**, e2949.
- de Jong EC, Vieira PL, Kalinski P, Schuitmaker JHN, Tanaka Y, Wierenga EA, Yazdanbakhsh M and Kapsenberg ML (2002) Microbial compounds selectively induce Th1 cell-promoting or Th2 cell-promoting dendritic cells *in vitro* with diverse Th cell-polarizing signals. *The Journal of Immunology* **168**, 1704–1709.
- De Oliveira Fraga LA, Torrero MN, Tocheva AS, Mitre E and Davies SJ (2010) Induction of type 2 responses by schistosome worms during prepatent infection. *The Journal of Infectious Diseases* **201**, 464.
- Dewals BG, Marillier RG, Hoving JC, Leeto M, Schwegmann A and Brombacher F (2010) IL-4R α -independent expression of mannose receptor and Ym1 by macrophages depends on their IL-10 responsiveness. *PLoS Neglected Tropical Diseases* **4**, e689.
- Doenhoff MJ (1998) A vaccine for schistosomiasis: alternative approaches. *Parasitology Today* **14**, 105–109.
- Doenhoff MJ, Butterworth AE, Hayes RJ, Sturrock RF, Ouma JH, Koeh D, Prentice M and Bain J (1993) Seroepidemiology and serodiagnosis of schistosomiasis in Kenya using crude and purified egg antigens of *Schistosoma mansoni* in ELISA. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **87**, 42–48.
- Du P, Ma Q, Zhu Z, Li G, Wang Y, Li QQ, Chen YF, Shang ZZ, Zhang J and Zhao L (2016) Mechanism of corilagin interference with IL-13/STAT6 signaling pathways in hepatic alternative activation macrophages in schistosomiasis-induced liver fibrosis in mouse model. *European Journal of Pharmacology* **793**, 119–126.
- Dunne DW and Cooke A (2005) A worm's eye view of the immune system: consequences for evolution of human autoimmune disease. *Nature Reviews Immunology* **5**, 420–426.
- Dunne DW, Jones FM and Doenhoff MJ (1991) The purification, characterization, serological activity and hepatotoxic properties of two cationic glycoproteins (alpha 1 and omega 1) from *Schistosoma mansoni* eggs. *Parasitology* **103**(Pt 2), 225–236.
- Egesa M, Lubyayi L, Tukahebwa EM, Bagaya BS, Chalmers IW, Wilson S, Hokke CH, Hoffmann KF, Dunne DW, Yazdanbakhsh M, Labuda LA and Cose S (2018) *Schistosoma mansoni* schistosomula antigens induce Th1/pro-inflammatory cytokine responses. *Parasite Immunology* **40**, e12592.
- El-Faham MH, Wheatcroft-Francklow KJ, Price HP, Sayers JR and Doenhoff MJ (2017) *Schistosoma mansoni* cercarial elastase (SmCE): differences in immunogenic properties of native and recombinant forms. *Parasitology* **144**, 1356–1364.
- El-Zayat SR, Sibaii H and Mannaa FA (2019) Toll-like receptors activation, signaling, and targeting: an overview. *Bulletin of the National Research Centre* **43**, 1–12.
- Enderlin Vaz Da Silva Z, Lehr HA and Velin D (2014) *In vitro* and *in vivo* repair activities of undifferentiated and classically and alternatively activated macrophages. *Pathobiology: Journal of Immunopathology, Molecular and Cellular Biology* **81**, 86–93.
- Everts B, Perona-Wright G, Smits HH, Hokke CH, van der Ham AJ, Fitzsimmons CM, Doenhoff MJ, van der Bosch J, Mohrs K, Haas H, Mohrs M, Yazdanbakhsh M and Schramm G (2009) Omega-1, a glycoprotein secreted by *Schistosoma mansoni* eggs, drives Th2 responses. *The Journal of Experimental Medicine* **206**, 1673–1680.
- Fang Y, Wu C, Chen Q, Wu J, Yang Y, Guo X, Chen G and Wang Z (2015) SJE16.7 activates macrophages and promotes *Schistosoma japonicum* egg-induced granuloma development. *Acta Tropica* **149**, 49–58.
- Fei-Yue L, Hong-Zhuan T, Jie Z, Rui-Hong Z, Jin-Hua Z, Xin-Ting C and Guang-Hui R (2017) Analysis of characteristics of medical assistance to advanced schistosomiasis patients in Hunan province, 2015. *Zhongguo xue xi chong bing fang zhi za zhi = Chinese Journal of Schistosomiasis Control* **29**, 281–285.
- Ferrante CJ, Pinhal-Enfield G, Elson G, Cronstein BN, Hasko G, Outram S and Leibovich SJ (2013) The adenosine-dependent angiogenic switch of macrophages to an M2-like phenotype is independent of interleukin-4 receptor alpha (IL4R α) signaling. *Inflammation* **36**, 921.
- Fraternali A, Brundu S and Magnani M (2015) Polarization and repolarization of macrophages. *Journal of Clinical and Cellular Immunology* **6**, 1000319.
- Freitas MS, Oliveira AF, Da Silva TA, Fernandes FF, Gonçalves RA, Almeida F and Roque-Barreira MC (2016) Paracoccin induces M1 polarization of macrophages via interaction with TLR4. *Frontiers in Microbiology* **7**, 1003.
- Gao Y, Chen L, Hou M, Chen Y, Ji M, Wu H and Wu G (2013) TLR2 directing PD-L2 expression inhibit T cells response in *Schistosoma japonicum* infection. *PLoS ONE* **8**, e82480.
- Gao S, Zhou J, Liu N, Wang L, Gao Q, Wu Y, Zhao Q, Liu P, Wang S, Liu Y, Guo N, Shen Y, Wu Y and Yuan Z (2015) Curcumin induces M2 macrophage polarization by secretion IL-4 and/or IL-13. *Journal of Molecular and Cellular Cardiology* **85**, 131–139.
- Ghosh S, Shang P, Yazdankhah M, Bhutto I, Hose S, Montezuma SR, Luo T, Chattopadhyay S, Qian J, Luty GA, Ferrington DA, Zigler JS and Sinha D (2017) Activating the AKT2-nuclear factor- κ B-lipocalin-2 axis elicits an inflammatory response in age-related macular degeneration. *The Journal of Pathology* **241**, 583–588.
- Gobbi F, Tamarozzi F, Buonfrate D, van Lieshout L, Bisoffi Z and Bottieau E (2020) New insights on acute and chronic schistosomiasis: do we need a redefinition? *Trends in Parasitology* **36**, 660–667.
- Goh F, Irvine KM, Lovelace E, Donnelly S, Jones MK, Brion K, Hume DA, Kotze AC, Dalton JP, Ingham A and Sweet MJ (2009) Selective induction of the notch ligand Jagged-1 in macrophages by soluble egg antigen from *Schistosoma mansoni* involves ERK signalling. *Immunology* **127**, 326.
- Gong M, Zhuo X and Ma A (2017) STAT6 upregulation promotes M2 macrophage polarization to suppress atherosclerosis. *Medical Science Monitor Basic Research* **23**, 240–249.
- Gong W, Huang F, Sun L, Yu A, Zhang X, Xu Y, Shen Y and Cao J (2018) Toll-like receptor-2 regulates macrophage polarization induced by excretory-secretory antigens from *Schistosoma japonicum* eggs and promotes liver pathology in murine schistosomiasis. *PLoS Neglected Tropical Diseases* **12**, e0007000.
- Gordon S (2003) Alternative activation of macrophages. *Nature Reviews Immunology* **3**, 23–35.
- Gordon S and Martinez-Pomares L (2017) Physiological roles of macrophages. *Pflugers Archiv: European Journal of Physiology* **469**, 365–374.
- Grenfell RFQ, Martins WH, Silva-Moraes V, Barata SVB, Ribeiro EG, Oliveira E and Coelho PMZ (2012) Antigens of worms and eggs showed

- a differentiated detection of specific IgG according to the time of *Schistosoma mansoni* infection in mice. *Revista da Sociedade Brasileira de Medicina Tropical* 45, 505–509.
- Grieco S, Sulekova LF, Nardelli S, Riggio O, Venditti M and Taliani G** (2016) Portal hypertension related to schistosomiasis treated with a transjugular intrahepatic portosystemic shunt. *Journal of Clinical Gastroenterology* 50, 608–610.
- Gryseels B, Polman K, Clerinx J and Kestens L** (2006) Human schistosomiasis. *The Lancet* 368, 1106–1118.
- Grzych JM, Pearce E, Cheever A, Caulada ZA, Caspar P, Heiny S, Lewis F and Sher A** (1991) Egg deposition is the major stimulus for the production of Th2 cytokines in murine *Schistosomiasis mansoni*. *The Journal of Immunology* 146, 1322–13227.
- Hai Y, Edwards JE, Van Zandt MC, Hoffmann KF and Christianson DW** (2014) Crystal structure of *Schistosoma mansoni* arginase, a potential drug target for the treatment of schistosomiasis. *Biochemistry* 53, 4671–4684.
- Hambrook JR and Hanington PC** (2021) Immune evasion strategies of schistosomes. *Frontiers in Immunology* 11, 624178.
- Hams E, Avielle G and Fallon PG** (2013) The schistosoma granuloma: friend or foe? *Frontiers in Immunology* 4, 89.
- Harizi H, Juzan M, Pittard V, Moreau JF and Gualde N** (2002) Cyclooxygenase-2-induced prostaglandin e(2) enhances the production of endogenous IL-10, which down-regulates dendritic cell functions. *The Journal of Immunology* 168, 2255–2263.
- He YX, Yu QF, Yu P, Mao CS and Hu YQ** (1990) Penetration of *Schistosoma japonicum* cercaria into host skin. *Chinese Medical Journal* 103, 34–44.
- He YX, Salafsky B and Ramaswamy K** (2005) Comparison of skin invasion among three major species of *Schistosoma*. *Trends in Parasitology* 21, 201–203.
- He B, Cai G, Ni Y, Li Y, Zong H and He L** (2011) Characterization and expression of a novel cystatin gene from *Schistosoma japonicum*. *Molecular and Cellular Probes* 25, 186–193.
- He Y, Gao Y, Zhang Q, Zhou G, Cao F and Yao S** (2020) IL-4 switches microglia/macrophage M1/M2 polarization and alleviates neurological damage by modulating the JAK1/STAT6 pathway following ICH. *Neuroscience* 437, 161–171.
- Hervé M, Angeli V, Pinzar E, Wintjens R, Faveeuw C, Narumiya S, Capron A, Urade Y, Capron M, Riveau G and Trottein F** (2003) Pivotal roles of the parasite PGD2 synthase and of the host D prostanoid receptor 1 in schistosome immune evasion. *European Journal of Immunology* 33, 2764–2772.
- Hesse M, Modolell M, La Flamme AC, Schito M, Fuentes JM, Cheever AW, Pearce EJ and Wynn TA** (2001) Differential regulation of nitric oxide synthase-2 and arginase-1 by type 1/type 2 cytokines *in vivo*: granulomatous pathology is shaped by the pattern of L-arginine metabolism. *The Journal of Immunology* 167, 6533–6544.
- Hesse M, Piccirillo CA, Belkaid Y, Pruffer J, Mentink-Kane M, Leusink M, Cheever AW, Shevach EM and Wynn TA** (2004) The pathogenesis of schistosomiasis is controlled by cooperating IL-10-producing innate effector and regulatory T cells. *The Journal of Immunology* 172, 3157–3166.
- Ho CH, Cheng CH, Huang TW, Peng SY, Lee KM and Cheng PC** (2022) Switched phenotypes of macrophages during the different stages of *Schistosoma japonicum* infection influenced the subsequent trends of immune responses. *Journal of Microbiology, Immunology and Infection* 55, 503–526.
- Hoffmann KF, Cheever AW and Wynn TA** (2000) IL-10 and the dangers of immune polarization: excessive type 1 and type 2 cytokine responses induce distinct forms of lethal immunopathology in murine schistosomiasis. *The Journal of Immunology* 164, 6406–6416.
- Hogg KG, Kumkate S, Anderson S and Mountford AP** (2003a) Interleukin-12 p40 secretion by cutaneous CD11c+ and F4/80+ cells is a major feature of the innate immune response in mice that develop Th1-mediated protective immunity to *Schistosoma mansoni*. *Infection and Immunity* 71, 3563–3571.
- Hogg KG, Kumkate S and Mountford AP** (2003b) IL-10 regulates early IL-12-mediated immune responses induced by the radiation-attenuated schistosome vaccine. *International Immunology* 15, 1451–1459.
- Houlder EL, Costain AH, Cook PC and MacDonald AS** (2021) Schistosomes in the lung: immunobiology and opportunity. *Frontiers in Immunology* 12, 1330.
- Hu S, Wu Z, Yang L and Fung MC** (2009) Molecular cloning and expression of a functional anti-inflammatory protein, Sj16, of *Schistosoma japonicum*. *International Journal for Parasitology* 39, 191–200.
- Hussell T and Bell TJ** (2014) Alveolar macrophages: plasticity in a tissue-specific context. *Nature Reviews. Immunology* 14, 81–93.
- Ingram RJ, Bartlett A, Brown MB, Marriott C and Whitfield PJ** (2003) Penetration of human skin by the cercariae of *Schistosoma mansoni*: an investigation of the effect of multiple cercarial applications. *Journal of Helminthology* 77, 27–31.
- Ingram JR, Rafi SB, Eroy-Reveles AA, Ray M, Lambeth L, Hsieh I, Ruelas D, Lim KC, Sakanari J, Craik CS, Jacobson MP and McKerrow JH** (2012) Investigation of the proteolytic functions of an expanded cercarial elastase gene family in *Schistosoma mansoni*. *PLoS Neglected Tropical Diseases* 6, e1589.
- Ismail HAHA, Hong ST, Babiker ATEB, Hassan RMAE, Sulaiman MAZ, Jeong HG, Kong WH, Lee SH, Cho HI, Nam HS, Oh CH and Lee YH** (2014) Prevalence, risk factors, and clinical manifestations of schistosomiasis among school children in the White Nile River basin, Sudan. *Parasites & Vectors* 7, 478.
- James SL and Glaven J** (1989) Macrophage cytotoxicity against schistosome of *Schistosoma mansoni* involves arginine-dependent production of reactive nitrogen intermediates. *The Journal of Immunology* 143, 4208–4212.
- James SL, Cheever AW, Caspar P and Wynn TA** (1998) Inducible nitric oxide synthase-deficient mice develop enhanced type 1 cytokine-associated cellular and humoral immune responses after vaccination with attenuated *Schistosoma mansoni* cercariae but display partially reduced resistance. *Infection and Immunity* 66, 3510.
- Jang-Lee J, Curwen RS, Ashton PD, Tissot B, Mathieson W, Panico M, Dell A, Wilson RA and Haslam SM** (2007) Glycomics analysis of *Schistosoma mansoni* egg and cercarial secretions. *Molecular & Cellular Proteomics* 6, 1485–1499.
- Jang E, Lee S, Kim JH, Kim JH, Seo JW, Lee WH, Mori K, Nakao K and Suk K** (2013) Secreted protein lipocalin-2 promotes microglial M1 polarization. *FASEB Journal* 27, 1176–1190.
- Janssen L, Silva Santos GL, Muller HS, Vieira ARA, De Campos TA and De Paulo Martins V** (2016) Schistosome-derived molecules as modulating actors of the immune system and promising candidates to treat auto-immune and inflammatory diseases. *Journal of Immunology Research* 2016, 5267485.
- Jenkins SJ, Hewitson JP, Ferret-Bernard S and Mountford AP** (2005a) Schistosome larvae stimulate macrophage cytokine production through TLR4-dependent and -independent pathways. *International Immunology* 17, 1409–1418.
- Jenkins SJ, Hewitson JP, Jenkins GR and Mountford AP** (2005b) Modulation of the host's immune response by schistosome larvae. *Parasite Immunology* 27, 385–393.
- Jenkins SJ, Ruckerl D, Cook PC, Jones LH, Finkelman FD, Van Rooijen N, MacDonald AS and Allen JE** (2011) Local macrophage proliferation, rather than recruitment from the blood, is a signature of Th2 inflammation. *Science (New York, N.Y.)* 332, 1284–1288.
- Jetten N, Verbruggen S, Gijbels MJ, Post MJ, De Winther MPJ and Donners MMPC** (2014) Anti-inflammatory M2, but not pro-inflammatory M1 macrophages promote angiogenesis *in vivo*. *Angiogenesis* 17, 109–118.
- Jin L, Yuan F, Chen C, Wu J, Gong R, Yuan G, Zeng H, Pei J and Chen T** (2019) Degradation products of polydopamine restrained inflammatory response of LPS-stimulated macrophages through mediation TLR-4/MyD88 dependent signaling pathways by antioxidant. *Inflammation* 42, 658–671.
- Kaisar MMM, Ritter M, del Fresno C, Jónasdóttir HS, van der Ham AJ, Pelgrom LR, Schramm G, Layland LE, Sancho D, Prazeres da Costa C, Giera M, Yazdanbakhsh M and Everts B** (2018) Dectin-1/2-induced auto-crine PGE2 signaling licenses dendritic cells to prime Th2 responses. *PLoS Biology* 16, e2005504.
- Kamdem SD, Moyou-Somo R, Brombacher F and Nono JK** (2018) Host regulators of liver fibrosis during human schistosomiasis. *Frontiers in Immunology* 9, 2781.
- Ke XD, Shen S, Song LJ, Yu CX, Kikuchi M, Hirayama K, Gao H, Wang J, Yin X, Yao Y, Liu Q and Zhou W** (2017) Characterization of *Schistosoma japonicum* CP1412 protein as a novel member of the ribonuclease T2 molecule family with immune regulatory function. *Parasites & Vectors* 10, 1–19.
- Khammo N, Bartlett A, Clothier RH and Whitfield PJ** (2002) The attachment of *Schistosoma mansoni* cercariae to human skin cells. *Parasitology* 124, 25–30.
- Kloc M, Ghobrial RM, Wosik J, Lewicka A, Lewicki S and Kubiak JZ** (2019) Macrophage functions in wound healing. *Journal of Tissue Engineering and Regenerative Medicine* 13, 99–109.

- Kloetzel K and Lewert RM (1966) Pigment formation in *Schistosoma mansoni* infections in the white mouse. *The American Journal of Tropical Medicine and Hygiene* 15, 28–31.
- Knuhr K, Langhans K, Nyenhuis S, Viertmann K, Overgaard Kildemoes AM, Doenhoff MJ, Haas H and Schramm G (2018) *Schistosoma mansoni* egg-released IPSE/alpha-1 dampens inflammatory cytokine responses via basophil interleukin (IL)-4 and IL-13. *Frontiers in Immunology* 9, 2293.
- Kumkate S, Jenkins GR, Paveley RA, Hogg KG and Mountford AP (2007) CD207+ Langerhans cells constitute a minor population of skin-derived antigen-presenting cells in the draining lymph node following exposure to *Schistosoma mansoni*. *International Journal for Parasitology* 37, 209–220.
- Lambertucci JR (2010) Acute *Schistosomiasis mansoni*: revisited and reconsidered. *Memórias do Instituto Oswaldo Cruz* 105, 422–435.
- Langenberg MCC, Hoogerwerf MA, Janse JJ, Van Lieshout L, Corstjens PLAM, Roestenberg M, Van Dam GJ, Van Diepen A, De Dood CJ, Feijt C, Ganesh MS, Gerritsma H, Hardeman G, Hokke CH, Koopman JPR, Kos-Van Oosterhoud J, Kruize YCM, Meij P, Ozir-Fazalalikhhan A, Van Schuijlenburg R, Smits HH, Verbeek-Menken PH, Visser LG, De Vries JJC, Winkel BMF and Yazdanbakhsh M (2019) Katayama syndrome without *Schistosoma mansoni* eggs. *Annals of Internal Medicine* 170, 732–733.
- Lechner A, Bohnacker S and Esser-von Bieren J (2021) Macrophage regulation & function in helminth infection. *Seminars in Immunology* 53, 101526.
- Lenzi HL, Kimmel E, Schechtman H, Pelajo-Machado M, Romanha WS, Pacheco RG, Mariano M and Lenzi JÁ (1998) Histoarchitecture of schistosomal granuloma development and involution: morphogenetic and biomechanical approaches. *Memórias do Instituto Oswaldo Cruz* 93, 141–151.
- Leontovych A, Ulrychová L, Horn M and Dvořák J (2020) Collection of excretory/secretory products from individual developmental stages of the blood fluke *Schistosoma mansoni*. *Methods in Molecular Biology* 2151, 55–63.
- Ley K (2017) M1 means kill; M2 means heal. *The Journal of Immunology* 199, 2191–2193.
- Li YQ, Chen YF, Dang YP, Wang Y, Shang ZZ, Ma Q, Wang YJ, Zhang J, Luo L, Li QQ and Zhao L (2017) Corilagin counteracts IL-13R α 1 signaling pathway in macrophages to mitigate schistosome egg-induced hepatic fibrosis. *Frontiers in Cellular and Infection Microbiology* 7, 443.
- Liu Y, Munker S, Müllenbach R and Weng HL (2012) IL-13 signaling in liver fibrogenesis. *Frontiers in Immunology* 3, 116.
- Liu YC, Zou XB, Chai YF and Yao YM (2014) Macrophage polarization in inflammatory diseases. *International Journal of Biological Sciences* 10, 520–529.
- Liu M, Ju C, Du XF, Shen HM, Wang JP, Li J, Zhang XM, Feng Z and Hu W (2015) Proteomic analysis on cercariae and schistosomula in reference to potential proteases involved in host invasion of *Schistosoma japonicum* larvae. *Journal of Proteome Research* 14, 4623–4634.
- Liu CP, Zhang X, Tan QL, Xu WX, Zhou CY, Luo M, Li X, Huang RY and Zeng X (2017) NF- κ B pathways are involved in M1 polarization of RAW 264.7 macrophage by polyporus polysaccharide in the tumor microenvironment. *PLoS ONE* 12, e0188317.
- Liu J, Zhu L, Wang J, Qiu L, Chen Y, Davis RE and Cheng G (2019) *Schistosoma japonicum* extracellular vesicle miRNA cargo regulates host macrophage functions facilitating parasitism. *PLoS Pathogens* 15, e1007817.
- Locati M, Curtale G and Mantovani A (2020) Diversity, mechanisms, and significance of macrophage plasticity. *Annual Review of Pathology* 15, 123–147.
- Lopes DM, Oliveira SC, Page B, Carvalho LP, Carvalho EM and Cardoso LS (2019) *Schistosoma mansoni* rSm29 antigen induces a regulatory phenotype on dendritic cells and lymphocytes from patients with cutaneous leishmaniasis. *Frontiers in Immunology* 9, 3122.
- LoVerde PT (2019) Schistosomiasis. *Advances in Experimental Medicine and Biology* 1154, 45–70.
- Lu Y, Li X, Liu S, Zhang Y and Zhang D (2018) Toll-like receptors and inflammatory bowel disease. *Frontiers in Immunology* 9, 72.
- Lundy SK and Lukacs NW (2013) Chronic schistosome infection leads to modulation of granuloma formation and systemic immune suppression. *Frontiers in Immunology* 4, 39.
- MacDonald AS, Straw AD, Dalton NM and Pearce EJ (2002) Cutting edge: Th2 response induction by dendritic cells: a role for CD40. *The Journal of Immunology* 168, 537–540.
- Martinez FO and Gordon S (2014) The M1 and M2 paradigm of macrophage activation: time for reassessment. *F1000Prime Reports* 6, 13.
- Masamba P and Kappo AP (2021) Immunological and biochemical interplay between cytokines, oxidative stress and schistosomiasis. *International Journal of Molecular Sciences* 22, 7216.
- Masi B, Perles-Barbacaru TA, Bernard M and Viola A (2020) Clinical and preclinical imaging of hepatosplenic schistosomiasis. *Trends in Parasitology* 36, 206–226.
- McManus DP, Dunne DW, Sacko M, Utzinger J, Vennervald BJ and Zhou XN (2018) Schistosomiasis. *Nature Reviews. Disease Primers* 4, 13.
- Meevissen MHJ, Wuhrer M, Doenhoff MJ, Schramm G, Haas H, Deelder AM and Hokke CH (2010) Structural characterization of glycans on omega-1, a major *Schistosoma mansoni* egg glycoprotein that drives Th2 responses. *Journal of Proteome Research* 9, 2630–2642.
- Mehta A and Baltimore D (2016) MicroRNAs as regulatory elements in immune system logic. *Nature Reviews. Immunology* 16, 279–294.
- Menson EN and Wilson RA (1990) Lung-phase immunity to *Schistosoma mansoni*: definition of alveolar macrophage phenotypes after vaccination and challenge of mice. *Parasite Immunology* 12, 353–366.
- Meyer NH, Mayerhofer H, Tripsianes K, Blindow S, Barths D, Mewes A, Weimar T, Köhli T, Bade S, Madl T, Frey A, Haas H, Mueller-Dieckmann J, Sattler M and Schramm G (2015) A crystallin fold in the interleukin-4-inducing principle of *Schistosoma mansoni* eggs (IPSE/ α -1) mediates IgE binding for antigen-independent basophil activation. *The Journal of Biological Chemistry* 290, 22111–22126.
- Miller P and Wilson RA (1978) Migration of the schistosomula of *Schistosoma mansoni* from skin to lungs. *Parasitology* 77, 281–302.
- Mills CD (2015) Anatomy of a discovery: M1 and M2 macrophages. *Frontiers in Immunology* 6, 212.
- Molehin AJ (2020) Current understanding of immunity against schistosomiasis: impact on vaccine and drug development. *Research and Reports in Tropical Medicine* 11, 119–128.
- Montagner S, Dehó L and Monticelli S (2014) MicroRNAs in hematopoietic development. *BMC Immunology* 15, 14.
- Mouser EEIM, Pollakis G, Smits HH, Thomas J, Yazdanbakhsh M, De Jong EC and Paxton WA (2019) *Schistosoma mansoni* soluble egg antigen (SEA) and recombinant omega-1 modulate induced CD4+ T-lymphocyte responses and HIV-1 infection *in vitro*. *PLoS Pathogens* 15, e1007924.
- Murray PJ (2017) Macrophage polarization. *Annual Review of Physiology* 79, 541–566.
- Murray PJ, Allen JE, Biswas SK, Fisher EA, Gilroy DW, Goerdts S, Gordon S, Hamilton JA, Ivashkiv LB, Lawrence T, Locati M, Mantovani A, Martinez FO, Mege JL, Mosser DM, Natoli G, Saeij JP, Schultze JL, Shirey KA, Sica A, Suttles J, Udalova I, van Ginderachter, Vogel SN and Wynn TA (2014) Macrophage activation and polarization: nomenclature and experimental guidelines. *Immunity* 41, 14–20.
- Mutengo MM, Mduluzi T, Kelly P, Mwansa JCL, Kwenda G, Musonda P and Chipeta J (2018) Low IL-6, IL-10, and TNF- α and high IL-13 cytokine levels are associated with severe hepatic fibrosis in *Schistosoma mansoni* chronically exposed individuals. *Journal of Parasitology Research* 73, 349–360.
- Nathan C and Cunningham-Bussell A (2013) Beyond oxidative stress: an immunologist's guide to reactive oxygen species. *Nature Reviews. Immunology* 13, 349–361.
- Nation CS, Da'dara AA, Marchant JK and Skelly PJ (2020) Schistosome migration in the definitive host. *PLoS Neglected Tropical Diseases* 14, 1–12.
- Nelwan ML (2019) Schistosomiasis: life cycle, diagnosis, and control. *Current Therapeutic Research, Clinical and Experimental* 91, 5–9.
- Neves LX, Sanson AL, Wilson RA and Castro-Borges W (2015) What's in SWAP? Abundance of the principal constituents in a soluble extract of *Schistosoma mansoni* revealed by shotgun proteomics. *Parasites & Vectors* 8, 1–9.
- Nowacki FC, Swain MT, Klychnikov OI, Niazi U, Ivens A, Quintana JF, Hensbergen PJ, Hokke CH, Buck AH and Hoffmann KF (2015) Protein and small non-coding RNA-enriched extracellular vesicles are released by the pathogenic blood fluke *Schistosoma mansoni*. *Journal of Extracellular Vesicles* 4, 28665.
- Oliveira MF, D'Avila JCP, Torres CR, Oliveira PL, Tempone AJ, Rumjanek FD, Braga CMS, Silva JR, Dansa-Petretski M, Oliveira MA, De Souza W and Ferreira ST (2000) Haemozoin in *Schistosoma mansoni*. *Molecular and Biochemical Parasitology* 111, 217–221.
- Orecchioni M, Ghosheh Y, Pramod AB and Ley K (2019) Macrophage polarization: different gene signatures in M1(LPS+) vs classically and M2(LPS-) vs alternatively activated macrophages. *Frontiers in Immunology* 10, 1084.
- Oswald IP, Wynn TA, Sher A and James SL (1994) NO as an effector molecule of parasite killing: modulation of its synthesis by cytokines.

- Comparative Biochemistry and Physiology Part C: Pharmacology, Toxicology and Endocrinology* **108**, 11–18.
- Oyesola OO, Shanahan MT, Kanke M, Mooney BM, Webb LM, Smita S, Matheson MK, Campioli P, Pham D, Früh SP, McGinty JW, Churchill MJ, Cahoon JL, Sundaravaradan P, Flitter BA, Mouli K, Nadjombati MS, Kamynina E, Peng SA, Cubitt RL, Gronert K, Lord JD, Rauch I, von Moltke J, Sethupathy P and Tait Wojno ED (2021) PGD2 and CRTH2 counteract type 2 cytokine-elicited intestinal epithelial responses during helminth infection. *The Journal of Experimental Medicine* **218**, e20202178.
- Paik YH, Kim J, Aoyama T, De Minicis S, Bataller R and Brenner DA (2014) Role of NADPH oxidases in liver fibrosis. *Antioxidants & Redox Signaling* **20**, 2854.
- Parisi L, Gini E, Baci D, Tremolati M, Fanuli M, Bassani B, Farronato G, Bruno A and Mortara L (2018) Macrophage polarization in chronic inflammatory diseases: killers or builders? *Journal of Immunology Research* **2018**, 8917804.
- Paveley RA, Aynsley SA, Cook PC, Turner JD and Mountford AP (2009) Fluorescent imaging of antigen released by a skin-invading helminth reveals differential uptake and activation profiles by antigen presenting cells. *PLoS Neglected Tropical Diseases* **3**, e528.
- Pearce EJ and MacDonald AS (2002) The immunobiology of schistosomiasis. *Nature Reviews Immunology* **2**, 499–511.
- Pearce EJ, Caspar P, Grzych JM, Lewis FA and Sher A (1991) Downregulation of Th1 cytokine production accompanies induction of Th2 responses by a parasitic helminth, *Schistosoma mansoni*. *The Journal of Experimental Medicine* **173**, 159–166.
- Pearce EJ, Kane CM, Sun J, Taylor JJ, McKee AS and Cervi L (2004) Th2 response polarization during infection with the helminth parasite *Schistosoma mansoni*. *Immunological Reviews* **201**, 117–126.
- Perona-Wright G, Jenkins SJ and MacDonald AS (2006) Dendritic cell activation and function in response to *Schistosoma mansoni*. *International Journal for Parasitology* **36**, 711–721.
- Pesce JT, Ramalingam TR, Wilson MS, Mentink-Kane MM, Thompson RW, Cheever AW, Urban JF, Wynn TA and Mansfield JM (2009) Retnla (relmalph/fizz1) suppresses helminth-induced Th2-type immunity. *PLoS Pathogens* **5**, e1000393.
- Piipponen M, Li D and Landén NX (2020) The immune functions of keratinocytes in skin wound healing. *International Journal of Molecular Sciences* **21**, 1–26.
- Rahman K, Vengrenyuk Y, Ramsey SA, Vila NR, Girgis NM, Liu J, Gusarova V, Gromada J, Weinstock A, Moore KJ, Loke P and Fisher EA (2017) Inflammatory Ly6Chi monocytes and their conversion to M2 macrophages drive atherosclerosis regression. *The Journal of Clinical Investigation* **127**, 2904–2915.
- Ramaswamy K, Kumar P and He YX (2000) A role for parasite-induced PGE2 in IL-10-mediated host immunoregulation by skin stage schistosomula of *Schistosoma mansoni*. *The Journal of Immunology* **165**, 4567–4574.
- Raposo G and Stoorvogel W (2013) Extracellular vesicles: exosomes, microvesicles, and friends. *The Journal of Cell Biology* **200**, 373–383.
- Rath M, Müller I, Kropf P, Closs EI and Munder M (2014) Metabolism via arginase or nitric oxide synthase: two competing arginine pathways in macrophages. *Frontiers in Immunology* **5**, 532.
- Ross AG, Vickers D, Olds GR, Shah SM and McManus DP (2007) Katayama syndrome. *The Lancet. Infectious Diseases* **7**, 218–224.
- Ross EA, Devitt A and Johnson JR (2021) Macrophages: the good, the bad, and the gluttony. *Frontiers in Immunology* **12**, 708186.
- Roupé KM, Nybo M, Sjöbring U, Alberius P, Schmidtchen A and Sørensen OE (2010) Injury is a major inducer of epidermal innate immune responses during wound healing. *The Journal of Investigative Dermatology* **130**, 1167–1177.
- Rückerl D and Cook PC (2019) Macrophages assemble! But do they need IL-4R during schistosomiasis? *European Journal of Immunology* **49**, 996–1000.
- Ruzicka T and Printz MP (1984) Arachidonic acid metabolism in skin: a review. *Reviews of Physiology, Biochemistry and Pharmacology* **100**, 121–160.
- Salter JP, Lim KC, Hansell E, Hsieh I and McKerrow JH (2000) Schistosome invasion of human skin and degradation of dermal elastin are mediated by a single serine protease. *Journal of Biological Chemistry* **275**, 38667–38673.
- Sanin DE and Mountford AP (2015) Sm16, a major component of *Schistosoma mansoni* cercarial excretory/secretory products, prevents macrophage classical activation and delays antigen processing. *Parasites & Vectors* **8**, 1–12.
- Schmall A, Al-Tamari HM, Herold S, Kampschulte M, Weigert A, Wietelmann A, Vipotnik N, Grimminger F, Seeger W, Pullamsetti SS and Savai R (2015) Macrophage and cancer cell cross-talk via CCR2 and CX3CR1 is a fundamental mechanism driving lung cancer. *American Journal of Respiratory and Critical Care Medicine* **191**, 437–447.
- Schramm G, Gronow A, Knobloch J, Wippersteg V, Grevelding CG, Galle J, Fuller H, Stanley RG, Chiodini PL, Haas H and Doenhoff MJ (2006) IPSE/alpha-1: a major immunogenic component secreted from *Schistosoma mansoni* eggs. *Molecular and Biochemical Parasitology* **147**, 9–19.
- Schramm G, Mohrs K, Wodrich M, Doenhoff MJ, Pearce EJ, Haas H and Mohrs M (2007) Cutting edge: IPSE/alpha-1, a glycoprotein from *Schistosoma mansoni* eggs, induces IgE-dependent, antigen-independent IL-4 production by murine basophils *in vivo*. *The Journal of Immunology* **178**, 6023–6027.
- Schwartz C and Fallon PG (2018) *Schistosoma* ‘eggs-iting’ the host: granuloma formation and egg excretion. *Frontiers in Immunology* **9**, 2492.
- Schwartz E, Rozenman J and Perelman M (2000) Pulmonary manifestations of early schistosome infection among nonimmune travelers. *The American Journal of Medicine* **109**, 718–722.
- Schwartz C, Oeser K, Prazeres da Costa C, Layland LE and Voehringer D (2014) T cell-derived IL-4/IL-13 protects mice against fatal *Schistosoma mansoni* infection independently of basophils. *The Journal of Immunology* **193**, 3590–3599.
- Shapouri-Moghaddam A, Mohammadian S, Vazini H, Taghadosi M, Esmaili SA, Mardani F, Seifi B, Mohammadi A, Afshari JT and Sahebkar A (2018) Macrophage plasticity, polarization, and function in health and disease. *Journal of Cellular Physiology* **233**, 6425–6440.
- Shen J, Wang L, Peng M, Liu Z, Zhang B, Zhou T, Sun X and Wu Z (2019) Recombinant Sj16 protein with novel activity alleviates hepatic granulomatous inflammation and fibrosis induced by *Schistosoma japonicum* associated with M2 macrophages in a mouse model. *Parasites & Vectors* **12**, 457.
- Shen H, Wang Z, Huang A, Zhu D, Sun P and Duan Y (2021) Lipocalin 2 is a regulator during macrophage polarization induced by soluble worm antigens. *Frontiers in Cellular and Infection Microbiology* **11**, 1.
- Shi Y, Luo P, Wang W, Horst K, Bläsius F, Relja B, Xu D, Hildebrand F and Greven J (2020) M1 but not M0 extracellular vesicles induce polarization of RAW264.7 macrophages via the TLR4-NFκB pathway *in vitro*. *Inflammation* **43**, 1611–1619.
- Shiels J, Cwiklinkiid K, Alvarado R, Thivierge K, Cotton S, Santana BG, Toid J, Donnelly S, Taggart CC, Weldonid S and Dalton JP (2020) *Schistosoma mansoni* immunomodulatory molecule Sm16/SPO-1/SmSLP is a member of the trematode-specific helminth defence molecules (HDMs). *PLoS Neglected Tropical Diseases* **14**, 1–25.
- Sica A and Mantovani A (2012) Macrophage plasticity and polarization: *in vivo* veritas. *The Journal of Clinical Investigation* **122**, 787–795.
- Sica A, Erreni M, Allavena P and Porta C (2015) Macrophage polarization in pathology. *Cellular and Molecular Life Sciences* **72**, 4111–4126.
- Smith H, Forman R, Mair I and Else KJ (2018) Interactions of helminths with macrophages: therapeutic potential for inflammatory intestinal disease. *Expert Review of Gastroenterology & Hepatology* **12**, 997–1006.
- Song LJ, Yin XR, Mu SS, Li JH, Gao H, Zhang Y, Dong PP, Mei CJ and Hua ZC (2020) The differential and dynamic progression of hepatic inflammation and immune responses during liver fibrosis induced by *Schistosoma japonicum* or carbon tetrachloride in mice. *Frontiers in Immunology* **11**, 2633.
- Sotillo J, Pearson M, Potriquet J, Becker L, Pickering D, Mulvenna J and Loukas A (2016) Extracellular vesicles secreted by *Schistosoma mansoni* contain protein vaccine candidates. *International Journal for Parasitology* **46**, 1–5.
- Starkey Lewis P, Campana L, Aleksieva N, Cartwright JA, Mackinnon A, O’Duibhir E, Kendall T, Vermeren M, Thomson A, Gadd V, Dwyer B, Aird R, Man TY, Rossi AG, Forrester L, Park BK and Forbes SJ (2020) Alternatively activated macrophages promote resolution of necrosis following acute liver injury. *Journal of Hepatology* **73**, 349–360.
- Steinmann P, Keiser J, Bos R, Tanner M and Utzinger J (2006) Schistosomiasis and water resources development: systematic review, meta-analysis, and estimates of people at risk. *Lancet Infectious Diseases* **6**, 411–425.
- Stingl P and Stingl T (2017) Schistosomiasis. *MMW Fortschritte der Medizin* **159**, 51–54.
- Suzuki Y, Shirai M, Asada K, Yasui H, Karayama M, Hozumi H, Furuhashi K, Enomoto N, Fujisawa T, Nakamura Y, Inui N, Shirai T, Hayakawa H and Suda T (2018) Macrophage mannose receptor, CD206, predict prognosis in patients with pulmonary tuberculosis. *Scientific Reports* **8**, 1–9.

- Takaki KK, Rinaldi G, Berriman M, Pagán AJ and Ramakrishnan L (2021) *Schistosoma mansoni* eggs modulate the timing of granuloma formation to promote transmission. *Cell Host & Microbe* **29**, 58.
- Tang Y, Shen Y, Hong Y, Zhang Z, Zhai Q, Fu Z, Li H, Lu K and Lin J (2021) miR-181a regulates the host immune response against *Schistosoma japonicum* infection through the TLR4 receptor pathway. *Parasites & Vectors* **14**, 548.
- Truscott M, Evans DA, Gunn M and Hoffmann KF (2013) *Schistosoma mansoni* hemozoin modulates alternative activation of macrophages via specific suppression of Retnla expression and secretion. *Infection and Immunity* **81**, 133–142.
- Tundup S, Srivastava L, Nagy T and Harn D (2014) CD14 influences host immune responses and alternative activation of macrophages during *Schistosoma mansoni* infection. *Infection and Immunity* **82**, 3240.
- Uribe-Querol E and Rosales C (2020) Phagocytosis: our current understanding of a universal biological process. *Frontiers in Immunology* **11**, 1066.
- van Liempt E, van Vliet SJ, Engering A, García Vallejo JJ, Bank CMC, Sanchez-Hernandez M, van Kooyk Y and van Die I (2007) *Schistosoma mansoni* soluble egg antigens are internalized by human dendritic cells through multiple C-type lectins and suppress TLR-induced dendritic cell activation. *Molecular Immunology* **44**, 2605–2615.
- Vannella KM and Wynn TA (2017) Mechanisms of organ injury and repair by macrophages. *Annual Review of Physiology* **79**, 593–617.
- Vannella KM, Barron L, Borthwick LA, Kindrachuk KN, Narasimhan PB, Hart KM, Thompson RW, White S, Cheever AW, Ramalingam TR and Wynn T (2014) Incomplete deletion of IL-4R α by LysMCre reveals distinct subsets of M2 macrophages controlling inflammation and fibrosis in chronic schistosomiasis. *PLoS Pathogens* **10**, e1004372.
- Verjee MA (2019) Schistosomiasis: still a cause of significant morbidity and mortality. *Research and Reports in Tropical Medicine* **10**, 153–163.
- Viola A, Munari F, Sánchez-Rodríguez R, Scolaro T and Castegna A (2019) The metabolic signature of macrophage responses. *Frontiers in Immunology* **10**, 1462.
- Wang L, Yang Z, Li Y, Yu F, Brindley PJ, McManus DP, Wei D, Han Z, Feng Z, Li Y and Hu W (2006) Reconstruction and *in silico* analysis of the MAPK signaling pathways in the human blood fluke, *Schistosoma japonicum*. *FEBS Letters* **580**, 3677–3686.
- Wang M, Altinoglu S, Takeda YS and Xu Q (2015) Integrating protein engineering and bioorthogonal click conjugation for extracellular vesicle modulation and intracellular delivery. *PLoS ONE* **10**, e0141860.
- Wang S, Ye Q, Zeng X and Qiao S (2019) Functions of macrophages in the maintenance of intestinal homeostasis. *Journal of Immunology Research* **2019**, 1512969.
- Wang L, Shang X, Qi X, Ba D, Lv J, Zhou X, Wang H, Shaxika N, Wang J and Ma X (2020) Clinical significance of M1/M2 macrophages and related cytokines in patients with spinal tuberculosis. *Disease Markers* **2020**, 2509454.
- Wei Y, Huang N, Chen S, Chen D, Li X, Xu J and Yang Z (2018) The diagnosis and treatment introspection of the first imported case of atypical cerebral schistosomiasis in Guangzhou city. *PLoS Neglected Tropical Diseases* **12**, e0006171.
- West HC and Bennett CL (2018) Redefining the role of Langerhans cells as immune regulators within the skin. *Frontiers in Immunology* **8**, 1941.
- Wheater PR and Wilson RA (1979) *Schistosoma mansoni*: a histological study of migration in the laboratory mouse. *Parasitology* **79**, 49–62.
- Whitfield PJ, Bartlett A, Brown MB and Marriott C (2003) Invasion by schistosome cercariae: studies with human skin explants. *Trends in Parasitology* **19**, 339–340.
- Wilson RA (1987) Cercariae to liver worms: development and migration in the mammalian host. In D Rollinson, AJG Simpson (eds), *The Biology of Schistosomes*. New York: From Genes to Latrines, pp. 115–146.
- Wilson RA (1998) Interferon gamma is a key cytokine in lung phase immunity to schistosomes but what is its precise role? *Brazilian Journal of Medical and Biological Research* **31**, 157–161.
- Wilson RA and Jones MK (2021) Fifty years of the schistosome tegument: discoveries, controversies, and outstanding questions. *International Journal for Parasitology* **51**, 1213–1232.
- Wilson MS, Mentink-Kane MM, Pesce JT, Ramalingam TR, Thompson R and Wynn TA (2007) Immunopathology of schistosomiasis. *Immunology and Cell Biology* **85**, 148.
- Wolde M, Laan LC, Medhin G, Gadissa E, Berhe N and Tsegaye A (2020) Human monocytes/macrophage inflammatory cytokine changes following *in vivo* and *in vitro* *Schistosoma mansoni* infection. *Journal of Inflammation Research* **13**, 35–43.
- World Health Organization (2020) Status of Schistosomiasis Endemic Countries: 2018. Available at https://apps.who.int/neglected_diseases/ntddata/sch/sch.html.
- Xu J, Zhang H, Chen L, Zhang D, Ji M, Wu H and Wu G (2014) *Schistosoma japonicum* infection induces macrophage polarization. *Journal of Biomedical Research* **28**, 299–308.
- Xu ZJ, Gu Y, Wang CZ, Jin Y, Wen XM, Ma JC, Tang LJ, Mao ZW, Qian J and Lin J (2019) The M2 macrophage marker CD206: a novel prognostic indicator for acute myeloid leukemia. *Oncoimmunology* **9**, 1683347.
- Xue L, Gyles SL, Wetley FR, Gazi L, Townsend E, Hunter MG and Pettipher R (2005) Prostaglandin D2 causes preferential induction of proinflammatory Th2 cytokine production through an action on chemoattractant receptor-like molecule expressed on Th2 cells. *The Journal of Immunology* **175**, 6531–6536.
- Yang Z and Ming XF (2014) Functions of arginase isoforms in macrophage inflammatory responses: impact on cardiovascular diseases and metabolic disorders. *Frontiers in Immunology* **5**, 533.
- Yang SH, Sharrocks AD and Whitmarsh AJ (2003) Transcriptional regulation by the MAP kinase signaling cascades. *Gene* **320**, 3–21.
- Yao Y, Xu XH and Jin L (2019) Macrophage polarization in physiological and pathological pregnancy. *Frontiers in Immunology* **10**, 792.
- Ye Z, Huang S, Zhang Y, Mei X, Zheng H, Li M, Chen J and Lu F (2020) Galectins, eosinophiles, and macrophages may contribute to *Schistosoma japonicum* egg-induced immunopathology in a mouse model. *Frontiers in Immunology* **11**, 146.
- Yin Z, Ma T, Lin Y, Lu X, Zhang C, Chen S and Jian Z (2018) IL-6/STAT3 pathway intermediates M1/M2 macrophage polarization during the development of hepatocellular carcinoma. *Journal of Cellular Biochemistry* **119**, 9419–9432.
- Yu Y, Wang J, Wang X, Gu P, Lei Z, Tang R, Wei C, Xu L, Wang C, Chen Y, Pu Y, Qi X, Yu B, Chen X, Zhu J, Li Y, Zhang Z, Zhou S and Su C (2021) Schistosome eggs stimulate reactive oxygen species production to enhance M2 macrophage differentiation and promote hepatic pathology in schistosomiasis. *PLoS Neglected Tropical Diseases* **15**, e0009696.
- Yunna C, Mengru H, Lei W and Weidong C (2020) Macrophage M1/M2 polarization. *European Journal of Pharmacology* **877**, 173090.
- Zhao P, Elks CM and Stephens JM (2014) The induction of lipocalin-2 protein expression *in vivo* and *in vitro*. *The Journal of Biological Chemistry* **289**, 5960–5969.
- Zheng S, Zhang P, Chen Y, Zheng S, Zheng L and Weng Z (2016) Inhibition of notch signaling attenuates schistosomiasis hepatic fibrosis *via* blocking macrophage M2 polarization. *PLoS ONE* **11**, e0166808.
- Zheng B, Zhang J, Chen H, Nie H, Miller H, Gong Q and Liu C (2020) T lymphocyte-mediated liver immunopathology of schistosomiasis. *Frontiers in Immunology* **11**, 61.
- Zhu J, Xu Z, Chen X, Zhou S, Zhang W, Chi Y, Li W, Song X, Liu F and Su C (2014) Parasitic antigens alter macrophage polarization during *Schistosoma japonicum* infection in mice. *Parasites & Vectors* **7**, 1–9.
- Zhu S, Wang S, Lin Y, Jiang P, Cui X, Wang X, Zhang Y and Pan W (2016) Release of extracellular vesicles containing small RNAs from the eggs of *Schistosoma japonicum*. *Parasites & Vectors* **9**, 1–9.
- Zussman RA, Bauman PM and Petruska JC (1970) The role of ingested hemoglobin in the nutrition of *Schistosoma mansoni*. *The Journal of Parasitology* **56**, 75–79.