

RESULTS OF A RADIO CONTINUUM SURVEY OF SPIRAL GALAXIES AT 10.55 GHz

S. NIKLAS AND R. WIELEBINSKI

*Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69,
D-53121 Bonn, Germany*

U. KLEIN

*Radioastronomisches Institut, Universität Bonn, Auf dem Hügel 71,
D-53121 Bonn, Germany*

AND

J. BRAINE

*IRAM, 300 Rue de la Piscine, F-38406 St. Martin d'Hères,
Franc*

The *Revised Shapley-Ames Catalog* (Sandage & Tamman, 1981) contains 335 spiral galaxies brighter than $B_T = 12$. A subsample of these galaxies with $\delta \geq -25^\circ$ and total flux densities at 1.49 GHz (Condon, 1987) above 10 mJy were observed at $\lambda 2.8$ cm using the 100-m radio telescope of the MPIfR Bonn. Depending on the expected flux density and extent of the source two observational methods were used: cross-scanning and mapping. In total, radio flux densities of 192 galaxies have been derived (Niklas *et al.*, 1995). Additionally, a literature search was made in order to get flux densities at other frequencies.

One can separate the thermal and non-thermal emission by fitting to the radio spectra of a galaxy an optically thin thermal component with a spectral index of -0.1 and a synchrotron component with a spectral index α_{nth} . The fit parameters are the thermal fraction $f_{\text{th}}^{1\text{GHz}}$ at 1 GHz and the non-thermal spectral index α_{nth} . The mean values of the derived distributions are: $\overline{f_{\text{th}}^{1\text{GHz}}} = 0.07 \pm 0.01$, $\sigma = 0.05$ and $\overline{\alpha_{\text{nth}}} = 0.85 \pm 0.02$, $\sigma = 0.12$. The thermal fraction seems to be independent of morphological type. The left diagram of Fig. 1 shows the distribution of α_{nth} . Sa and Sab galaxies and some irregular galaxies tend to have flatter spectra than Sb/Sc galaxies. A test of the separation of thermal and non-thermal emission was made. We have calculated the flux density of the thermal radio emission at the corre-




Figure 1. The left diagram shows the distribution of the derived α_{nth} . The different grey scales represents the morphological types. The right diagrams shows the plot of $\alpha_{\text{nth}}^{\text{H}\alpha}$ versus $\alpha_{\text{nth}}^{\text{radio}}$. The solid line corresponds to perfect agreement.

sponding frequencies using the $\text{H}\alpha$ data of Kennicutt & Kent (1983). These thermal fluxes were subtracted from the composite radio spectra. We fitted a power law to the residual spectrum in order to derive the non-thermal spectral index $\alpha_{\text{nth}}^{\text{H}\alpha}$. The right diagram of Fig. 1 shows $\alpha_{\text{nth}}^{\text{H}\alpha}$ versus $\alpha_{\text{nth}}^{\text{radio}}$. There exists a good correlation between the derived quantities. The shift towards flatter $\alpha_{\text{nth}}^{\text{H}\alpha}$ may be due to optical absorption in the host galaxies.

References

- Sandage, A., Tammann, G.A. 1981, *A Revised Shapley-Ames Catalog of Bright Galaxies* (Washington, D.C.: Carnegie Institution of Washington)(RSA)
 Condon, J. 1987, *ApJS*, **65**, 465
 Niklas, S., Klein, U., Braine, J., Wielebinski, R. 1995, *A&AS*, **113**
 Kennicutt, R.C., Kent, S.M. 1983, *ApJ*, **88**, 1094