

## SUBSETS CHARACTERIZING THE CLOSURE OF THE NUMERICAL RANGE

K. C. DAS, S. MAJUMDAR and BRAILEY SIMS

(Received 28 May 1984)

Communicated by J. H. Chabrowski

### Abstract

For an operator on a Hilbert space, points in the *closure* of its numerical range are characterized as either extreme, non-extreme boundary, or interior in terms of various associated sets of bounded sequences of vectors. These generalize similar results due to Embry, for points in the numerical range.

1980 *Mathematics subject classification (Amer. Math. Soc.):* 47 A 12

### 1. Introduction

Let  $T$  be an operator (that is, a bounded linear transformation) on a complex Hilbert space  $H$  with inner product  $\langle \cdot, \cdot \rangle$  and associated norm  $\|\cdot\|$ . It is well known that the numerical range

$$W(T) = \{ \langle Tx, x \rangle : \|x\| = 1, x \in H \}$$

is a convex subset of the complex plane. Denote the closure of  $W(T)$  by  $W(T)^-$ . Theorem 1 of M. R. Embry (1970) characterizes every point  $z$  of  $W(T)$  as either an extreme point, a non-extreme boundary point or an interior point in terms of the subset  $M_z(T)$  and its linear span, where

$$M_z(T) = \left\{ x \in H : \langle Tx, x \rangle - z\|x\|^2 = 0 \right\} \quad (z \in W(T)).$$

This theorem, though very interesting, does not characterize the unattained boundary points of the numerical range. In this note we attempt to fill this gap by

a generalization which can be applied to every point of  $W(T)^-$ . For any  $z \in W(T)^-$ , let

$$N_z(T) = \{(x_n) \in l_\infty(H) : \langle Tx_n, x_n \rangle - z\|x_n\|^2 \rightarrow 0\},$$

$$N'_z(T) = \{(x_n) \in l_\infty(H) : \langle Tx_n, x_n \rangle / \|x_n\|^2 \rightarrow z\},$$

$$N^L(T) = \bigcup_z \{N_z(T) : z \in L \cap W(T)^-\}$$

and

$$N_L(T) = \{(x_n) \in l_\infty(H) : \inf_{z \in L} |\langle Tx_n, x_n \rangle - z\|x_n\|^2| \rightarrow 0\}$$

where  $l_\infty(H)$  is the set of all bounded sequences of vectors from  $H$  and  $L$  is a line of support for  $W(T)^-$ . Let  $\gamma N_z(T)$  be the linear span of  $N_z(T)$ . Since  $N_z(T)$  is homogeneous,  $\gamma N_z(T) = N_z(T) + N_z(T)$ . It is readily seen that  $N_L(T)$  is a subspace (Majumdar and Sims (to appear)).

### 2. Basic lemmas

In order to establish our characterization for points of  $W(T)^-$  we need the following two lemmas. The first, stated without proof, is an easy corollary to Lemma 3 of Majumdar and Sims (to appear).

**LEMMA 1.** *If  $b$  is an extreme point of  $W(T)^-$  and  $L$  is a line of support for  $W(T)$  passing through  $b$ , then  $\lim \langle (T - b)x_n, y_n \rangle = 0$  and  $\lim \langle (T - b)y_n, x_n \rangle = 0$  for all  $(x_n) \in N_b(T)$  and  $(y_n) \in N_L(T)$ .*

**LEMMA 2.** *Let  $z$  be in the interior of a line segment lying in  $W(T)^-$  with end points  $a$  and  $b$ . Then  $N'_a(T) \subset \gamma N_z(T)$ .*

**PROOF.** Without loss of generality we may take  $a = 1, b = 0$  and  $(x_n) \in N'_1(T)$  to have  $\|x_n\| = 1$ . Let  $(y_n) \in N_0(T)$  be such that  $\|y_n\| = 1$  and  $\text{Re} \langle \text{Im } Tx_n, y_n \rangle = 0$ . For any bounded sequence  $(r_n)$ , let  $h_n = r_n x_n + y_n$ ; then we have  $\langle \text{Im } Th_n, h_n \rangle \rightarrow 0$ . We show the existence of two such distinct sequences  $(r_n)$  for which

$$(1) \quad \langle \text{Re } Th_n, h_n \rangle - z\|h_n\|^2 = 0$$

for all sufficiently large  $n$ . The equations in  $r_n$  given by (1) are equivalent to

$$r_n^2(1 - z + \epsilon_n) + 2r_n \text{Re} \langle (\text{Re } T - z)x_n, y_n \rangle + (\epsilon'_n - z) = 0$$

where  $\epsilon_n = \langle \text{Re } Tx_n, x_n \rangle - 1$  and  $\epsilon'_n = \langle \text{Re } Ty_n, y_n \rangle$ , both of which tend to zero. Thus the equations in (1) are of the form  $A_n r_n^2 + B_n r_n + C_n = 0$  where  $A_n, B_n, C_n$  are real numbers independent of  $r_n$ .

Let  $D_n = B_n^2 - 4A_nC_n$ , then

$$D_n = 4[\operatorname{Re}\langle \operatorname{Re} T - z, x_n, y_n \rangle]^2 + 4z(1 - z) + \delta_n$$

where  $\delta_n \rightarrow 0$ . Hence there are positive constants  $\alpha, \beta$  such that for all sufficiently large  $n$ ,  $\alpha \leq A_n$ ,  $D_n \leq \beta$  and  $|B_n| \leq \beta$ . This shows the existence of two distinct sequences solving (1) both of which are bounded and whose differences  $d_n = \sqrt{D_n}/A_n$  are eventually bounded away from zero. Thus we have for both these sequences that  $h_n \in N_z(T)$ . Subtraction and the fact that  $d_n$  is uniformly bounded away from zero gives  $(x_n) \in \gamma N_z(T)$ .

**REMARK.** A simplified version of the above argument applied to a pair of points  $a, b$  lying in a line segment in  $W(T)$  shows the existence of a real number  $r$  and a vector  $y$  such that  $a = \langle Tx, x \rangle$ ,  $b = \langle Ty, y \rangle$ ,  $\|x\| = \|y\| = 1$  and  $\langle T(rx + y), rx + y \rangle / \|rx + y\|^2 = ta + (1 - t)b$ ,  $0 < t < 1$ , yielding the convexity of  $W(T)$ . In contrast with the proof of convexity given by Halmos (1967), this argument gives two explicit values of  $r$ .

### 3. Characterization of $W(T)^-$

**THEOREM 3.** *Every element  $z$  of  $W(T)^-$  can be characterized as follows.*

- (i)  $z$  is an extreme point of  $W(T)^-$  if and only if  $N_z(T)$  is a subspace.
- (ii) If  $z$  is a nonextreme boundary point of  $W(T)^-$  and  $L$  the line of support for  $W(T)$  passing through  $z$ , then (a)  $\gamma N_z(T) = N_L(T) + N_z(T)$  and (b)  $N_L(T) = l_\infty(H)$  if and only if  $W(T)^- \subset L$ .
- (iii) If  $W(T)^-$  is not a straight line segment, then  $z$  is an interior point of  $W(T)^-$  if and only if  $N'_a(T) \subset \gamma N_z(T)$  for all  $z \in W(T)^-$ .

**PROOF.** (i) See Das and Craven (1983) and also Majumdar and Sims (to appear). Also note that the result  $N_z(T)$  is a subspace when  $z$  is an extreme point of  $W(T)^-$  can be deduced as a corollary to Lemma 1. Homogeneity being obvious, we prove linearity. Let  $(x_n^{(1)}), (x_n^{(2)}) \in N_z(T)$ . Thus  $(x_n^{(1)}), (x_n^{(2)}) \in N_L(T)$  where  $L$  is a line of support for  $W(T)$  passing through  $z$ . But  $N_L(T)$  is a subspace. So  $(x_n^{(1)} + x_n^{(2)}) \in N_L(T)$ . Now since  $(x_n^{(i)}) \in N_z(T)$ ,  $i = 1, 2$  and  $(x_n^{(1)} + x_n^{(2)}) \in N_L(T)$ , by Lemma 1 we have  $\lim \langle (T - z)x_n^{(1)}, x_n^{(1)} + x_n^{(2)} \rangle = 0$  for  $i = 1, 2$  and hence  $\lim \langle (T - z)(x_n^{(1)} + x_n^{(2)}), x_n^{(1)} + x_n^{(2)} \rangle = 0$  as required.

(ii) (a) We first show  $N_a(T) \subset \gamma N_z(T)$  for each  $a \in L \cap W(T)^-$ . Without loss of generality we may take  $L$  as the real axis and  $\operatorname{Im} W(T) \geq 0$ . Let  $(x_n) \in N_a(T)$  and  $(y_n) \in N_b(T)$ ,  $\|y_n\| = 1$  where  $b \in L$  is the extreme point of  $W(T)^-$  such that  $(a - z)/(z - b) \geq 0$ . Then  $(y_n)$  can be chosen so that  $\operatorname{Re}\langle y_n, x_n \rangle = 0$  and

Lemma 1 gives  $\operatorname{Re}\langle Ty_n, x_n \rangle \rightarrow 0$ . Also  $\operatorname{Im} W(T) \geq 0$  implies  $\operatorname{Im} Ty_n \rightarrow 0$ . Let  $r_n = [(a - z)/(z - b)]^{1/2} \|x_n\|$ . Then easy calculations show that with our assumptions  $\langle T(x_n \pm r_n y_n), x_n \pm r_n y_n \rangle - z \|x_n \pm r_n y_n\|^2 \rightarrow 0$ . That is  $(x_n \pm r_n y_n) \in N_z(T)$ . As in the proof of Lemma 2, adding these two sequences and using the homogeneity of  $N_z(T)$  we have  $(x_n) \in \gamma N_z(T)$ . Thus  $N_a(T) \subset \gamma N_z(T)$  for all  $a \in L \cap W(T)^-$  and so we have  $N^L(T) \subset \gamma N_z(T)$ . Since  $N_z(T) \subset N^L(T) \subset \gamma N_z(T)$ , by taking the vector sum of  $N_z(T)$  with each of these subsets we obtain  $\gamma N_z(T) = N^L(T) + N_z(T)$ .

(b) As before, if we take  $L$  as the real axis, we have  $N_L(T) = \{(x_n) \in l_\infty(H) : \operatorname{Im}\langle Tx_n, x_n \rangle \rightarrow 0\}$ . Now if  $W(T)^- \subset L$ ,  $(x_n) \in l_\infty(H)$  implies  $\operatorname{Im}\langle Tx_n, x_n \rangle = 0$  and so  $(x_n) \in N_L(T)$ . Hence  $N_L(T) = l_\infty(H)$ . Conversely if  $W(T)^-$  is not a subset of  $L$ , there exists  $(x_n) \in l_\infty(H)$ ,  $\|x_n\| = 1$  such that  $\operatorname{Im}\langle Tx_n, x_n \rangle$  does not tend to zero, or equivalently,  $(x_n) \notin N_L(T)$ . Hence  $N_L(T) \neq l_\infty(H)$ .

(iii) If  $z$  is an interior point of  $W(T)^-$ , by Lemma 2,  $N'_a(T) \subset \gamma N_z(T)$  whenever  $a \in W(T)^-$ . On the other hand, if  $z$  is a boundary point of  $W(T)^-$ , without loss of generality we may take  $L$ , the line of support for  $W(T)$  passing through  $z$ , as the real axis, in which case,  $N_L(T) = \{(x_n) \in l_\infty(H) : \operatorname{Im}\langle Tx_n, x_n \rangle \rightarrow 0\}$ . Thus  $\gamma N_z(T) \subset N_L(T)$  since  $N_L(T)$  is a subspace, but as  $W(T)^-$  does not lie in  $L$ , there exists an  $a \in W(T)$  such that  $\operatorname{Im} a \neq 0$ . Hence  $N'_a(T)$  is not a subset of  $\gamma N_z(T)$ .

## References

- K. C. Das and B. D. Craven (1983), 'Linearity and weak convergence on the boundary of numerical range', *J. Austral. Math. Soc. Ser. A* **35**, 221–226.  
 M. R. Embry (1970), 'The numerical range of an operator', *Pacific J. Math.* **32**, 647–650.  
 M. R. Embry (1975), 'Orthogonality and the numerical range,' *J. Math. Soc. Japan* **27**, 405–411.  
 P. R. Halmos (1967), *Hilbert space problem book* (Van Nostrand, New York).  
 S. Majumdar and B. Sims (to appear), 'Subspaces associated with boundary points of the numerical range', *J. Austral. Math. Soc. Ser. A*.  
 J. G. Stampfil (1966), 'Extreme points of the numerical range of a hyponormal operator', *Michigan Math. J.* **13**, 87–89.

K. C. Das  
 Indian Institute of Technology  
 Karagpur 721302  
 India

S. Majumdar and Brailey Sims  
 University of New England  
 Armidale, N.S.W. 2351  
 Australia