

## Instructions to authors

**Submission procedure.** Submission of a paper to *European Psychiatry* is understood to imply that the data contained therein has not previously been published (except in abstract form) or is being considered for publication elsewhere.

There are no submission or page charges.

Only submissions in English will be considered. One original plus three copies (one set of photographs must be original prints) should be submitted to one of the following *five receiving centres* according to geographical location:

Pr C BALLUS  
Unidad de Psiquiatría  
Facultad de Medicina  
Casanova, 143  
08036 Barcelona, Spain  
Fax: (34) 32 54 66 91

Pr M MAJ  
Università di Napoli  
Ist di Psichiatria  
Largo Madonna delle Grazie  
80138 Napoli Italy  
Fax: (39) 81 44 99 38

Pr JD GUELFI  
Clinique de la Faculté  
100, rue de la Santé  
75674 Paris Cedex 14 France  
Fax: (33) 1 45 65 89 43

Pr CB PULL  
Centre Hospitalier de Luxembourg  
4, rue Barblé  
1210 Luxembourg  
Fax: (352) 45 87 62

Pr H HEIMANN  
Goethestrasse 23  
7400 Tübingen Germany  
Fax: (49) 70 71 29 41 41

Pr M ACKENHEIL  
Psychiatrische Klinik  
und Poliklinik  
der Universität München  
Nussbaumstr 7  
8000 München 2 Germany  
Fax: (49) 89 51 60 47 41

Pr RM MURRAY  
Institute of Psychiatry  
De Crespigny Park  
London SE5 8AF UK  
Fax: (44) 71 70 35 796

Dr P BECH  
Dept of Psychiatry  
Frederiksborg General Hospital  
DK-3400 Hillerød Denmark  
Fax: (45) 48 24 20 16

### Deputy Editors

Dr P BOYER  
Inserm, Clinique des maladies  
mentales et de l'encéphale  
100, rue de la Santé  
75014 Paris  
Fax: (33) 1 45 65 89 43

Dr Y LECRUBIER  
Hôpital de la Salpêtrière  
Pavillon Clémambault U 302  
47, boulevard de l'Hôpital  
75651 Paris Cedex 13  
Fax: (33) 1 44 23 92 49

Submissions from Eastern Europe  
should, for the time being, be sent  
to the Deputy Editors.

Submissions from non European  
countries can be sent to any of the  
preceding editors.

Papers that do not conform to the general criteria for publication in *European Psychiatry* will be returned immediately to authors to avoid unnecessary delay in submission elsewhere.

**Organisation of the manuscript.** Manuscripts should be typewritten/printed double-spaced with wide margins throughout. Title page, abstracts, tables, legends to figures and reference list should each be provided on separate sheets of paper. The title page should include: the title, the name(s) and affiliation(s) of the author(s), an address for correspondence, and telephone/fax numbers for editorial queries. All original and review articles should include an abstract (a single paragraph of no more than 150 words) and 3-6 key words for abstracting and indexing purposes. For original and review articles, the text should be ordered under the following headings: Introduction, Subjects and methods/Materials and methods, Results, Discussion (may be combined with Results), Conclusion, Acknowledgments (when appropriate), References.

• **Short communications** should not exceed 1500 words or the equivalent space including figures and tables, with abstracts of no more than 50 words.

• **Rapid communications** should not exceed 1000 words or the equivalent space. The summary should consist of no more than 50 words. Only one table and one figure are accepted. Figures (glossy) should be submitted in a form suitable for direct reproduction. References should be limited to a maximum of 10 and are in addition to the 1000 words. To ensure rapid publication, articles must meet a high standard, both in terms of scientific content and presentation. Following acceptance of an article in this category, no further modifications by the author will be allowed. Rapid communications are considered to be articles comprising preliminary but consistent results and will be published within three months following acceptance.

• **Letters to the Editor** (maximum of 500 words) will be processed rapidly and therefore should be sent to the Deputy Editors. To ensure speedy publication, please adhere strictly to the general instructions on style and arrangement; provide only figures and tables suitable for direct reproduction.

• **Illustrations.** Photographs should be presented as glossy prints with high contrast. Figures should be completely and consistently lettered, the size of the lettering being appropriate to that of the illustration, taking into account the necessary reduction in size. Illustrations should be designed to fit either a single column or the full text width. Each illustration should be clearly marked on the reverse side with the name of the author(s), the number of the illustration and its orientation (top).

Colour figures will be included subject to the author's agreement to defray part of the cost.

• **Tables.** All tables must be cited in the text, have titles and be numbered consecutively with roman numerals. Only horizontal lines should be included, and kept to a minimum.

• **References** (Harvard System) Authors are responsible for the accuracy of the references. Only published articles and those in press (the journal should be stated) may be included; unpublished results and personal communications should be cited as such in the text. In the text, a reference should be cited by author and date; when there are more than two authors, state the first author's name followed by '*et al.*'. References should be arranged alphabetically at the end of the paper and include, in the following order: all authors (surnames followed by initials), year in parentheses, journal title (abbreviated according to the Serial Sources for the Biosis Data Base), volume number, and inclusive page numbers.

For a book, the editors' names, book title, publisher and publisher's location should appear before volume and page numbers.

### Book

Takahashi R, R Flor-Henry P, Gruzelier J, Niwa SI (1987) *Dynamics, Laterality and Psychopathology*. Elsevier, Amsterdam

### Chapter in a book

Pinard G, Tetreault L (1974) Concerning semantic problems in psychological evaluation. In: *Psychological Measurements in Psychopharmacology. Modern Problems in Pharmacopsychiatry*, Vol 7 (Pichot P, ed) Karger, Basel, 8-22

• **Nomenclature.** Metric units must be used throughout, laboratory units must be followed by SI units. The generic name of a drug should be used unless the specific trade name of the drug is directly relevant to the discussion.

• **Proofs and reprints.** Proofs will be sent to the author indicated on the title page; no changes to the original manuscript will be allowed at this stage. They should be carefully corrected and returned (preferably by fax) to the publisher within 48 hours after reception. If this period is exceeded, the galley proofs will be proofread by the editorial staff of the publishing house only and printed without the author's corrections.

25 free reprints are supplied per contribution. Additional reprints may be ordered at the prices quoted on the order form sent to the corresponding author.

• **Copyright.** As soon as the article is published, the author is considered to have transferred his rights to the publisher. Requests for reproduction should be sent to the latter.

### Examples:

#### Journal article

Löö H, Rein W, Souche A, Dufour H, Guelfi JD, Malka R, Olié JP (1990) Psychopathological and sociodemographic characteristics of 1231 depressed patients with and without co-existing alcoholism. *Psychiatr & Psychobiol* 5, 249-256

# PROZAC® 20mg

Chlorhydrate de Fluoxétine

## UN PROGRÈS MAJEUR DANS LE TRAITEMENT DES ÉTATS DÉPRESSIFS



**Gélules :** boîte de 14. **Composition :** Gélules chlorhydrate de fluoxétine correspondant à 20 mg de fluoxétine base. **Indications thérapeutiques :** états dépressifs. **Contre-indications :** association avec les IMAO - hypersensibilité connue à la fluoxétine - enfant de moins de 15 ans. **Mises en garde :** un délai de 14 jours doit être respecté entre l'arrêt de l'IMAO et le début du traitement par fluoxétine. Un délai de cinq semaines doit être respecté entre l'arrêt de la fluoxétine et le début du traitement par un IMAO. En cas de survenue d'éruption cutanée ou d'autres phénomènes probablement allergiques, interrompre le traitement. **Précautions d'emploi :** le suicide constitue un risque inhérent à l'état dépressif. L'association de la fluoxétine avec une sismothérapie n'est pas recommandée. En cas de dysfonctionnement hépatique, une réduction de la posologie doit être envisagée. L'apparition de troubles respiratoires persistants peut demander une évaluation plus approfondie. Des cas d'hyponatrémie ont été rapportés, pour la plupart, chez des patients âgés, prenant des diurétiques ou encore hypovolémiques. Chez les patients diabétiques, la fluoxétine est susceptible de modifier l'équilibre glycémique. Une adaptation du traitement antidiabétique à l'instauration ou à l'arrêt de Prozac peut être nécessaire. L'innocuité durant la grossesse et l'allaitement n'a pas été établie. Prévenir les conducteurs d'automobiles et les utilisateurs de machine d'une éventuelle modification de leur comportement. **Interactions médicamenteuses :** la fluoxétine peut allonger la demi-vie du diazépam. Une augmentation des taux plasmatiques des anti-dépresseurs tricycliques ou de la carbamazépine a pu être observée en cas de co-administration avec PROZAC 20 mg. Des variations de la lithiéme ont été rapportées lors de l'association du lithium et de la fluoxétine. Il convient donc de surveiller la lithiéme en cas de coprescription. **Effets indésirables :** peuvent être observés : nausées, diarrhées, bouche sèche, céphalées, nervosité, insomnie, anxiété, somnolence, tremblements et éruptions cutanées, rarement associées à une urticaire, un syndrome du canal carpien, de la fièvre et/ou un dysfonctionnement hépatique. Eruptions et symptômes concomitants sont réversibles à l'arrêt du traitement. Des réactions maniaques ou psychotiques peuvent apparaître chez certains sujets prédisposés. La fluoxétine peut induire une perte de poids banale n'entraînant que très rarement l'arrêt du traitement. Exceptionnellement, on peut noter une élévation des transaminases sériques et/ou une leucopénie asymptomatique, réversibles à l'arrêt du traitement. Confusion mentale, idées suicidaires, dyskinésies, thrombocytopénie, hyperprolactinémie, saignements vaginaux ont été enregistrés. Des symptômes respiratoires telle une dyspnée persistante ou une augmentation de la toux ont été exceptionnellement rapportés (cf. précautions d'emploi). **Mode d'emploi et posologie :** la posologie recommandée est de 20 mg/jour. Cette posologie peut être augmentée jusqu'à 60 mg/jour. Elle doit être diminuée en cas de cirrhose hépatique. **Surdosage :** des crises comitiales ont été rapportées. Pour informations complémentaires, se reporter au dictionnaire des spécialités pharmaceutiques. Liste 1 - AMM. 331 009.5 (1988) gélules. **Prix :** 88.20 F (14 gélules). Remb. Soc. à 70%. Agréé au collect. **Cout du traitement journalier :** 6,30 F.

