

Chemical Abundances of Metal-poor stars in Dwarf Galaxies

Kim A. Venn¹, Pascale Jablonka², Vanessa Hill³, Else Starkenburg⁴,
Bertrand Lemasle⁵, Matthew Shetrone⁶, Mike Irwin⁷, John Norris⁸,
David Yong⁸, Gerry Gilmore⁷, Stephania Salvadori⁹, Asa Skuladottir⁹
and Eline Tolstoy⁹

¹Dept. of Physics & Astronomy, University of Victoria, Victoria, BC, V8P 5C2, Canada
email: kvenn@uvic.ca

²Laboratoire d'astrophysique, EPFL, Observatoire de Sauverny, 1290 Versoix, Switzerland

³Obs. de la Cote d'Azur, 06304 Nice Cedex 4, France

⁴Leibniz-Institute für Astrophysik Potsdam, 14482 Potsdam, Germany

⁵Anton Pannekoek Inst. for Astronomy, Univ. of Amsterdam, 1090 GE Amsterdam,
Netherlands

⁶McDonald Observatory, Univ. of Texas at Austin, Austin TX 78712, USA

⁷Institute of Astronomy, Univ. of Cambridge, Madingley Road, Cambridge, CB3 0HA, UK

⁸RSAA, Australian National Univ., Mount Stromlo Observatory, Weston ACT 2611, Australia

⁹Kapteyn Astronomical Inst., Univ. of Groningen, 9747 AD Groningen, Netherlands

Abstract. Stars in low-mass dwarf galaxies show a larger range in their chemical properties than those in the Milky Way halo. The slower star formation efficiency make dwarf galaxies ideal systems for testing nucleosynthetic yields. Not only are alpha-poor stars found at lower metallicities, and a higher fraction of carbon-enhanced stars, but we are also finding stars in dwarf galaxies that appear to be iron-rich. These are compared with yields from a variety of supernova predictions.

Keywords. stars: abundances, galaxies: dwarf, evolution, Local Group, stellar content.

1. Introduction

It has been known for more than a decade now that “the overwhelming majority of Milky Way stars, those in the Galactic thick disk and thin disk, seem to have nothing at all to do with dwarf galaxy origins” (Gilmore 2012). As summarized by Tolstoy *et al.* (2009), the ratio of alpha-elements to iron are lower in the stars in dwarf galaxies for metallicities above $[\text{Fe}/\text{H}] \sim -2$. This has been found for stars in the Sculptor, Carina, Fornax, and LMC dwarf galaxies, and the Sagittarius dwarf galaxy remnant. The early results have been confirmed by larger data samples (e.g., Lemasle *et al.* 2012, 2014, Hendricks *et al.* 2014), and extended to lower mass dwarfs, such as Bootes I and Segue I (e.g., Feltz *et al.* 2009, Norris *et al.* 2010, Frebel *et al.* 2014).

The picture is less clear when comparing the chemistry of the stars in the Galactic halo with those in dwarf galaxies with metallicities $[\text{Fe}/\text{H}] < -2.0$. An examination of the stellar abundances of $[\text{Ca}/\text{Fe}]$ and $[\text{Mg}/\text{Fe}]$ in stars below metallicity $[\text{Fe}/\text{H}] = -2$ is shown in Fig. 1. The uncertainties in each individual stars are typically < 0.2 dex. While the general trend of high $[\alpha/\text{Fe}]$ ratios (above the solar ratio) appears in all systems, the dispersion is larger in the dwarf galaxies, and includes several stars with very low $[\alpha/\text{Fe}]$ abundances (below the solar ratio).




Figure 1. $[\text{Mg}/\text{Fe}]$ and $[\text{Ca}/\text{Fe}]$ for stars in the Milky Way (grey) and dwarf galaxies. These include abundances summarized by Frebel 2010, and additional new references for Sculptor (blue; Jablonka *et al.* 2015, Simon *et al.* 2015, Starkenburg *et al.* 2013, Tafelmeyer *et al.* 2010), Sextans (green; Tafelmeyer *et al.* 2010) Carina (orange; Norris *et al.* 2016, Venn *et al.* 2012, Lemasle *et al.* 2012) Draco and UMi (yellow), and the ultra faint dwarf galaxies (black; Frebel 2014 for Segue 1 and Aden *et al.* 2011 for Hercules).

2. Element Abundances in Metal-Poor Stars

At the lowest metallicities, the detailed chemical abundances have been modelled as enrichment by a variety of core collapse SN, including a range of progenitor masses, SN explosion energies, and mixing and fallback prescriptions. Iwamoto *et al.* (2005) have found that "faint supernovae" with extensive mixing and fallback during the SN II explosion produce decreasing yields with increasing atomic number (mitigated by the odd-even effect); and that these yields reproduce the element distribution for the ultra metal-poor stars HE1327-2326 and HE0107-5240 better than predictions from the metal-free massive models by Heger & Woosley (2002, 2008). Neither of these models produce the heavy neutron capture elements.

The core collapse supernova models by Wanaajo (2013) can reproduce the solar-system r-process distribution by tuning the initial core masses. These models are also able to reproduce the heavy element distribution in metal-poor Galactic stars, such as CS 31082-001. However, Wanaajo *et al.* (2014) have also been able to make these same heavy element distributions with a variety of compact binary mergers. This latter scenario is an interesting alternative for the site of the r-process (also see Shen *et al.* 2015), though rely on timescale arguments for making and merging two neutron stars.

In Fig. 2, the $[\text{Ba}/\text{Fe}]$ ratios for metal-poor stars in the Galaxy and dwarf galaxies are shown. The general distribution and dispersions are quite similar for the majority of stars, implying the sites and yields for the r-process are similar in these systems. Again, the uncertainties in the abundances ratios are estimated as < 0.2 dex. Only the upper limits on $[\text{Ba}/\text{Fe}]$ for stars in the dwarf galaxies show an unexpected result, as highlighted by Frebel *et al.* (2014). The intermediate metallicity stars (with $[\text{Fe}/\text{H}] > -2.5$) in the ultra faint dwarf galaxies, Segue 1, Com Ber, and Hercules, have extremely low $[\text{Ba}/\text{Fe}]$ values or upper limits. This suggests these stars formed from gas that was not enriched in r-process elements, at all, yet they have high iron abundances. This is not seen in the




Figure 2. $[\text{Ba}/\text{Fe}]$ for stars in the Milky Way (grey) and dwarf galaxies (colours as in 1). Upper limits for stars in dwarf galaxies (only) are noted as open downward-pointing open triangles. Detection limits for the Ba II 4554 Å spectral line are shown as green lines (10 mÅ as dotted, 3 mÅ as dashed, 1 mÅ as solid), for a red giant with $T = 4500$ K, following Roederer (2013).

Galaxy nor the classical dwarf galaxies. This observation has been interpreted by Frebel & Norris (2015) as the result of a single (or few) high-mass metal-free supernova event that also removed the gas in these lowest mass galaxies.

3. Iron-rich Stars in Dwarf Galaxies

At intermediate metallicities, $[\text{Fe}/\text{H}] > -2.5$, another phenomenon is found in dwarf galaxies, where some stars appear to be enriched in iron-group elements. This is seen as an unusually low $[\text{X}/\text{Fe}]$ ratios for nearly all elements, except those in the iron-group. One example is Star 612 in the Carina dwarf galaxy (Venn *et al.* 2012), though others have now been found in Carina (Norris *et al.* 2016) and Sculptor (Jablonka *et al.* 2015). Similar stars have been found in the (outer) halo of the Galaxy by Ivans *et al.* (2003) and Jacobson *et al.* (2015).

One possibility is that the iron-enrichment comes from inhomogeneous mixing of the interstellar medium at later times. Pockets of SN Ia material may occur if the gas is poorly mixed. Some chemical evolution models (e.g., Revaz & Jablonka 2012, Wise *et al.* 2012, Romano *et al.* 2015) support inhomogeneous mixing on sufficiently long timescales only in the lower mass dwarf galaxies.

Another option has been suggested by Kobayashi *et al.* (2015) where SN Iax have a more significant role in the chemical evolution of a dwarf galaxy. SN Iax are those from a higher mass (hybrid, C+O+Ne) white dwarf, formed at lower metallicities (even as low as $[\text{Fe}/\text{H}] = -2.5$).

4. Carbon-rich Stars in Dwarf Galaxies

Carbon-enhancements ($[\text{C}/\text{Fe}] > 0.7$) are commonly found in metal-poor stars in the Galaxy and typically associated with binary systems (e.g., Starkeburg *et al.* 2014). However, the CEMP-no stars may be a different class of these stars, formed from primordial faint supernovae and dominating the early metal enrichment in dwarf galaxies. While CEMP-no stars are found in the Galaxy and the ultra faint dwarf galaxies (e.g., Beers & Christlieb 2005, Norris *et al.* 2013), they have only recently been found also in the classical dwarf galaxies (Skuladottir *et al.* 2015). Initial interpretations were that the early chemical evolution of the classical dwarf galaxies may have differed from those of the ultra faint dwarfs, however a recent analysis of cosmological models of dwarf galaxies has shown that CEMP-no stars would have formed in all systems, and with equal fractions in the populations with $[\text{Fe}/\text{H}] < -4$ (Salvadori *et al.* 2015). The difficulty is in finding these stars once the peak in the metallicity distribution function moves from $[\text{Fe}/\text{H}] = -3$ to -2 , as in the higher mass classical dwarf galaxies. Also the classical dwarf galaxies are further away, meaning that only the brightest stars can be observed. Comparisons of the metallicities of the stars from CaT surveys versus those from high resolution studies shows that more metal-poor stars are detected at lower magnitudes (e.g., Lemasle *et al.* 2012).

5. Conclusions

The element abundance ratios in the most metal-poor stars ($[\text{Fe}/\text{H}] < -3$) in dwarf galaxies are excellent tests for the variety of SN II yields and other supernova models. Most elements show larger dispersions in $[\text{X}/\text{Fe}]$ in dwarf galaxies, than similar stars in the Galactic halo. Spectral lines of some of the more interested elements (such as barium) can be so weak that we are near the detection limits of our high resolution spectrographs, even on the 8-10 meter telescopes. Also, the CEMP stars are likely in all dwarf galaxies, though can be harder to find in the more distant and higher mass (higher mean metallicity) classical dwarf galaxies.

The intermediate-metallicity stars ($[\text{Fe}/\text{H}] \sim -2$) can show the largest deviations from similar metallicity stars in the Galactic halo, due to differences in the later chemical evolution stages of the dwarf galaxies. This can include effects due to inhomogeneous mixing of the interstellar medium, including SN Ia pockets of iron-enrichment, in the lower mass dwarf galaxies.

References

- Aden, D. *et al.* 2011, *A&A*, 525, 153
 Feltzing, S. *et al.* 2009, *A&A*, 508, 1
 Frebel, A. 2010, in *Astron. Nachr.* 331, 474
 Frebel, A., Simon, J. D., Kirby, E. N. 2014, *ApJ*, 786, 74
 Frebel, A., Norris, J. E. 2015, *ARAA*, 53, 631
 Gilmore, G. 2012, in *Assembling the Puzzle of the Milky Way*, Le Grand-Bourmand, France.
 Heger, A., Woosley, S. E. 2003, *ApJ*, 567, 532
 Heger, A., Woosley, S. E. 2010, *ApJ*, 724, 341
 Hendricks, B. *et al.* 2014, *ApJ*, 785, 102
 Ivans I., *et al.* 2003, *ApJ*, 592, 906
 Iwamoto, N., *et al.* 2005, *Sci*, 309, 451
 Jablonka, P., *et al.* 2015, *A&A*, 583, 67
 Jablonka, P., *et al.* 2016, in prep
 Kobayashi, C., *et al.* 2015, *ApJ*, 804, 24

- Lemasle, B., *et al.* 2012, *A&A*, 538, 100
Lemasle, B., *et al.* 2014, *A&A*, 572, 88
Norris, J. E., *et al.* 2010, *ApJ*, 723, 1632
Norris, J. E., *et al.* 2013, *ApJ*, 762, 28
Norris, J. E., *et al.* 2016, in prep
Revaz, Y., Jablonka, P 2012, *A&A*, 538, 82
Roederer, I.U. 2013 *AJ*, 145, 26
Romano, D., Bellazzini, M., Starkeburg, E., Leaman, R. 2015, *MNRAS*, 446, 4220
Salvadori, S., Skuladottir, A., Tolstoy, E. 2015 *MNRAS*, 454, 1320
Shen, S., Cooke, R. J., Ramirez-Ruiz, E., *et al.* 2015, *ApJ*, 807, 115
Simon, J. D. *et al.* 2015, *ApJ*, 802, 93
Skuladottir, A., *et al.* 2015, *A&A*, 574, 129
Starkeburg, E., *et al.* 2014, *MNRAS*, 441, 1217
Starkeburg, E., *et al.* 2013, *MNRAS*, 429, 725
Tafelmeyer, M., *et al.* 2010, *A&A*, 524, A58
Tolstoy, E., *et al.* 2009, *ARAA*, 47, 371
Venn, K. A., *et al.* 2012, *ApJ*, 751, 102
Wanajo, S. 2013, *ApJ*, 770, 22
Wanajo, S. *et al.* 2014, *ApJ*, 789, 39
Wise, J. *et al.* 2012, *ApJ*, 745, 50