


# Are COVID-19 survivors at increased risk for suicide?

Leo Sher 

James J. Peters Veterans' Administration Medical Center, Bronx, NY, USA and Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, NY, USA

## Letter to the Editor

**Cite this article:** Sher L. (2020) Are COVID-19 survivors at increased risk for suicide? *Acta Neuropsychiatrica* 32:270. doi: [10.1017/neu.2020.21](https://doi.org/10.1017/neu.2020.21)

Received: 27 April 2020  
Revised: 30 April 2020  
Accepted: 2 May 2020  
First published online: 4 May 2020

**Author for correspondence:**  
Leo Sher, Email: [Leo.Sher@mssm.edu](mailto:Leo.Sher@mssm.edu)

The coronavirus disease 19 (COVID-19) outbreak first emerged in China late last year. The COVID-19 epidemic has spread to all continents. Millions of people got sick with COVID-19. In this letter to the editor, I suggest that the COVID-19 survivors especially individuals who had severe COVID-19 are at increased suicide risk. Suicidality among individuals who had COVID-19 may be related to both psychological and neurobiological factors.

Psychological factors that may increase suicide risk among COVID-19 patients include learning about their diagnosis, anxiety, and distress related to symptoms of the disease and stress related to hospitalisation and hospital treatment. Realising that they have COVID-19 may be very stressful, especially for low-resilient individuals. Symptoms of the disease, especially severe symptoms, as well as social isolation and fear of infecting other people may lead to a serious psychological trauma. Individuals who needed an admission to an intensive care unit (ICU) are at especially high risk of developing post-traumatic stress disorder (PTSD), depression, anxiety, sleep abnormalities, and cognitive impairments (McGiffin *et al.*, 2016). Ventilation adjustments, gaps in anaesthesia and/or analgesia, long sedation, restraint use, and other ICU-related factors contribute to a profound psychological effect of ICU hospitalisations. A recent study in China indicated that 96.2% of recovering COVID-19 patients had significant post-traumatic stress symptoms (Bo *et al.*, 2020).

Multiple lines of evidence indicate that stress-related disorders including depression, PTSD, and sleep disorders are associated with suicidal ideation, suicide attempts, and death by suicide (Sher, 2019). Sleep abnormalities are a stand-alone risk factor for suicidal behaviour. Cognitive impairments are also associated with suicidality (Sher, 2019). Many recovering COVID-19 patients have physical symptoms for a long time and experience psychosocial difficulties such as loss of employment and financial issues. Both physical symptoms and psychosocial stressors contribute to suicidal behaviour (Sher, 2019).

COVID-19 has neurobiological effects (Asadi-Pooya & Simani, 2020). Studies have shown human coronavirus infections are associated with neuroinvasion and neurotropism (Asadi-Pooya & Simani, 2020). A recent review of the effect of COVID-19 on the central nervous system indicates that neurological manifestations are present in about 25% of the patients (Asadi-Pooya & Simani, 2020). Headache, dizziness, acute ischaemic stroke, ataxia, seizures, and other neurological conditions have been observed in COVID-19 patients. Neurological conditions including ischaemic stroke and headache are associated with increased suicide risk (Hudzik & Marek, 2014).

COVID-19 survivors should be regarded as individuals at elevated risk for suicide. The single most significant predictor of suicide is the presence of depression. Recovered COVID-19 patients need to be screened for depression and suicidality. Many coronavirus disease survivors will need long-term psychological interventions. There should be specific strategies to enhance the psychological condition of COVID-19 survivors and reduce suicidality in this population. We need to examine what kind of early interventions in coronavirus disease survivors may decrease psychiatric morbidity and suicidality in the future.

## References

- Asadi-Pooya AA and Simani L** (2020) Central nervous system manifestations of COVID-19: a systematic review. *Journal of Neurological Sciences* 413, 116832. doi: [10.1016/j.jns.2020.116832](https://doi.org/10.1016/j.jns.2020.116832).
- Bo HX, Li W, Yang Y, Wang Y, Zhang Q, Cheung T, Wu X and Xiang YT** (2020) Posttraumatic stress symptoms and attitude toward crisis mental health services among clinically stable patients with COVID-19 in China. *Psychological Medicine* doi: [10.1017/S0033291720000999](https://doi.org/10.1017/S0033291720000999) [Epub ahead of print].
- Hudzik TJ and Marek GJ** (2014) Neurological disease and suicidal behavior. In Cannon KE, Hudzik TJ (eds.). *Suicide: Phenomenology And Neurobiology*. Cham, Switzerland: Springer International Publishing.
- McGiffin JN, Galatzer-Levy IR and Bonanno GA** (2016) Is the intensive care unit traumatic? what we know and don't know about the intensive care unit and posttraumatic stress responses. *Rehabilitation Psychology* 61, 120–131. doi: [10.1037/rep0000073](https://doi.org/10.1037/rep0000073).
- Sher L** (2019) Resilience as a focus of suicide research and prevention. *Acta Psychiatrica Scandinavica* 140, 169–180. doi: [10.1111/acps.13059](https://doi.org/10.1111/acps.13059).

© The Author(s) 2020. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

