

Kolakoski-(3, 1) Is a (Deformed) Model Set

Michael Baake and Bernd Sing

Abstract. Unlike the (classical) Kolakoski sequence on the alphabet $\{1, 2\}$, its analogue on $\{1, 3\}$ can be related to a primitive substitution rule. Using this connection, we prove that the corresponding bi-infinite fixed point is a regular generic model set and thus has a pure point diffraction spectrum. The Kolakoski-(3, 1) sequence is then obtained as a deformation, without losing the pure point diffraction property.

1 Introduction

A one-sided infinite sequence ω over the alphabet $\mathcal{A} = \{1, 2\}$ is called a (classical) *Kolakoski sequence* (named after W. Kolakoski who introduced it in 1965, see [21]), if it equals the sequence defined by its run lengths, e.g.:

$$(1) \quad \omega = \underbrace{2\ 2}_2 \underbrace{1\ 1}_2 2\ 1 \underbrace{2\ 2}_2 1 \underbrace{2\ 2}_2 \underbrace{1\ 1}_2 2 \underbrace{1\ 1}_2 \dots = \omega.$$

Here, a *run* is a maximal subword consisting of identical letters. The sequence $\omega' = 1\omega$ is the only other sequence which has this property.

One way to obtain ω of (1) is by starting with 2 as a seed and iterating the two substitutions

$$\sigma_0: \begin{matrix} 1 & \mapsto & 2 \\ 2 & \mapsto & 22 \end{matrix} \quad \text{and} \quad \sigma_1: \begin{matrix} 1 & \mapsto & 1 \\ 2 & \mapsto & 11, \end{matrix}$$

alternatingly, i.e., σ_0 substitutes letters on even positions and σ_1 letters on odd positions (we begin counting at 0):

$$2 \mapsto 22 \mapsto 2211 \mapsto 221121 \mapsto 221121221 \mapsto \dots$$

Clearly, the iterates converge to the Kolakoski sequence ω (in the obvious product topology), and ω is the unique (one-sided) fixed point of this iteration.

One can generalize this by choosing a different alphabet $\mathcal{A} = \{p, q\}$ (we are only looking at alphabets with $\text{card}(\mathcal{A}) = 2$), e.g., $\mathcal{A} = \{1, 3\}$, which is the main focus of this paper. Such a (generalized) Kolakoski sequence, which is also equal to the sequence of its run lengths, can be obtained by iterating the two substitutions

$$\sigma_0: \begin{matrix} q & \mapsto & p^q \\ p & \mapsto & p^p \end{matrix} \quad \text{and} \quad \sigma_1: \begin{matrix} q & \mapsto & q^q \\ p & \mapsto & q^p \end{matrix}$$

Received by the editors June 20, 2002.
 AMS subject classification: 52C23, 37B10, 28A80, 43A25.
 ©Canadian Mathematical Society 2004.

alternatingly. Here, the starting letter of the sequence is p . We will call such a sequence Kolakoski- (p, q) sequence, or $\text{Kol}(p, q)$ for short. The classical Kolakoski sequence ω of (1) is therefore denoted by $\text{Kol}(2, 1)$ (and ω' by $\text{Kol}(1, 2)$).

While little is known about the classical Kolakoski sequence (see [15]), and the same holds for all $\text{Kol}(p, q)$ with p odd and q even or vice versa (see [31]), the situation is more favourable if p and q are either both even or both odd. If both are even, one can rewrite the substitution as a substitution of constant length by building blocks of 4 letters (see [31, 32]). Spectral properties can then be deduced by a criterion of Dekking [13]. The case where both symbols are odd will be studied in this paper exemplarily on $\text{Kol}(3, 1)$.

It is our aim to determine structure and order of the sequence $\text{Kol}(3, 1)$. This will require two steps: First, we relate it to a unimodular substitution of Pisot type and prove that the corresponding aperiodic point set is a regular generic model set. Second, we relate this back to the original $\text{Kol}(3, 1)$ by a deformation. Here, the first step is a concrete example of the general conjecture that all unimodular substitutions of Pisot type are regular model sets (however, not always generic). This general conjecture cannot be proved by an immediate application of our strategy, but we hope that our method sheds new light on it.

Remark Every $\text{Kol}(p, q)$ can uniquely be extended to a bi-infinite (or two-sided) sequence. The one-sided sequence (to the right) is $\text{Kol}(p, q)$ as explained above. The added part to the left is a reversed copy of $\text{Kol}(q, p)$, e.g., in the case of the classical Kolakoski sequence of (1), this reads as

$$\dots 11221221211221|22112122122112 \dots,$$

where “|” denotes the seamline between the one-sided sequences. Note that, if $q = 1$ (or $p = 1$), the bi-infinite sequence is mirror symmetric around the first position to the left (right) of the seamline. The bi-infinite sequence equals the sequence of its run lengths, if counting is begun at the seamline. Alternatively, one can get such a bi-infinite sequence by starting with $q|p$ and applying the two substitutions to get $\sigma_1(q)|\sigma_0(p)$ in the first step and so forth. This also implies that $\text{Kol}(p, q)$ and $\text{Kol}(q, p)$ will have the same spectral properties, and it suffices to study one of them.

2 Kol(3, 1) as Substitution

If both letters are odd numbers, one can build blocks of 2 letters and obtain an (ordinary) substitution. Setting¹ $A = 33, B = 31$ and $C = 11$ in the case of $\text{Kol}(3, 1)$, this substitution σ and its *substitution matrix* \mathbf{M} (sometimes called *incidence matrix* of the substitution) are given by

$$(2) \quad \begin{array}{l} A \mapsto ABC \\ \sigma: B \mapsto AB \\ C \mapsto B \end{array} \quad \text{and} \quad \mathbf{M} = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 0 \\ 0 & 1 & 0 \end{pmatrix},$$

¹That $\text{Kol}(3, 1)$ can be related to a substitution is well-known, e.g., in [14], a substitution over an alphabet with four letters is given, while [33] uses the same substitution with three letters as we do. We thank the referee for pointing this last reference out to us.

where the entry M_{ij} is the number of occurrences of j in $\sigma(i)$ ($i, j \in \{A, B, C\}$; sometimes the transposed matrix is used). A bi-infinite fixed point can be obtained as follows:

$$(3) \quad B|A \mapsto AB|ABC \mapsto ABCAB|ABCABB \mapsto \dots$$

This corresponds to

$$(4) \quad \dots 3331113331|333111333131 \dots$$

which is the unique bi-infinite Kol(3, 1) according to our above convention. The matrix \mathbf{M} is primitive because \mathbf{M}^3 has positive entries only. The characteristic polynomial $P(x)$ of \mathbf{M} is

$$(5) \quad P(x) := \det(x\mathbf{1} - \mathbf{M}) = x^3 - 2x^2 - 1,$$

which is irreducible over \mathbb{Z} (there is no solution mod 3) and over \mathbb{Q} (every rational algebraic integer is an integer). The discriminant D of $P(x)$ is $D = \frac{59}{108}$, so $P(x)$ has one real root α and two complex conjugate roots β and $\bar{\beta}$. One gets

$$2.21 \approx \alpha > 1 > |\beta| \approx 0.67 > 0,$$

wherefore α is a *Pisot-Vijayaraghavan number* (i.e., an algebraic integer greater than 1 whose algebraic conjugates are all less than 1 in modulus), and σ is a substitution of *Pisot type*. Since $\det(\mathbf{M}) = 1$, the roots α, β and $\bar{\beta}$ are also *algebraic units*, and the associated substitution is said to be *unimodular*. Note that $\operatorname{Re}(\beta) = 1 - \frac{\alpha}{2}$. If necessary, we will choose β such that $\operatorname{Im}(\beta) > 0$ in the following calculations (the other possibility only leads to overall minus signs).

There is a natural geometric representation of such a substitution by inflation, compare [24]. Here, one associates bond lengths (or intervals) ℓ_A, ℓ_B and ℓ_C to each letter. These bond lengths are given by the components of the right eigenvector which belongs to the (real) eigenvalue α and is unique (up to normalization) by the Perron-Frobenius theorem. The normalization can be chosen so that

$$\ell_A = \alpha^2 - \alpha \approx 2.66, \quad \ell_B = \alpha \approx 2.21 \quad \text{and} \quad \ell_C = 1.$$

Inflating the bond lengths by a factor of α and dividing them into original intervals just corresponds to the substitution (because $\alpha \cdot \ell_A = \ell_A + \ell_B + \ell_C$, etc.). We will denote this realization of the bi-infinite fixed point with natural bond lengths (respectively the point set associated with this realization where we mark the left endpoints of the intervals by their name) by $\Sigma \text{Kol}(3, 1)$, reserving “Kol(3, 1)” for the case of unit (or integer) bond lengths.

On the other hand, the frequencies ρ_A, ρ_B and ρ_C of the letters in the infinite sequence are given by the components of the left eigenvector of \mathbf{M} to the eigenvalue α . This gives

$$\rho_A = \frac{1}{2}(-\alpha^2 + 3\alpha - 1) \approx 0.38, \quad \rho_B = \alpha^2 - 2\alpha \approx 0.45, \quad \rho_C = \frac{1}{2}(-\alpha^2 + \alpha + 3) \approx 0.17,$$

with $\rho_A + \rho_B + \rho_C = 1$. Therefore, the average bond length ℓ in the geometric representation is

$$(6) \quad \ell = \rho_A \cdot \ell_A + \rho_B \cdot \ell_B + \rho_C \cdot \ell_C = \frac{1}{2}(-\alpha^2 + \alpha + 7) \approx 2.17,$$

and the frequencies of 3s and 1s in Kol(3, 1) can easily be calculated to be $\rho_3 = \frac{1}{2}(\alpha - 1) \approx 0.60$ and $\rho_1 = \frac{1}{2}(-\alpha + 3) \approx 0.40$.

Remark In the case where p and q are odd (positive) integers, one gets unimodular substitutions of Pisot type iff $p = q \pm 2$. More generally, one gets substitutions of Pisot type iff $2 \cdot (p + q) \geq (p - q)^2$ holds. Otherwise, all the eigenvalues are greater than 1 in modulus, see [31].

3 Model Set and IFS

A model set $\Lambda(\Omega)$ (or *cut-and-project set*) in physical space \mathbb{R}^d is defined within the following general cut-and-project scheme [26, 3]

$$(7) \quad \begin{array}{ccccc} & & \mathbb{R}^d & \xleftarrow{\pi} & \mathbb{R}^d \times H & \xrightarrow{\pi_{\text{int}}} & H \\ & & & & \cup & & \\ & & & & \Gamma & & \end{array}$$

\swarrow 1-1 dense \searrow

where the *internal space* H is a locally compact Abelian group, and $\Gamma \subset \mathbb{R}^d \times H$ is a *lattice*, i.e., a co-compact discrete subgroup of $\mathbb{R}^d \times H$. The projection $\pi_{\text{int}}(\Gamma)$ is assumed to be dense in internal space, and the projection π into physical space has to be one-to-one on Γ . The model set $\Lambda(\Omega)$ is

$$\Lambda(\Omega) = \{ \pi(x) \mid x \in \Gamma, \pi_{\text{int}}(x) \in \Omega \} \subset \mathbb{R}^d,$$

where the *window* $\Omega \subset H$ is a relatively compact set with non-empty interior. If we set $L = \pi(\Gamma) \subset \mathbb{R}^d$, we can define, for $x \in L$, the *star map* $^* : L \mapsto H$ by $x^* = \pi_{\text{int}} \circ (\pi|_L)^{-1}(x)$, see [5]. So we have $\Gamma = \{(x, x^*) \mid x \in L\}$ and $L^* = \pi_{\text{int}}(\Gamma)$. If the boundary $\partial\Omega$ of the window has vanishing Haar measure in H , we call $\Lambda(\Omega)$ a *regular model set*. If, in addition, $\pi_{\text{int}}(\Gamma) \cap \partial\Omega = \emptyset$, the model set is called *non-singular* or *generic*.

Every model set is also a *Delone set* (or *Delaunay set*), i.e., it is both *uniformly discrete*² and *relatively dense*³. A Delone set X is a *Meyer set*, if also $X - X$ is a Delone set. Every model set is a Meyer set, see [25].

We will now construct a model set $\Lambda(\Omega)$ and—in a first step—show that this model set differs from Σ Kol(3, 1) at most on positions of density 0. By Galois conjugation (see [24, 11]), which here corresponds to the star map as we will see, we find

²A set Λ is *uniformly discrete* if $\exists r > 0$ s.t. every open ball of radius r contains at most one point of Λ .
³A set Λ is *relatively dense* if $\exists R > 0$ s.t. every closed ball of radius R contains at least one point of Λ .

a lattice

$$(8) \quad \Gamma = \mathbb{Z} \cdot \mathbf{v}_A + \mathbb{Z} \cdot \mathbf{v}_B + \mathbb{Z} \cdot \mathbf{v}_C \subset \mathbb{R} \times \mathbb{C} \simeq \mathbb{R}^3$$

where

$$\mathbf{v}_A = \begin{pmatrix} \alpha^2 - \alpha \\ \operatorname{Re}(\beta^2 - \beta) \\ \operatorname{Im}(\beta^2 - \beta) \end{pmatrix}, \quad \mathbf{v}_B = \begin{pmatrix} \alpha \\ \operatorname{Re}(\beta) \\ \operatorname{Im}(\beta) \end{pmatrix} \quad \text{and} \quad \mathbf{v}_C = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}.$$

The projection π (i.e., the projection on the first coordinate) is injective on Γ because $\mathbb{Q}(\alpha)$ is a \mathbb{Q} -vector space of dimension 3 with (\mathbb{Q} -)linearly independent elements 1, α and α^2 . Also, $\pi(\Gamma) = \mathbb{Z}[\alpha]$ is dense. To see that $\pi_{\text{int}}(\Gamma)$ is dense, we note that $\pi_{\text{int}}(\Gamma) = \mathbb{Z}[\beta]$ and that 1 and β are linearly independent. So, β^n and β^{n+1} are also linearly independent for all $n \in \mathbb{N}$, and their \mathbb{Z} -span forms a two-dimensional lattice in \mathbb{C} , which is a uniformly discrete subset of $\mathbb{Z}[\beta]$. Since $|\beta| < 1$, one can choose, for every $\varepsilon > 0$, an n , such that there is a lattice point (of the lattice $\mathbb{Z} \cdot \beta^n + \mathbb{Z} \cdot \beta^{n+1}$) in every ball of radius ε , so $\mathbb{Z}[\beta]$ is dense in \mathbb{C} . Note, that π_{int} is also injective on Γ (this can be seen from $\operatorname{Re}(\beta) = 1 - \frac{\alpha}{2}$ and $\operatorname{Re}(\beta^2) = 2 - \frac{\alpha^2}{2}$). So we have established:

Proposition 1 *With $\mathbb{C} \simeq \mathbb{R}^2$, Γ of (8) and the natural projections π and π_{int} , we obtain the following cut-and-project scheme:*

$$(9) \quad \begin{array}{ccccc} & & \mathbb{R} & \xleftarrow{\pi} & \mathbb{R} \times \mathbb{R}^2 & \xrightarrow{\pi_{\text{int}}} & \mathbb{R}^2 & & \\ & & & & \cup & & & & \\ & & \mathbb{R} & & \Gamma & & \mathbb{R}^2 & & \\ & \swarrow & & & & & & \searrow & \\ & \text{dense} & & & & & & \text{dense} & \\ & \text{1-1} & & & & & & \text{1-1} & \end{array}$$

Furthermore, we have

$$\pi(\Gamma) = \mathbb{Z}[\alpha] \quad \text{and} \quad \pi_{\text{int}}(\Gamma) = \mathbb{Z}[\beta],$$

where α is the real root of (5) and β one of the complex conjugate ones. ■

In order to describe $\Sigma \operatorname{Kol}(3, 1)$, the main task is now to determine the appropriate windows Ω_A, Ω_B and Ω_C (one for each letter; $\Omega = \Omega_A \cup \Omega_B \cup \Omega_C$). For these windows, the substitution rule σ of (2) induces the following *iterated function system* (IFS for short) in internal space⁴, cf. [24]:

$$(10) \quad \begin{aligned} \Omega_A &= \beta\Omega_A \cup \beta\Omega_B \\ \Omega_B &= \beta\Omega_A + \beta^2 - \beta \cup \beta\Omega_B + \beta^2 - \beta \cup \beta\Omega_C \\ \Omega_C &= \beta\Omega_A + \beta^2. \end{aligned}$$

⁴For later reference, we write:

$$(10') \quad \Omega_A = f_1(\Omega_A) \cup f_1(\Omega_B), \quad \Omega_B = f_3(\Omega_A) \cup f_3(\Omega_B) \cup f_1(\Omega_C), \quad \Omega_C = f_0(\Omega_A),$$

where f_1 and f_3 are defined as in (13), and $f_0(z) = \beta z + \beta^2$.

This IFS is obtained as follows: We denote by Λ_A the subset of $\Sigma \text{Kol}(3, 1)$ of left endpoints of intervals of type A (of length ℓ_A), and similar for Λ_B and Λ_C (we have $\Sigma \text{Kol}(3, 1) = \Lambda_A \dot{\cup} \Lambda_B \dot{\cup} \Lambda_C$, where $\dot{\cup}$ denotes disjoint union). Then, the substitution σ of (2) induces the following equations for these Delone sets in \mathbb{R} :

$$\begin{aligned}
 \Lambda_A &= \alpha\Lambda_A \cup \alpha\Lambda_B \\
 (11) \quad \Lambda_B &= \alpha\Lambda_A + \alpha^2 - \alpha \cup \alpha\Lambda_B + \alpha^2 - \alpha \cup \alpha\Lambda_C \\
 \Lambda_C &= \alpha\Lambda_A + \alpha^2.
 \end{aligned}$$

Applying the star map to these equations yields (10). In this sense, the iteration of the IFS (10) in internal space corresponds to the iteration (11) in physical space (note that by Proposition 1 the star map is bijective on $\mathbb{Z}[\alpha]$).

Setting $\Omega_{AB} = \Omega_A \cup \Omega_B$ in (10), the system decouples and we remain with the simpler IFS

$$(12) \quad \Omega_{AB} = f_1(\Omega_{AB}) \cup f_2(\Omega_{AB}) \cup f_3(\Omega_{AB})$$

where

$$(13) \quad f_1(z) = \beta z, \quad f_2(z) = \beta^3 z + \beta^3 \quad \text{and} \quad f_3(z) = \beta z + \beta^2 - \beta.$$

The mappings $f_i: \mathbb{C} \mapsto \mathbb{C}$ are contractions ($|\beta| < 1$), so that Hutchinson’s theorem [20, Section 3.1(3)] guarantees a unique compact solution of (12), called the *attractor* of the IFS. The sets Ω_A, Ω_B and Ω_C can be calculated from Ω_{AB} as

$$(14) \quad \Omega_A = f_1(\Omega_{AB}), \quad \Omega_B = f_2(\Omega_{AB}) \cup f_3(\Omega_{AB}), \quad \text{and} \quad \Omega_C = f_4(\Omega_{AB}),$$


where $f_4(z) = \beta^2 z + \beta^2$. They are also compact sets in the plane. For the components of Ω_{AB} , see Figure 1; the windows Ω_A , etc., are shown in Figure 6. Note that the decoupling of the IFS (i.e., the step from (10) to (12)) lies at the heart of our argument and seems to be the reason that we cannot immediately generalize our method to other unimodular substitutions of Pisot type⁵, because no such decoupling emerges in general.

The *similarity dimension* s of a set given by an IFS is the unique non-negative number s such that the contraction constants to the power of s add up to 1 (see [16]). For Ω_{AB} , this means

$$|\beta|^{3s} + 2|\beta|^s = 1$$

with solution $s = s(\Omega_{AB}) = 2$ (because $\alpha \cdot |\beta|^2 = 1$, the substitution is unimodular). The similarity dimension s of a set is connected to its *Hausdorff dimension* h by $h \leq s$ where equality holds if the *open set condition* (OSC for short) is satisfied [16]. An IFS with mappings f_i satisfies the OSC iff there exists a nonempty open set U such that $f_i(U) \cap f_j(U) = \emptyset$ for $i \neq j$ and $f_i(U) \subset U$ for all i . It is easy to see that the corresponding self-similar set Ω_{AB} must be contained in the closure \overline{U} so that the pieces $f_i(\Omega_{AB}) \subset \overline{f_i(U)}$ can intersect at their boundaries but cannot have interior points in common [8]. If their boundaries do intersect, the IFS is called *just touching*.

⁵The substitution (2) can be analyzed by the balanced pair algorithm as described in [34]. This algorithm also confirms that it has pure point spectrum, but one does not get the model set property.



Point	Coordinate
P_1	$\frac{1}{2}(\beta^2 - 3\beta - 1)$
P_2	$\frac{1}{2}(\beta^2 - 3\beta + 1)$
P_3	$\frac{1}{2}(-\beta^2 + \beta + 1)$
P_4	$\frac{1}{2}(-\beta^2 + \beta - 1)$
P_5	$-\frac{1}{2}\beta$
P_6	$\frac{1}{2}(\beta^2 - \beta - 1)$
P_7	$\frac{1}{2}(-\beta^2 - \beta - 1)$
P_8	$\frac{1}{2}(\beta^2 - \beta + 1)$
P_9	$\frac{1}{2}(-\beta^2 - \beta + 1)$
P_{10}	$\frac{1}{2}(-\beta + 1)$

Figure 1: Components $f_i(\Omega_{AB})$ of the set Ω_{AB} and the points P_1, \dots, P_{10} used in the proof of Proposition 2. Ω_{AB} is the union of the three shaded areas.

Table 1: Coordinates of the points P_1, \dots, P_{10} in Figure 1.

Proposition 2 *The IFS of (12) for Ω_{AB} is just touching.*

Proof To determine the boundary of Ω_{AB} , we choose special points P_i in Ω_{AB} , see Figure 1 (for illustration) and Table 1 (for details)⁶. We first show how these points are determined. Demanding

$$P_2 = f_3(P_1), \quad P_3 = f_1(P_2), \quad P_4 = f_1(P_3), \quad \text{and} \quad P_1 = f_3(P_4)$$

one gets the following fixed point equation

$$(15) \quad P_1 = f_3 \circ f_1 \circ f_1 \circ f_3(P_1) = \beta^4 P_1 + 6\beta^2 + 2$$

for P_1 , and similar results hold for P_2, P_3 and P_4 . The unique solution of (15) is

$$P_1 = \frac{1}{2}(\beta^2 - 3\beta - 1).$$

Choosing

$$P_5 = \frac{1}{2}(P_1 + P_3), \quad P_6 = f_3(P_3), \quad P_7 = f_1(P_4),$$

$$P_8 = f_2(P_3), \quad P_9 = f_1(P_1), \quad P_{10} = \frac{1}{2}(P_2 + P_3)$$

⁶We use the two dimensional geometry of the internal space here explicitly, and, instead of going into cumbersome notations and explanations, show some figures to clarify and assist the proofs.

and setting τ to be the inversion in the center P_5 ($\tau: z \mapsto -z - \beta$) and κ the one in the center P_{10} ($\kappa: z \mapsto -z - \beta + 1$), one can verify the following equations:

$$\begin{aligned} P_1 &= \tau(P_3) \\ P_2 &= \tau(P_4) = \kappa(P_3) \\ P_5 &= \frac{1}{2}(P_2 + P_4) = \frac{1}{2}(P_6 + P_9) = \frac{1}{2}(P_7 + P_8) \\ P_6 &= \tau(P_9) \\ P_7 &= f_2(P_1) = f_3(P_9) = \tau(P_8) \\ P_8 &= f_1(P_6) = f_3(P_2) = \tau(P_8) = \kappa(P_9) \\ P_{10} &= \frac{1}{2}(P_8 + P_9). \end{aligned}$$

For the mappings, one finds

$$f_1 \circ \tau = \tau \circ f_3, \quad f_2 \circ \tau = \tau \circ f_2 \quad \text{and} \quad f_3 \circ \tau = \tau \circ f_1,$$

showing that Ω_{AB} is inversion symmetric in the center P_5 , i.e., $\tau(\Omega_{AB}) = \Omega_{AB}$.

Denoting by $[P_2, P_3]$ the “boundary” between P_2 and P_3 (the “right edge”), one finds

$$\begin{aligned} [P_2, P_8] &= f_3 \circ \tau \circ f_1([P_2, P_3]) \\ [P_8, P_9] &= f_1 \circ \tau \circ f_1 \circ \tau \circ f_1([P_2, P_3]) \\ [P_9, P_3] &= f_1 \circ \tau \circ f_1([P_2, P_3]) \end{aligned}$$

and therefore the following IFS for $[P_2, P_3]$:

$$(16) \quad [P_2, P_3] = g_1([P_2, P_3]) \cup g_2([P_2, P_3]) \cup g_3([P_2, P_3]),$$

where

$$(17) \quad g_1(z) = -\beta^2 z - \beta, \quad g_2(z) = (2\beta^2 + 1)z + \beta^2 + 1, \quad \text{and} \quad g_3(z) = -\beta^2 z - \beta^2.$$

Of course, we have not shown yet that $[P_2, P_3]$ really is (a piece of) the boundary of Ω_{AB} , so we just define $[P_2, P_3]$ to be the unique compact solution of the IFS (16), which is inversion symmetric in the center P_{10} because

$$g_1 \circ \kappa = \kappa \circ g_3, \quad g_2 \circ \kappa = \kappa \circ g_2 \quad \text{and} \quad g_3 \circ \kappa = \kappa \circ g_1.$$

Also, we know that $[P_2, P_3]$ is connected since we can start the iteration with the straight line from P_2 to P_3 . In each iteration, the image remains a (piecewise smooth) path from P_2 to P_3 .

With the mappings f_1 and τ , we get a boundary

$$\begin{aligned} [P_2, P_3] \cup f_1([P_2, P_3]) \cup \tau([P_2, P_3]) \cup \tau \circ f_1([P_2, P_3]) \\ = [P_2, P_3] \cup [P_3, P_4] \cup [P_4, P_1] \cup [P_1, P_2] \end{aligned}$$

around a simply connected open set U (we will prove in the next proposition that this boundary is non-self-intersecting). Now one can show that only the boundaries of $\overline{f_i(U)}$ intersect. Consider, for example, the region between $f_2(U)$ and $f_3(U)$. Then the boundary $[P_7, P_8]$ on $f_2(U)$ is given by

$$[P_7, P_8] = f_2 \circ \tau \circ f_1([P_2, P_3]) \cup f_2([P_2, P_3]),$$

while the one on $f_3(U)$ is given by (taking orientation into account)

$$[P_7, P_8] = f_3 \circ \kappa \circ g_2([P_2, P_3]) \cup f_3 \circ g_3([P_2, P_3]).$$

It is easy to verify that

$$f_2 \circ \tau \circ f_1 = f_3 \circ \kappa \circ g_2 \quad \text{and} \quad f_2 = f_3 \circ \kappa \circ g_3.$$

So the boundaries coincide. Similarly, one can check the region between $f_1(U)$ and $f_2(U)$ (note that P_7 and P_8 belong to all three sets $\overline{f_i(U)}$), and that the boundary of U coincides with pieces of the boundaries of the $f_i(U)$ —so the situation is as expected from Figure 1. Therefore, the IFS is just touching. Also, we now know that $[P_2, P_3]$ is really a piece of the boundary of Ω_{AB} . ■

Proposition 3 *Let Ω_{AB} be the unique compact solution of the IFS (12). Then its boundary is non-self-intersecting.*

Proof We specify a (closed) rhombus R (which surrounds the boundary $[P_2, P_3]$ and the straight line from P_2 to P_3) such that its iteration in (12) will not leave R , i.e., such that

$$(18) \quad (g_1(R) \cup g_2(R) \cup g_3(R)) \cap (\mathbb{R}^2 \setminus R) = \emptyset.$$

Furthermore, we also require that

$$(19) \quad g_1(R) \cap g_3(R) = \emptyset$$

and

$$(20) \quad \begin{aligned} g_1(R) \cap g_2(g_i(R)) &= \emptyset \quad \text{for } i \neq 1 \\ g_2(R) \cap g_1(g_i(R)) &= \emptyset \quad \text{for } i \neq 3 \\ g_2(R) \cap g_3(g_i(R)) &= \emptyset \quad \text{for } i \neq 1 \\ g_3(R) \cap g_2(g_i(R)) &= \emptyset \quad \text{for } i \neq 3. \end{aligned}$$




Figure 2: A rhombus R with corners E_1, \dots, E_4 , such that $g_i(R) \subset R$ for all i (see (18)). The $g_i(R)$ are shaded in gray. For the coordinates of E_1, \dots, E_4 , see Table 2.




Figure 3: In addition to Figure 2, the second iteration of the rhombus R is also shown, shaded in the dark gray. (20) is satisfied.




Figure 4: The situation around the “joint” P_8 . We have an inversion symmetry in the center P_{14} . For the coordinates, see Table 2.

Such a rhombus R exists, see Figure 2 for a picture of such a rhombus that satisfies the conditions of (18) and (19) and Table 2 for the coordinates of its corners (of course, we could also use a shape different from a rhombus). This rhombus also satisfies (20), see Figure 3.

Here, (19) tells us that $[P_2, P_8]$ and $[P_9, P_3]$ do not have a point in common; similar statements apply for (20). Each iterate of the rhombus is associated to a corresponding iterate of the boundary $[P_2, P_3]$ or the straight line from P_2 to P_3 (denoted by (P_2, P_3)). We call two rhombi at the same iteration level neighbouring if their corresponding iteration of (P_2, P_3) have a common endpoint. We see in Figure 3 that only neighbouring rhombi intersect at the second iteration level. We show that for any iteration level only neighbouring rhombi intersect.

We have verified the assertion for the first and second iteration level and proceed inductively. Since g_1, g_2 and g_3 are affine, we get the third iteration level as follows: The associated rhombi between P_2 and P_8 are a scaled down (by g_1) version of those of the second level, therefore the assertion holds for them. Similarly for the rhombi between P_8 and P_9 (by g_2) and between P_9 and P_3 (by g_3). So the only critical points remaining are the “joints” at P_8 and P_9 . We show that, at these points, also only neighbouring rhombi intersect and for this, we make use of the self-similar structure of the boundary, see Figure 4: The boundary $[P_{12}, P_{11}]$ is inversion symmetric in

the center P_{14} . This is clear for $[P_{13}, P_8] = g_1(g_3([P_2, P_3]))$ (and therefore $P_{14} = g_1(g_3(P_{10}))$). But it also holds for $[P_{12}, P_{13}] = g_1(g_2([P_2, P_3]))$ and $[P_8, P_{11}] = g_2(g_1(\kappa([P_2, P_3])))$. So, since the assertion holds around P_{13} , it also holds around P_8 by symmetry. Similar arguments apply around P_9 . So the assertion holds for the third iteration level, *i.e.*, for the third iteration level only neighbouring rhombi intersect. But the same argument applies to all further iteration levels. So the assertion is true, *i.e.*, for a given iteration level only neighbouring rhombi intersect. Also note that each rhombus has two neighbouring rhombi (with the exception of the “starting” and “ending” rhombi at P_2 and P_3 which only have one) and that there is no “rhombus loop”, *i.e.*, going from P_2 to P_3 we cross each rhombus only once.

Now, suppose $[P_2, P_3]$ is self-intersecting. Then there exist points $x, y \in [P_2, P_3]$ ($x \neq y$) such that they are connected in $[P_2, P_3]$ in two different ways, W_1 and W_2 (and we have a loop). We can choose points $u \in W_1$ and $v \in W_2$ such that $d(u, W_2) = \min\{d(v, z) \mid z \in W_2\} > 0$ ($W_2 \subset [P_2, P_3]$ is compact) and $d(v, W_1) > 0$. But then, u, v are in non-neighbouring rhombi for some iteration level N (and then for all iteration levels $n \geq N$), since the length of a rhombus of the N -th iteration level is at most $|\beta|^{2N} \cdot |E_2 - E_4|$. So, we get a “rhombus loop” for this iteration level by the rhombi which overlay W_1 and W_2 . This is a contradiction, therefore $[P_2, P_3]$ is non-self-intersecting.

From this single edge we proceed to all of the boundary. Here, critical are the “joints” $P_2, P_3, \text{etc.}$, again, because we get the other three parts by an affine map of this edge (*e.g.*, $[P_3, P_4] = f_1([P_2, P_3])$) and opposite edges (*i.e.*, $[P_2, P_3]$ and $[P_4, P_1]$) do not overlap, *cf.* Figure 5. But at P_2 , an argument like the one at P_8 above applies, *i.e.*, we have an inversion symmetry of part of the boundary in the center $\frac{1}{2}(P_{12} + P_{13})$ (and similar for the other “joints”). This extends our findings to the entire boundary. ■

This also implies that the boundaries of Ω_A, Ω_B and Ω_C , respectively their union Ω , are non-self-intersecting. Also, from the proof of the last proposition, we can deduce the following.

Corollary 1 *The point 0 is an inner point of $f_1(\Omega_{AB}) \subset \Omega_A$ and $-\beta$ is an inner point of $f_3(\Omega_{AB}) \subset \Omega_B$.*

Proof We again use the iteration of rhombi as in Proposition 3 to show that the two points are really inner points in the respective areas. For this, see Figure 5, where the first iteration of the rhombi is used for all parts of the boundary. Clearly, the points $0, -\beta$ are inner points, which can easily be checked by a simple (though somewhat tedious) calculation of distances. ■

Proposition 4 *Let Ω_{AB} be the unique compact solution of the IFS (12).*

- (i) Ω_{AB} is inversion symmetric in the center $P_5 = -\frac{1}{2}\beta$.
- (ii) Ω_{AB} has Hausdorff dimension $h(\Omega_{AB}) = 2$.
- (iii) Ω_{AB} has positive (Hausdorff and Lebesgue) measure (area).

Point	Coordinate
E_1	$P_{10} - i\frac{2}{5}(\beta^2 - 2\beta)$
E_2	$P_2 + \frac{2}{5}(\beta^2 - 2\beta)$
E_3	$P_{10} + i\frac{2}{5}(\beta^2 - 2\beta)$
E_4	$P_3 - \frac{2}{5}(\beta^2 - 2\beta)$
P_8	$\frac{1}{2}(\beta^2 - \beta + 1)$
P_{11}	$\frac{1}{2}(9\beta^2 + \beta + 5)$
P_{12}	$\frac{1}{2}(-3\beta^2 - 3\beta - 1)$
P_{13}	$\frac{1}{2}(5\beta^2 - \beta + 3)$
P_{14}	$\frac{1}{2}(3\beta^2 - \beta + 2)$

Table 2: Coordinates of the points used in Figures 2 and 4.




Figure 5: The points 0 and $-\beta$ are represented as black dots. Also, the boundary is shown and the area of the rhombi (first iteration for all parts of the boundary) is shaded in gray. The boundary runs inside this shaded region, so 0 and $-\beta$ are inner points.

- (iv) The boundary $\partial\Omega_{AB}$ has vanishing (Lebesgue) measure.
- (v) There is a periodic tiling of the plane with Ω_{AB} as prototile.

Proof (i) See proof of Proposition 2.

(ii) Just touching implies the OSC, therefore $h(\Omega_{AB}) = s(\Omega_{AB}) = 2$.

(iii) The OSC for Ω_{AB} (or any self-similar set with similarity dimension s) is equivalent to the positive Hausdorff measure condition $\mu^s(\Omega_{AB}) > 0$, where μ^s denotes the s -dimensional Hausdorff measure, see [8] and references therein. For Euclidean dimensions, Hausdorff and Lebesgue measure are connected by a nonzero multiplicative constant.

(iv) The similarity dimension $\bar{s} = s(\partial\Omega_{AB})$ of the boundary is the solution of (contraction constants given in (17))

$$2|\beta|^{2\bar{s}} + |\beta|^{3\bar{s}} = 1,$$

which is $\bar{s} = -\log(\tau)/\log(|\beta|) \approx 1.22$ (where $\tau = \frac{1}{2}(1 + \sqrt{5})$ is the golden ratio; the previous equation is solved by $|\beta|^{-\bar{s}} = \tau$). Therefore, the statement follows from $h(\partial\Omega_{AB}) \leq s(\partial\Omega_{AB})$.

(v) Because of the inversion symmetries τ of Ω_{AB} and κ of $\partial\Omega_{AB}$ from P_2 to P_3 (see proof of Proposition 2), the “right edge” and the “left edge” differ only by a translation $P_2 - P_1 = 1$, and, similarly, the “upper edge” and the “lower edge” differ by $P_2 - P_3 = \beta^2 - 2\beta$. ■

Proposition 5 *Let $\Omega_A, \Omega_B, \Omega_C$ be the solution of the IFS (10), and $\Omega = \Omega_A \cup \Omega_B \cup \Omega_C$. Then Ω is a compact set, homeomorphic to a disc, with positive area. The boundary $\partial\Omega$ is a fractal of vanishing Lebesgue measure, which is non-self-intersecting. The set Ω admits a lattice tiling of \mathbb{R}^2 , where the lattice is spanned by $P_2 - P_6 = -\beta + 1$ and $P_2 - P_3 = \beta^2 - 2\beta$.*

Proof It is clear from our construction that Ω is a compact set with simply connected interior. We have also seen that the boundary is connected and consists of finitely many pieces, each of which is obtained from a construction as used in the proof of Proposition 2. So, Ω must be homeomorphic to a disc. The remaining statements follow directly from Propositions 3 and 4, because the mappings in (14) are affine and the just touching property also holds for $\Omega = \Omega_A \cup \Omega_B \cup \Omega_C$. Since we also know the boundary of Ω (we have an IFS for every part of it), we can also verify the translation vectors by comparing the corresponding iterated function systems. Also, see Figure 6 for a depiction of these vectors. ■

Corollary 2 *$\Lambda(\Omega)$ is a regular model set.* ■

We can calculate the volume⁷

$$(21) \quad |\Gamma| = |\det(\mathbf{v}_A, \mathbf{v}_B, \mathbf{v}_C)| = |\operatorname{Im}(\beta)|(3\alpha^2 - 4\alpha) = \frac{1}{2}\sqrt{59} \approx 3.84$$

of the fundamental domain of Γ . And because of the periodic tiling of the plane with Ω as a prototile, it is also easy to calculate the area $\mu_{\text{int}}(\Omega)$ of Ω : as $\mu_{\text{int}}(\partial\Omega) = 0$, $\mu_{\text{int}}(\Omega)$ equals the area of a fundamental domain of the corresponding lattice of periods. This gives

$$(22) \quad \begin{aligned} \mu_{\text{int}}(\Omega) &= |\det((P_2 - P_6), (P_2 - P_3))| = |\operatorname{Im}(\beta)|(\alpha^2 - \alpha) \\ &= \frac{1}{2} \frac{1}{\sqrt{59}}(3\alpha^2 - 2\alpha + 17) \approx 1.77. \end{aligned}$$

Then the following lemma applies.

Lemma 1 *Let Γ be a lattice in $\mathbb{R} \times \mathbb{R}^m$, $|\Gamma|$ be the volume of a measurable fundamental domain of Γ in $\mathbb{R} \times \mathbb{R}^m$ with respect to the product $\mu \otimes \mu_{\text{int}}$ of the Lebesgue measures μ, μ_{int} on \mathbb{R}, \mathbb{R}^m , respectively. Assume that we have a cut-and-project scheme as in (7). If Ω is a bounded subset of \mathbb{R}^m with almost no boundary, then the density $\operatorname{dens}(\Lambda(\Omega))$ of the corresponding regular model set in \mathbb{R} is*

$$\operatorname{dens}(\Lambda(\Omega)) = \frac{\mu_{\text{int}}(\Omega)}{|\Gamma|}.$$

⁷Note that the discriminant of $\mathbb{Q}(\alpha)$ is -59 . The volume $|\Gamma|$ is proportional to the square root of the absolute value of the discriminant. The proportional constant is one factor of $\frac{1}{2}$ because there is one complex conjugate pair $\beta, \bar{\beta}$ of algebraic conjugates of α , see [11, Chapter II, Section 4.2, Theorem 2]. Note that we also have a formula for $|\operatorname{Im}(\beta)|$ in terms of α by this:

$$|\operatorname{Im}(\beta)| = \frac{1}{2 \cdot \sqrt{59}}(-8\alpha^2 + 25\alpha - 6).$$




Figure 6: Components of the set Ω , the periodic tiling of the plane (\mathbb{C}) with it and the corresponding translation vectors.

Proof This follows from [28, Proposition 2.1] because the projection π is one-to-one on Γ by construction. ■

With (6), (21) and (22), it is now easy to check that the density of the model set $\Lambda(\Omega)$ and the density of $\Sigma \text{Kol}(3, 1)$ are equal, i.e.,

$$(23) \quad \frac{\mu_{\text{int}}(\Omega)}{|\Gamma|} = \frac{1}{\ell}.$$

Proposition 6 *The sequence $\Sigma \text{Kol}(3, 1)$ is a subset of $\Lambda(\Omega)$. Further, they differ at most on positions of zero density and therefore have the same pure point diffraction spectrum.*

Proof We choose $\mathbf{0}, -\mathbf{v}_B \in \Gamma$. Then their projections into internal space are elements of the attractor Ω , because $f_1(0) = 0$ and $f_3(-\beta) = -\beta$. But starting with these two points, the iteration of the IFS in internal space just corresponds to the iteration in (3), respectively (11), in physical space. Therefore, $\Sigma \text{Kol}(3, 1) \subset \Lambda(\Omega)$, because all iterates of the star map of 0 and $-\alpha$ (i.e., 0 and $-\beta$) stay in Ω . Equation (23) shows that both sequences have the same density. So they can at most differ on positions of zero density.

Regular model sets have a pure point diffraction spectrum, see [9, 6, 29] and references therein. Therefore, the diffraction spectrum of $\Lambda(\Omega)$ is pure point, and $\Sigma \text{Kol}(3, 1)$, differing at most at positions of zero density, has the same spectrum by an argument in [18]. ■

Theorem 1 $\Sigma \text{Kol}(3, 1)$ is a regular model set (except possibly for positions of zero density) and has a pure point diffraction spectrum. Its autocorrelation is a norm almost periodic point measure, supported on a uniformly discrete subset of $\mathbb{Z}[\alpha]$.

Proof The first assertion follows from Proposition 6. The autocorrelation measure (see [18] for details) is supported on $\Lambda(\Omega) - \Lambda(\Omega) \subset \mathbb{Z}[\alpha]$. Since $\Lambda(\Omega)$ is a model set, it is also a Meyer set, hence $\Lambda(\Omega) - \Lambda(\Omega)$ is Delone. The norm almost periodicity follows from [6, Theorem 5]. ■

What we have proved so far is enough to calculate the diffraction spectrum of $\Sigma \text{Kol}(3, 1)$ and $\text{Kol}(3, 1)$, see Section 5. But in the next section, we want to show that $\Lambda(\Omega)$ really equals $\Sigma \text{Kol}(3, 1)$.

Remarks For unimodular substitutions of Pisot type, i.e., $\text{Kol}(2m \pm 1, 2m \mp 1)$ with $m \geq 1$, the procedure is essentially the same. Unfortunately, the IFS does not decouple like in (12) for $m > 1$, which makes it technically more involved. For non-unimodular substitutions of Pisot type, the internal space is more complicated in having additional p -adic type components, see [30, 17, 7, 22, 23] for further details and examples.

The sequences $\Sigma \text{Kol}(p, q)$, which are not of Pisot type, do not have a pure point spectral component outside $k = 0$ by an argument in [10], see [31] for details.

Some of the results given have been studied extensively under the name of “Rauzy fractal”, e.g., that the windows have non-empty interior, that the windows do not overlap in this case (this follows from the so-called strong coincidence condition) and also the periodic tilability seems to follow from results in [1, 12, 35]. But we also need the lattice of the periodic tiling explicitly, as well as the induced IFS (16) for the boundary. Therefore, we opted to give an elementary and complete derivation here.

4 $\Sigma \text{Kol}(3, 1)$ is a Generic Model Set

We set $\overset{\circ}{\Omega} = \overset{\circ}{\Omega}_A \cup \overset{\circ}{\Omega}_B \cup \overset{\circ}{\Omega}_C \subset \Omega$ and $\partial\overset{\circ}{\Omega} = \Omega \setminus \overset{\circ}{\Omega} = \partial\Omega_A \cup \partial\Omega_B \cup \partial\Omega_C$. Then we can improve a statement of Proposition 6.

Proposition 7 $\Sigma \text{Kol}(3, 1)$ is equal to the model set $\Lambda(\overset{\circ}{\Omega})$.

Proof $\Sigma \text{Kol}(3, 1) \subset \Lambda(\overset{\circ}{\Omega})$: Note that the mappings f_i ($i \in \{0, 1, 3\}$) of (10') are similarities (all directions are contracted by the same factor, here $|\beta|$). Therefore, they map balls around x to balls around $f_i(x)$. Furthermore, they map balls in $\overset{\circ}{\Omega}_k$ to balls in $\overset{\circ}{\Omega}_\ell$ ($k, \ell \in \{A, B, C\}$). Since the starting points of the iteration (3) in internal space, namely $-\beta$ and 0 , are inner points of Ω_B and Ω_A by Corollary 1, one can also find balls of radius $\varepsilon > 0$ around $-\beta$ and 0 which lie entirely in $\overset{\circ}{\Omega}_B$ and $\overset{\circ}{\Omega}_A$, respectively. Since the iteration in physical space corresponds to the IFS in internal space, the star map of an arbitrary point in $\Sigma \text{Kol}(3, 1)$ is thus a point of $\overset{\circ}{\Omega}$.

$\Lambda(\hat{\Omega}) \subset \Sigma \text{Kol}(3, 1)$: Suppose $x \in \Lambda(\hat{\Omega}) \setminus \Sigma \text{Kol}(3, 1)$. Then x^* and all its iterates of the IFS (10) are in $\hat{\Omega}$, by the same reasoning as before. Furthermore, the mappings f_i ($i \in \{0, 1, 3\}$) are affine similarities and therefore all iterates of x^* are disjoint to all of the iterates of $-\beta$ and 0. But then

$$\text{dens } \Sigma \text{Kol}(3, 1) < \text{dens } \Lambda(\hat{\Omega}),$$

because the set of iterates of x under inflation has positive density in $\Lambda(\hat{\Omega})$. This contradicts Proposition 6. ■

Note that not only the original sequence with 3's and 1's is inversion symmetric (see (4)), but also the positions in $\Lambda(\Omega_{AB})$.

Corollary 3 $\Lambda(\Omega_{AB})$ is inversion symmetric in the center $-\frac{1}{2}\alpha$.

Proof The starting points $-\beta$ and 0 and the IFS (12) are inversion symmetric in the center P_5 . This corresponds, in physical space (by Galois conjugation), to inversion symmetry in the center $-\frac{1}{2}\alpha$. ■

For the following, we need some more definitions, see [3] and [26]. If Λ is a discrete point set in \mathbb{R}^d , we call $\mathcal{P}_r(u)$ an r -patch of a point $u \in \Lambda$, if $\mathcal{P}_r(u) = \Lambda \cap B_r(u)$, where $B_r(u)$ is the ball of radius r about u . Often, we are only interested in the set of $\{\mathcal{P}_r(u) \mid r > 0, u \in \Lambda\}$ and an element of this set is simply called a patch. Two structures Λ_1 and Λ_2 are locally indistinguishable (or locally isomorphic or LI) if each patch of Λ_1 is, up to translation, also a patch of Λ_2 and vice versa. The corresponding equivalence class is called LI-class.

A discrete structure Λ is repetitive, if for every $r > 0$ there is a radius $R(r) > 0$ such that within each ball of radius $R(r)$, no matter its position in \mathbb{R}^d , there is at least one translate of each r -patch. Note that every primitive substitution generates a repetitive sequence, wherefore $\Sigma \text{Kol}(3, 1)$ is repetitive.

We now look at generic model sets and show that $\Lambda(\Omega)$ is actually generic.

Lemma 2 The set $C = \{c \in \mathbb{R}^2 \mid (c + \partial\hat{\Omega}) \cap L^* = \emptyset\}$ is dense in \mathbb{R}^2 , especially $c + L^* \subset C$ for $c \in C$.

Proof The set C is not the empty set by Baire's category theorem ($\partial\hat{\Omega}$ is a meager set, L^* is countable), by standard arguments, which in this context first appeared in [27, Section 2.2.2], also see [7]. But if $c \in C$, then $c + t \in C, \forall t \in L^*$ (notice that L^* is an Abelian group), and L^* is dense. ■

Proposition 8

- (i) The model set $\Lambda(c + \Omega)$ is repetitive and generic for $c \in C$.
- (ii) The model sets $\Lambda(c + \Omega)$ and $\Lambda(\bar{c} + \Omega)$ are LI for $c, \bar{c} \in C$.

(iii) $\Sigma \text{Kol}(3, 1)$ and $\Lambda(c + \Omega)$ are LI for $c \in C$.

Proof (i) The model set $\Lambda(c + \Omega)$ is generic by the definition of the set C . It is repetitive by [28, Theorem 6] and [29, Proposition 3.1].

(ii) This is a by now standard argument, apparently first used in [27, Lemma 2.1].

(iii) Since $\Lambda(c + \Omega)$ is repetitive, by [27, Lemma 1.2] it is enough to check that every patch of $\Sigma \text{Kol}(3, 1)$ also occurs⁸ as a patch of $\Lambda(c + \Omega)$: Let \mathcal{P} be a patch of $\Sigma \text{Kol}(3, 1)$.

Then $\mathcal{P}^* \subset \overset{\circ}{\Omega}$ by Proposition 7, and since \mathcal{P} is a finite patch, we even know that there is an $\varepsilon > 0$ such that $\mathcal{P}^* \subset \overset{\circ}{\Omega}_\varepsilon$, where $\overset{\circ}{\Omega}_\varepsilon = \{t \in \mathbb{R}^2 \mid t \in \overset{\circ}{\Omega}, \text{dist}(t, \partial\overset{\circ}{\Omega}) > \varepsilon\}$. The set C is dense by Lemma 2, therefore there is a $\tilde{c} \in C$ such that $\mathcal{P}^* \subset \tilde{c} + \Omega$. Then $\mathcal{P} \subset \Lambda(\tilde{c} + \Omega)$ and $\Sigma \text{Kol}(3, 1)$ and $\Lambda(\tilde{c} + \Omega)$ are LI. Since LI is an equivalence relation, it follows from (ii) that $\Sigma \text{Kol}(3, 1)$ and $\Lambda(c + \Omega)$ are LI for every $c \in C$. ■

Proposition 9 Define $C_0 = \{c \in C \mid 0 \in c + \Omega\}$. Then, for every $r > 0$ and $c \in C_0$, we get

$$B_r(0) \cap \Sigma \text{Kol}(3, 1) = B_r(0) \cap (\Lambda(c + \Omega) + t) = B_r(0) \cap \Lambda(c + t^* + \Omega)$$

for an appropriate $t \in \mathbb{Z}[\alpha]$. Additionally, it follows that $c + t^* \in C_0$.

Proof By Proposition 8(iii) we know that $\Lambda(c + \Omega)$ and $\Sigma \text{Kol}(3, 1)$ are LI, therefore the patch $B_r(0) \cap \Sigma \text{Kol}(3, 1)$ occurs somewhere in $\Lambda(c + \Omega)$. By the choice of C_0 , we can translate this patch in $\Lambda(c + \Omega)$ with $t \in \mathbb{Z}[\alpha]$ to the origin. In internal space, this is a translation t^* and by Lemma 2 we have $c + t^* \in C$. But $\Sigma \text{Kol}(3, 1)$ has a point at the origin, so we even have $c + t^* \in C_0$. ■

Theorem 2 $\Sigma \text{Kol}(3, 1)$ is a regular generic model set.

Proof Since $\Sigma \text{Kol}(3, 1)$ is the fixed point of a primitive substitution, it is repetitive, and the corresponding dynamical system is minimal, see [29, Proposition 3.1].

By Proposition 8(iii) the model set $\Lambda(c + \Omega)$ for $c \in C$ is in the LI-class of $\Sigma \text{Kol}(3, 1)$, hence the latter must be the limit of some sequence $(t_i)_i$ of translations of $\Lambda(c + \Omega)$, where the translates t_i can be restricted to elements of $\mathbb{Z}[\alpha]$ and $c + t_i \in C_0$ by Proposition 9.

However, if $\partial\Omega \cap L^* \neq \emptyset$, there is a point $\tilde{x} \in \partial\overset{\circ}{\Omega} \cap L^* \cap \overset{\circ}{\Omega}$ with $\tilde{x} \in \partial\Omega_A \cap \partial\Omega_B$ (so \tilde{x} lies on the common boundary of Ω_A and Ω_B). This is because by appropriate combinations of the mappings $g_1, g_2, g_3, \tau, \kappa, f_0, f_1, f_2, f_3, f_4$ and the translations by $P_2 - P_1, P_2 - P_3, P_2 - P_6$ of Section 3, which all map L^* onto L^* , we can “move” points on $\partial\overset{\circ}{\Omega}$ from every “edge” to every other “edge”⁹. We have $\varepsilon_0 = \text{dist}(\tilde{x}, \partial\overset{\circ}{\Omega} \setminus (\partial\Omega_A \cap \partial\Omega_B)) > 0$.

⁸That every patch of $\Lambda(c + \Omega)$ is also one of $\Sigma \text{Kol}(3, 1)$, follows then together with the repetitivity of $\Lambda(c + \Omega)$.

⁹We even get that, with one point $x \in \partial\overset{\circ}{\Omega} \cap L^*$, there is a dense set of points in $\partial\overset{\circ}{\Omega} \cap L^*$, because we can always “move” x to the edge $[P_2, P_3]$, apply the IFS (16) there and “move” this edge, with now dense points, to every other edge.

The inverse star image of this point must then be in any limit of sequences $\Lambda(t_i^* + c + \Omega)$ with $c + t_i^* \rightarrow 0$, but it is not in $\Sigma \text{Kol}(3, 1)$ —which is a contradiction. So no such point can exist and $\partial\Omega \cap L^* = \emptyset$.

Regularity was established in Theorem 1 together with Proposition 7. ■

This argument is rather general and applies in other situations as well. For a more elementary proof see the appendix.

Remark By Proposition 5 we know that $\bigcup_{t \in G} (t + \Omega) = \mathbb{R}^2$, where $G = \langle -\beta + 1, \beta^2 - 2\beta \rangle_{\mathbb{Z}}$ is a rank 2 free Abelian group (a 2-dimensional lattice) and by Theorem 2 that $L^* = \mathbb{Z}[\beta] \subset \dot{\bigcup}_{t \in G} (t + \overset{\circ}{\Omega})$, where $\dot{\cup}$ denotes disjoint union. In physical space, we get

$$L = \mathbb{Z}[\alpha] = \dot{\bigcup}_{s \in G'} (s + \Sigma \text{Kol}(3, 1)),$$

where

$$G' = \langle -\alpha + 1, \alpha^2 - 2\alpha \rangle_{\mathbb{Z}} = \langle \ell_C - \ell_B, \ell_A - \ell_B \rangle_{\mathbb{Z}},$$

i.e., $\mathbb{Z}[\alpha]$ is the disjoint union of translates of the regular, generic model set $\Sigma \text{Kol}(3, 1)$. The set of translations needed is a rank 2 subgroup, whose Galois dual in $\mathbb{Z}[\beta]$ is a lattice. But we can also write

$$L = \mathbb{Z}[\alpha] = \dot{\bigcup}_{r \in \Sigma \text{Kol}(3,1)} (r + G'),$$

so L/G' is a coset system with the structure of a model set. Now, let $\lambda(m)$ be the m -th element of $\Sigma \text{Kol}(3, 1)$ ($m \in \mathbb{Z}$). Then one can show that the induced group structure on this coset system is $(\lambda(m)) + (\lambda(n)) = (\lambda(m + n))$, i.e., it is the action of \mathbb{Z} on $\Sigma \text{Kol}(3, 1)$. This group structure lines up with the deformation in the next section.

5 Deformation and Diffraction

In the cut-and-project scheme (7) with $H = \mathbb{R}^m$, let $\varphi: \mathbb{R}^m \rightarrow \mathbb{R}^d$ be a continuous function with compact support (e.g., Ω). We call

$$\Lambda_\varphi(\Omega) = \{x + \varphi(x^*) \mid x \in \Lambda(\Omega)\}$$

a *deformed model set* if it is also a Delone set, see [9]. The model set $\Lambda(\Omega)$ can be seen as deformed model set where the associated function φ is trivial, i.e., $\varphi \equiv 0$. The diffraction spectrum of (deformed) model sets (where each of its points is represented by a normalized Dirac measure, say) can be calculated explicitly, see [9] for details. We write δ_k for the Dirac measure at k , i.e., $\delta_k(f) = f(k)$ for f continuous. Also, we need the dual of a lattice $\Gamma \subset \mathbb{R}^n$ defined as

$$\Gamma^* := \{y \in \mathbb{R}^n \mid x \cdot y \in \mathbb{Z}, \forall x \in \Gamma\},$$

with $x \cdot y$ denoting the Euclidean scalar product.

Proposition 10 ([9]) *Let $\Lambda_\varphi(\Omega)$ be a deformed model set in \mathbb{R}^d constructed with a regular model set $\Lambda(\Omega)$ and a continuous function φ of compact support. Then, the diffraction pattern of $\Lambda_\varphi(\Omega)$ is the positive pure point measure*

$$\hat{\gamma} = \sum_{k \in \pi(\Gamma^*)} |c_k(\Lambda_\varphi(\Omega))|^2 \delta_k,$$

where Γ^* is the dual lattice of Γ , δ_k is the Dirac measure at k and $c_k(\Lambda_\varphi(\Omega))$ is the Fourier-Bohr coefficient of $\Lambda_\varphi(\Omega)$ at k . This Fourier-Bohr coefficient exists and has the value

$$(24) \quad c_k(\Lambda_\varphi(\Omega)) = \begin{cases} \frac{1}{|\Gamma|} \int_{\Omega} e^{-2\pi i(k \cdot \varphi(y) - k^* \cdot y)} dy, & \text{if } (k, k^*) \in \Gamma^*, \\ 0, & \text{otherwise.} \end{cases} \quad \blacksquare$$

For a regular model set $\Lambda(\Omega)$ (where $\varphi \equiv 0$), the Fourier-Bohr coefficient is just given by the (inverse) Fourier transform of the characteristic function of the window Ω .

For $\Sigma \text{ Kol}(3, 1)$, we note that the support F of the spectrum is dense in \mathbb{R} since it is given by the \mathbb{Z} -span of the projection of the dual lattice vectors, *i.e.*,

$$(25) \quad F = \mathbb{Z} \cdot \pi(\mathbf{v}_A^*) + \mathbb{Z} \cdot \pi(\mathbf{v}_B^*) + \mathbb{Z} \cdot \pi(\mathbf{v}_C^*),$$

but $\pi(\mathbf{v}_B^*) = (\alpha - 1)\pi(\mathbf{v}_A^*)$ and therefore they are linearly independent over \mathbb{Q} .

To deform $\Sigma \text{ Kol}(3, 1)$ to $\text{Kol}(3, 1)$, we make the linear ansatz $\varphi(x^*) = ax_1^* + bx_2^*$, where x_i^* denotes the i -th Cartesian component of the vector $x^* \in \mathbb{R}^2$. With this φ , we now deform all bond lengths ℓ_i to the average bond length ℓ , *i.e.*, we have to solve the following linear system of equations ($i \in \{A, B, C\}$):

$$(\mathbf{v}_i)_1 + a(\mathbf{v}_i)_2 + b(\mathbf{v}_i)_3 = \ell.$$

This over-determined system is solved by

$$a = \ell - 1 = \frac{1}{2}(-\alpha^2 + \alpha + 5) \approx 1.17 \quad \text{and}$$

$$b = \frac{\text{Im}(\beta)}{59} \cdot (-\alpha^2 - 17\alpha + 31) = \frac{1}{2\sqrt{59}}(-413\alpha^2 + 885\alpha - 59) \approx -0.13.$$

Due to the linearity of φ (and the positivity of the bond lengths involved), this deformation does not alter the order of the points (*i.e.*, for $x, x' \in \Lambda(\Omega)$ with $x < x'$, we always have $x + \varphi(x^*) < x' + \varphi(x'^*)$). Note that the support F of the spectrum stays the same as in (25), only the Fourier-Bohr coefficients change.

The positions in $\Lambda_\varphi(\Omega_i)$ are now subsets of $\ell \cdot \mathbb{Z}$. To be more precise, we even have

$$\Lambda_\varphi(\Omega_A) \cup \Lambda_\varphi(\Omega_B) \cup \Lambda_\varphi(\Omega_C) = \ell \cdot \mathbb{Z}.$$

Because of this embedding into $\ell \cdot \mathbb{Z}$, the diffraction spectrum of each of the aperiodic sets $\Lambda_\varphi(\Omega_i)$ is $(\ell \cdot \mathbb{Z})^*$ -periodic [2], *i.e.*, it is periodic with period $1/\ell$ (note that

$\mathbb{Z}^* = \mathbb{Z}$; the diffraction spectrum of $\Sigma \text{Kol}(3, 1)$ is not periodic). This might not be obvious from (24) at first sight, but for $n \in \mathbb{Z}$ we have (note that $\pi(\mathbf{v}_i^*) = \rho_i/\ell$)

$$(26) \quad \frac{n}{\ell} = n\pi(\mathbf{v}_A^* + \mathbf{v}_B^* + \mathbf{v}_C^*) = \pi(n\mathbf{v}_A^* + n\mathbf{v}_B^* + n\mathbf{v}_C^*),$$

therefore, for every $k' = k + \frac{n}{\ell}$ with $(k, k^*) \in \Gamma^*$, there is also a k'^* with $(k', k'^*) \in \Gamma^*$ given by

$$k'^* = k^* + \pi_{\text{int}}(n\mathbf{v}_A^* + n\mathbf{v}_B^* + n\mathbf{v}_C^*).$$

But with the chosen φ we get

$$(27) \quad (k' - k) \cdot \varphi(y) - (k'^* - k^*) \cdot y = ny_1 \left[\frac{a}{\ell} - (\pi_{\text{int}}(\mathbf{v}_A^* + \mathbf{v}_B^* + \mathbf{v}_C^*))_1 \right] \\ + ny_2 \left[\frac{b}{\ell} - (\pi_{\text{int}}(\mathbf{v}_A^* + \mathbf{v}_B^* + \mathbf{v}_C^*))_2 \right] = 0$$

because each of the terms in square brackets vanishes. Therefore

$$c_k(\Lambda_\varphi(\Omega_i)) = c_{k'}(\Lambda_\varphi(\Omega_i))$$

holds, and the spectrum is periodic with period $1/\ell$.

To obtain the diffraction spectrum of $\text{Kol}(3, 1)$ from here, one only has to rescale the positions in $\Lambda_\varphi(\Omega)$ by a factor of $1/\ell$. To summarize:

Theorem 3 *The bi-infinite sequence $\text{Kol}(3, 1)$, represented with equal bond lengths, is a deformed model set and has a pure point diffraction spectrum. ■*

Remarks By the same method, we can also find a deformation $\tilde{\varphi}(x^*) = \tilde{a}x_1^* + \tilde{b}x_2^*$ such that we represent the letter ‘1’ of $\text{Kol}(3, 1)$ with an interval of length $\tilde{\ell}$ and the letter ‘3’ with one of length $3\tilde{\ell}$. For this, the letters A, B, C have bond lengths $6\tilde{\ell}, 4\tilde{\ell}, 2\tilde{\ell}$, respectively. For the parameters of the deformation (the average bond length must be ℓ again), we get

$$\tilde{\ell} = \frac{1}{4}(7\alpha^2 - 15\alpha + 1) \approx 0.49, \quad \tilde{a} = \frac{1}{2}(7\alpha^2 - 15\alpha - 1) \approx -0.016 \quad \text{and} \\ \tilde{b} = \frac{\text{Im}(\beta)}{59} \cdot (-179\alpha^2 + 379\alpha + 3) = \frac{1}{2\sqrt{59}^3}(1239\alpha^2 - 767\alpha - 4661) \approx -0.36.$$

Now (26) changes to

$$\frac{n}{\tilde{\ell}} = \pi(6n\mathbf{v}_A^* + 4n\mathbf{v}_B^* + 2n\mathbf{v}_C^*),$$

and with the same calculation as before one gets an equation which corresponds to (27), where the two terms in square brackets

$$\left[\frac{\tilde{a}}{\tilde{\ell}} - (\pi_{\text{int}}(6\mathbf{v}_A^* + 4\mathbf{v}_B^* + 2\mathbf{v}_C^*))_1 \right] \quad \text{and} \quad \left[\frac{\tilde{b}}{\tilde{\ell}} - (\pi_{\text{int}}(6\mathbf{v}_A^* + 4\mathbf{v}_B^* + 2\mathbf{v}_C^*))_2 \right]$$

also both vanish. Therefore, the spectrum is periodic with period $1/\tilde{\ell}$ as expected [2], since $\Lambda_{\tilde{\varphi}}(\Omega) \subsetneq \tilde{\ell} \cdot \mathbb{Z}$. This representation with integer bond lengths (after rescaling) has the advantage that the union of the three aperiodic sets $\Lambda_{\tilde{\varphi}}(\Omega_i)$ is still an aperiodic set. Clearly, it is also pure point diffractive.

$\text{Kol}(3, 1)$ in its natural setting with intervals of length 1, or of lengths 3 and 1, can be obtained as a deformation of the model set $\Sigma \text{Kol}(3, 1)$ derived above, where the intervals have incommensurate length. The basic theory of this is fully developed in [9, 19], but one can also understand, from a dynamical systems point of view, which deformations are stable in the sense that they do not change the spectral type of the dynamical spectrum (and hence of the diffraction spectrum, due to unique ergodicity), see [4].

Acknowledgments

It is a pleasure to thank Christoph Bandt for fractal advice, Robert V. Moody for helpful discussions and the German Research Council (DFG) for financial support. Also, we like to thank the referee for useful suggestions which led to an improvement of this article.

Appendix: An Alternative Proof of Theorem 2

By Proposition 9 we can choose a sequence $(c_i)_i$ with $c_i = c + t_i^* \in C_0$ ($t_i^* \in L^*$) such that $B_{r_i} \cap \Sigma \text{Kol}(3, 1) = B_{r_i} \cap \Lambda(c_i + \Omega)$ for every sequence $(r_i)_i$ with $r_i > i$. Also, this statement holds for every subsequence $(c_{i_j})_j$.

Now, assume $\partial\hat{\Omega} \cap L^* \neq \emptyset$. Then we have a point $\tilde{x} \in \partial\hat{\Omega} \cap L^* \cap \hat{\Omega}$ with $\tilde{x} \in \partial\Omega_A \cap \partial\Omega_B$. Set $\varepsilon_0 = \text{dist}(\tilde{x}, \partial\hat{\Omega} \setminus (\partial\Omega_A \cap \partial\Omega_B)) > 0$. Then a translation $y \neq 0$ of Ω with $y \in B_{\varepsilon_0}(0) \cap C$ has the following effect: $\tilde{x} \in y + \hat{\Omega}$, because by the definition of C , \tilde{x} cannot be on the boundary $y + \partial\hat{\Omega}$, and by the choice of ε_0 , it must either be in $y + \Omega_A$ or $y + \Omega_B$.

Now take a sequence $(c_i)_i$ as above. Clearly, this sequence must converge to 0. Therefore, there is an N such that $|c_i| < \varepsilon_0$ for all $i > N$. By choosing an appropriate subsequence $(c_{i_j})_j$ we get a sequence $(\tilde{c}_j)_j$ with $\tilde{c}_j = c_{i_j}$ such that \tilde{x} is always either in $\tilde{c}_j + \Omega_A$ or in $\tilde{c}_j + \Omega_B$. Also we have $B_{r_i} \cap \Sigma \text{Kol}(3, 1) = B_{r_i} \cap \Lambda(\tilde{c}_i + \Omega)$ for $r_i > i$. But both π and π_{int} are one-to-one. Therefore, the inverse of the star map of \tilde{x} must be a point of each $\Lambda(\tilde{c}_i + \Omega)$ and it also must be in $B_R(0)$ for some $R < \infty$. But by Proposition 7, it is not in $\Sigma \text{Kol}(3, 1)$. Therefore we get a contradiction and our assumption is wrong. So, $0 \in C$ and $\Sigma \text{Kol}(3, 1)$ is generic by Proposition 8(i). ■

References

- [1] P. Arnoux and S. Ito, *Pisot substitutions and Rauzy fractals*. Bull. Belg. Math. Soc. Simon Stevin **8**(2001), 181–207.
- [2] M. Baake, *Diffraction of weighted lattice subsets*. Canad. Math. Bull. **45**(2002), 483–498. math.MG/0106111

- [3] ———, *A guide to mathematical quasicrystals*. In: Quasicrystals, (eds., J.-B. Suck, M. Schreiber and P. Häussler), Springer, Berlin, 2002, 17–48. [math-ph/9901014](#)
- [4] M. Baake and D. Lenz, *Deformation of Delone dynamical systems and topological conjugacy*. Preprint (2003).
- [5] M. Baake and R. V. Moody, *Self-similar measures for quasicrystals*. In: Directions in Mathematical Quasicrystals, (eds., M. Baake and R. V. Moody), Amer. Math. Soc., Providence, 2000, 1–42. [math.MG/0008063](#)
- [6] ———, *Weighted Dirac combs with pure point diffraction*. To appear in: J. Reine Angew. Math. [math.MG/0203030](#)
- [7] M. Baake, R. V. Moody and M. Schlottmann, *Limit-(quasi)periodic point sets as quasicrystals with p -adic internal spaces*. J. Phys. A **31**(1998), 5755–5765. [math-ph/9901008](#)
- [8] C. Bandt, *Self-similar tilings and patterns described by mappings*. In: The Mathematics of Long-Range Aperiodic Order, (ed., R. V. Moody), Kluwer, Dordrecht (1997), 45–83.
- [9] G. Bernuau and M. Duneau, *Fourier Analysis of deformed model sets*. In: Directions in Mathematical Quasicrystals, (eds., M. Baake and R. V. Moody), AMS, Providence (2000), 43–60.
- [10] E. Bombieri and J. E. Taylor, *Which distributions of matter diffract? An initial investigation*. J. Physique Coll. C **3**(1986), 19–29.
- [11] S.I. Borewicz and I.R. Šafarevič, *Zahlentheorie*. Birkhäuser, Basel, 1966.
- [12] V. Canterini and A. Siegel, *Geometric representation of substitutions of Pisot type*. Trans. Amer. Math. Soc. **353**(2001), 5121–5144.
- [13] F. M. Dekking, *The spectrum of dynamical systems arising from substitutions of constant length*. Z. Wahrsch. Verw. Gebiete **41**(1978), 221–239.
- [14] ———, *Regularity and irregularity of sequences generated by automata*. Sémin. Th. Nombres Bordeaux 1979–80, exposé 9, 901–910.
- [15] ———, *What is the long range order in the Kolakoski sequence?* In: The Mathematics of Long-Range Aperiodic Order, (ed., R. V. Moody), Kluwer, Dordrecht, 1997, 115–125.
- [16] G. A. Edgar, *Measure, Topology and Fractal Geometry*. Springer, New York, 1990.
- [17] F. Gähler and R. Klitzing, *The diffraction pattern of self-similar tilings*. In: The Mathematics of Long-Range Aperiodic Order, (ed., R. V. Moody), Kluwer, Dordrecht, 1997, 141–174.
- [18] A. Hof, *On diffraction by aperiodic structures*. Comm. Math. Phys. **169**(1995), 25–43.
- [19] ———, *Diffraction by aperiodic structures*. In: The Mathematics of Long-Range Aperiodic Order, (ed., R. V. Moody), Kluwer, Dordrecht, 1997, 239–268.
- [20] J. E. Hutchinson, *Fractals and self-similarity*. Indiana Univ. Math. J. **30**(1981), 713–747.
- [21] W. Kolakoski, *Self generating runs*, Problem 5304. Amer. Math. Monthly **72**(1965), 674.
- [22] J.-Y. Lee and R. V. Moody, *Lattice substitution systems and model sets*. Discrete Comput. Geom. **25**(2001), 173–201. [math.MG/0002019](#)
- [23] J.-Y. Lee, R. V. Moody and B. Solomyak, *Pure point dynamical and diffraction spectra*. Ann. Inst. H. Poincaré **3**(2002), 1003–1018. [mp_arc/02-39](#)
- [24] J. M. Luck, C. Godrèche, A. Janner and T. Janssen, *The nature of the atomic surfaces of quasiperiodic self-similar structures*. J. Phys. A **26**(1993), 1951–1999.
- [25] R. V. Moody, *Meyer sets and their duals*. In: The Mathematics of Long-Range Aperiodic Order, (ed., R. V. Moody), Kluwer, Dordrecht, 1997, 403–441.
- [26] ———, *Model sets: a survey*. In: From Quasicrystals to More Complex Systems, (eds., F. Axel, F. Dénoyer and J.P. Gazeau), EDP Sciences, Les Ulis, and Springer, Berlin, 2000, 145–166. [math.MG/0002020](#)
- [27] M. Schlottmann, *Geometrische Eigenschaften quasiperiodischer Strukturen*. Dissertation, Universität Tübingen, 1993.
- [28] ———, *Cut-and-project sets in locally compact Abelian groups*. In: Quasicrystals and Discrete Geometry, (ed., J. Patera), Amer. Math. Soc., Providence, 1998, 247–264.
- [29] ———, *Generalized model sets and dynamical systems*. In: Directions in Mathematical Quasicrystals, (eds., M. Baake and R. V. Moody), Amer. Math. Soc., Providence, 2000, 43–60.
- [30] A. Siegel, *Représentation des systèmes dynamiques substitutifs non unimodulaires*. Ergodic Theory Dynam. Systems **23**(2003), 1247–1273.
- [31] B. Sing, *Spektrale Eigenschaften der Kolakoski-Sequenzen*. Diploma Thesis, Universität Tübingen, 2002, available from the author.
- [32] ———, *Kolakoski-(2m, 2n) are limit-periodic model sets*. J. Math. Phys. **44**(2003), 899–912. [math-ph/0207037](#)
- [33] V. F. Sirvent, *Modelos geométricos asociados a substituciones*. Habilitation (trabajo de ascenso), Universidad Simón Bolívar (1998); available at: <http://www.ma.usb.ve/~vsirvent/publi.html>.

- [34] V. F. Sirvent and B. Solomyak, *Pure discrete spectrum for one-dimensional substitutions of Pisot type*. *Canad. Math. Bull.* **45**(2002), 697–710.
- [35] V. F. Sirvent and Y. Wang, *Self-affine tiling via substitution dynamical systems and Rauzy fractals*. *Pacific J. Math.* **206**(2002), 465–485.

*Fakultät für Mathematik
Universität Bielefeld
Postfach 100131
33501 Bielefeld
Germany*

e-mail: mbaake@mathematik.uni-bielefeld.de

*Institut für Mathematik
Universität Greifswald
Jahnstr. 15a
17487 Greifswald
Germany*

e-mail: sing@uni-greifswald.de

<http://schubert.math-inf.uni-greifswald.de>