

Stellar Populations of Kinematically Decoupled Cores in E/S0 Galaxies†

Richard M. McDermid¹, Eric Emsellem², Kristen L. Shapiro³,
Roland Bacon², Martin Bureau⁴, Michele Cappellari^{1,4}, Roger L.
Davies⁴, Tim de Zeeuw¹, Jesús Falcón-Barroso^{1,5}, Davor Krajnović⁴,
Harald Kuntschner⁶ Reynier F. Peletier⁷, and Marc Sarzi⁸

¹Leiden Observatory, Leiden University, Postbus 9513, 2300 RA Leiden, The Netherlands
email: mcdermid@strw.leidenuniv.nl

²Université de Lyon 1, CRAL, Observatoire de Lyon, 9 av. Charles André, F-69230
Saint-Genis Laval; CNRS, UMR 5574 ; ENS de Lyon, France

³UC Berkeley Department of Astronomy, Berkeley, CA 94720, USA

⁴Denys Wilkinson Building, University of Oxford, Keble Road, Oxford, United Kingdom

⁵European Space and Technology Centre (ESTEC), Keplerlaan 1, Postbus 299, 2200 AG
Noordwijk, The Netherlands

⁶Space Telescope European Coordinating Facility, European Southern Observatory,
Karl-Schwarzschild-Str 2, 85748 Garching, Germany

⁷Kapteyn Astronomical Institute, Postbus 800, 9700 AV Groningen, The Netherlands

⁸Centre for Astrophysics Research, Science & Technology Research Institute, University of
Hertfordshire, Hatfield, United Kingdom

Abstract. In this poster contribution, we present results from high spatial resolution integral-field spectroscopy of elliptical (E) and lenticular (S0) galaxies from the SAURON representative survey, obtained with the OASIS and GMOS spectrographs. These seeing-limited observations explore the central $\sim 10'' \times 10''$ (typically one kiloparsec diameter) regions of these galaxies using a spatial sampling four times higher than SAURON ($0''.27$ vs. $0''.94$ spatial elements), resulting in almost a factor of two improvement in the median PSF. These data allow accurate study of the central regions to complement the large-scale view provided by SAURON. We derive the stellar and gas kinematics, stellar absorption-line strengths and nebular emission-line strengths for our sample, and derive maps of the luminosity-weighted stellar age, metallicity and abundance ratio via stellar population models. From these data we find a wealth of structures either not seen or poorly resolved in the SAURON data, including a number of kinematically-decoupled cores (KDCs) in the centres of some galaxies. We compare the intrinsic size and luminosity-weighted stellar age of all the visible KDCs in the full SAURON sample, and find two types of components: kiloparsec-scale KDCs, which are older than 8 Gyr, and are found in galaxies with little net rotation; and compact KDCs, which have intrinsic diameters of less than a few hundred parsec, show a range of stellar ages from 0.5 - 15 Gyr (with 5/6 younger than 5 Gyr), are found exclusively in fast-rotating galaxies, and are close to counter-rotating around the same axis as their host. Of the 7 galaxies in the SAURON sample with integrated luminosity-weighted ages less than 5 Gyr, 5 show such compact KDCs, suggesting a link between counter-rotation and recent star-formation. We show that this may be partly due to a combination of small sample size at young ages, and an observational bias, since young KDCs are easier to detect than their older and/or co-rotating counterparts.

Keywords. galaxies: elliptical and lenticular, cD; evolution; formation; kinematics and dynamics; stellar content; structure

† Poster available at: www.strw.leidenuniv.nl/~mcdermid/mcdermid_iau_2006_poster.pdf