

traumatic stress disorders in those recovering from Covid-19. The results confirmed the continuing effect of the program after the follow-up period.

Conclusions: Using metacognitive therapy has an effective effect in reducing post-traumatic stress disorder, and it can be used with many psychologically disturbed people.

Disclosure of Interest: None Declared

O0092

Prevalence of Prolonged Grief Disorder and Related Clinical Factors During the COVID-19 Pandemic in Turkey

R. Tekdemir^{1*}, A. Kandeger², Ö. Tan², M. Aydın² and K. Altınbaş³

¹Psychiatry, Selcuk University Faculty of Medicine, Mazhar Osman Mood Clinic; ²Psychiatry, Selcuk University Faculty of Medicine, Konya and ³Freelance, Consulting doctor, Istanbul, Türkiye

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.215

Introduction: Prolonged grief disorder has recently been added to the Diagnostic and Statistical Manual of Mental Disorders 5, Text Revision. To understand the health burden and then allocate economic and professional resources, it is necessary to provide epidemiological data for this new disorder. More information on the characteristics of people suffering from PGD is also beneficial to better identify individuals at risk.

Objectives: This study, therefore, aimed to estimate the prevalence of the PGD criteria in a special period such as the Covid-19 pandemic and in a representative population-based sample, evaluate the sociodemographic, and loss-related correlates of PGD case-ness and explore possible predictors.

Methods: The study included 126 people (97 females/29 males) who lost a relative for any reason during the Covid-19 pandemic period (March 2019-January 2022) in Turkey. We used self-reported data from participants who all completed questions on socio-demographic and loss-related characteristics plus Hospital Anxiety and Depression Scale (HADS), Prolonged Grief Disorder Scale (PG-13), Multidimensional Scale of Perceived Social Support (MSPSS), Adult Separation Anxiety Questionnaire (ASA-27).

Results: Median age was 34 years, range (18-63); 12 participants were diagnosed with PGD (9.5%). No difference was detected between deaths due to COVID-19 and its complications and deaths due to other causes in terms of PGD diagnosis and PGD symptom severity. When we divide the participants into two groups according to PGD diagnosis (PGD and nonPGD): The average age of the PGD group was higher ($Z=-2.068$; $p=0,31$) and they had more additional medical conditions ($\chi^2=7.21$; $p=0,007$). Thoughts of guilt were more common in the PGD group ($\chi^2=7.92$; $p=0,005$). Additionally, HADS-total, HADS -depression, HADS -anxiety and ASA-27 were higher in the PGD group (respectively: $Z=-4.047$; $P=0,00$, $Z=-4.209$; $P=0,00$, $Z=-3.437$; $P=0,001$, $Z=-1.975$; $P=0,048$). PGD occurred most frequently after first-degree losses ($\chi^2=13.67$; $p=0,00$) and was inversely proportional to the age of the loss ($Z=-1.979$; $P=0,04$). In the nonPGD group, the rate of believing in any religion ($\chi^2=5.807$; $p=0,016$). and the level of fulfilling the requirements of the religion were

higher ($\chi^2=10.584$; $p=0,05$). In the linear regression analysis examining the predictors associated with the severity of prolonged grief; the deceased person was a first-degree relative ($t= 6.23$; $p<0,001$) and younger in age ($t=-3.71$; $p<0,001$), the presence of guilt ($t= 3.28$; $p=0,001$), and increased separation anxiety ($t= 4.13$; $p<0,001$) and depression scores ($t= 4.29$; $p<0,001$) were significant boost of prolonged grief severity.

Conclusions: Although higher PGD rates were expected in deaths due to Covid-19 compared to deaths due to other causes, we did not detect any significant difference in this study. However, this study identified some possible predictors associated with PGD.

Disclosure of Interest: None Declared

O0093

Physical healthcare gap among patients with severe mental illness through the COVID-19 pandemic. Preliminary results from a real-world investigation in Lombardy, Italy

C. Conflitti^{1,2,3*}, M. Monzio Compagnoni^{1,2}, G. Corrao^{2,4} and A. Lora^{2,3}

¹Statistics and Quantitative Methods, University of Milan-Bicocca;

²National Centre of Healthcare Research & Pharmacoepidemiology, Milan;

³Department of Mental Health, ASST Lecco, Lecco and

⁴University of Milan-Bicocca, Milan, Italy

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.216

Introduction: Patients suffering from mental disorders tend to be less adherent to the recommended therapies. Moreover, the COVID-19 pandemic had a global impact on physical and social well-being, which turned out stronger in the most fragile patients, like those with a mental condition.

Objectives: To assess whether the COVID-19 pandemic influenced the physical healthcare gap between patients with and without severe mental illness (SMI) treated for chronic conditions.

Methods: Data were retrieved from Healthcare Utilization Databases of Lombardy region (Italy). Prevalent users of antihypertensive drugs, statins or antidiabetic drugs, receiving healthcare in Lombardy during 2020, were identified. Among them, those with a previous diagnosis of schizophrenic or bipolar disorder were selected and matched with up to 3 patients without any sign of mental disorder by sex, age and number of contacts with the NHS during the previous year. 3 cohorts (not necessarily independent) were formed.

High adherence to specific recommended drug therapies and discontinuation during 2020 were evaluated.

Association between presence of SMI and high adherence was evaluated by using a log-binomial model (risk ratios, RR with 95% CI); a Cox model (hazard ratios, HR) was used for discontinuation.

As comparison, same analyses were performed to the cohorts of prevalent users in 2019, to evaluate the impact of the COVID-19 pandemic. Results were stratified according to the type of mental disorder.

Results: 36'436, 14'136 and 12'597 prevalent users of antihypertensives, statins or antidiabetics respectively were identified, of which 25% with SMI (9'109, 3'536 and 3'152 respectively).